

Linear Codes Constructed From Two Weakly Regular Plateaued Functions with Index $(p - 1)/2$

Shudi Yang^{a,*}, Tonghui Zhang^{b,a}, Zheng-An Yao^c

^a*School of Mathematical Sciences, Qufu Normal University, Shandong 273165, P.R.China*

^b*School of Mathematics and Statistics, Fujian Normal University, Fujian 350117, P.R.China*

^c*School of Mathematics, Sun Yat-Sen University, Guangzhou 510275, P.R.China*

Abstract

Linear codes are the most important family of codes in cryptography and coding theory. Some codes have only a few weights and are widely used in many areas, such as authentication codes, secret sharing schemes and strongly regular graphs. By setting $p \equiv 1 \pmod{4}$, we construct an infinite family of linear codes using two distinct weakly regular unbalanced (and balanced) plateaued functions with index $(p - 1)/2$. Their weight distributions are completely determined by applying exponential sums and Walsh transform. As a result, most of our constructed codes have a few nonzero weights and are minimal.

Keywords: linear code, plateaued function, the weight distribution, Walsh transform.

2010 MSC: 94B15, 11T71

1. Introduction

Throughout the paper, we always let p be an odd prime. We will use the symbol \mathbb{F}_p to denote the finite field with p elements. A linear code C over \mathbb{F}_p with length n , dimension k and minimum distance d is said to have parameters $[n, k, d]$, which means that C is a k -dimensional linear subspace of \mathbb{F}_p^n with Hamming distance d . The code C is called projective if its dual code has minimum

*Corresponding author

Email addresses: yangshudi@qfnu.edu.cn (Shudi Yang), zhangthvvs@126.com (Tonghui Zhang), mcsyao@mail.sysu.edu.cn (Zheng-An Yao)

distance larger than 2. The Hamming weight of a codeword \mathbf{c} , denoted by $\text{wt}(\mathbf{c})$, is defined as the number of nonzero entries in \mathbf{c} . Let $A_w = \#\{\mathbf{c} \in C : \text{wt}(\mathbf{c}) = w\}$ for $0 \leq w \leq n$. Then the sequence $(A_0, A_1, A_2, \dots, A_n)$ stands for the weight distribution of C , where $A_0 = 1$. The code C is called t -weight if the number of nonzero A_w for $1 \leq w \leq n$ equals t . The weight distribution is of vital importance since it contains the information of computing the error probability of error detection and correction. In recent decades, a large number of linear codes have been investigated, most of which have a few weights and good parameters [3, 4, 7, 8, 10, 12, 16, 17, 22, 23, 25, 26]. The construction of linear codes is usually based on different functions, such as, Boolean functions [3], bent functions [19, 26], square functions [20] and weakly regular plateaued functions [4, 16, 17].

Let us introduce an efficient way to construct linear codes, which was proposed by Ding *et al.* [5]. Let $q = p^m$ for an integer m , and D be a subset of \mathbb{F}_q^* of size n . Define

$$C_D = \{\mathbf{c}(a) = (\text{Tr}(ax))_{x \in D} : a \in \mathbb{F}_q\},$$

where Tr is the absolute trace function. It can be checked that C_D is a linear code of length n . The set D is called the defining set of C_D . This approach is soon generalized by Li *et al.* [11], who defined a class of codes by

$$C_D = \left\{ \mathbf{c}(a, b) = (\text{Tr}(ax + by))_{(x, y) \in D} : a, b \in \mathbb{F}_q \right\}, \quad (1.1)$$

where the defining set D is a subset of $\mathbb{F}_q^2 \setminus \{(0, 0)\}$ of size n . Based on this method, Wu *et al.* [21] offered a new approach to linear codes using the defining set

$$D = \{(x, y) \in \mathbb{F}_q^2 \setminus \{(0, 0)\} : f(x) + g(y) = 0\}, \quad (1.2)$$

where f and g are weakly regular bent functions from \mathbb{F}_q to \mathbb{F}_p . Later, Cheng *et al.* in [4] introduced several linear codes C_D of (1.1) with a few weights by considering f and g to be weakly regular unbalanced s -plateaued functions in the defining set (1.2), where $0 \leq s \leq m$. In 2022, Snak [18] went deeper by choosing weakly regular unbalanced and balanced s_f -plateaued function f and

s_g -plateaued function g in the defining set (1.2), where $0 \leq s_f, s_g \leq m$. All of them studied the indexes of functions f and g among the set $\{2, p-1\}$, that is, $l_f, l_g \in \{2, p-1\}$.

Along the research line studied in [4, 18, 21], we further consider the index of $\frac{p-1}{2}$, where $p \equiv 1 \pmod{4}$. Now the defining set is denoted by

$$D_{f,g} = \{(x, y) \in \mathbb{F}_q^2 \setminus \{(0, 0)\} : f(x) + g(y) = 0\}, \quad (1.3)$$

where f and g are weakly regular unbalanced and balanced s -plateaued and t -plateaued functions, respectively, for $0 \leq s, t \leq m$. For clarity, we only concentrate on the case $l_g = \frac{p-1}{2}$ and $l_f \in \{2, \frac{p-1}{2}, p-1\}$, where $p \equiv 1 \pmod{4}$. In this paper, we consider the constructed codes $C_{D_{f,g}}$ of (1.1) and (1.3). In details, we will determine their parameters and their weight distributions using Walsh transform. Their punctured codes are also determined. As we will show later, they are projective, and some of them are optimal since they meet the Griesmer bound.

The rest of this paper is arranged as follows. We firstly present in Section 2 an introduction to the mathematical foundations, including cyclotomic fields and weakly regular plateaued functions. Section 3 gives necessary results for our computation. Our main results are proposed in Section 4, where we study the weight distributions and the parameters of our constructed codes and their punctured ones. Section 5 shows the minimality and applications of these codes. Finally, we conclude the whole paper in Section 6.

2. Mathematical background

In this section, we give a brief exposition of exponential sums, cyclotomic fields, cyclotomic classes and weakly regular plateaued functions. First of all, we set up notation and terminology. Let $q = p^m$ for an integer $m \geq 2$. The set of square (resp. non-square) elements in \mathbb{F}_p^* is denoted by S_q (resp. N_{sq}). Let

η be the quadratic character of \mathbb{F}_p , that is,

$$\eta(x) = \begin{cases} 0, & \text{if } x = 0, \\ 1, & \text{if } x \in S_q, \\ -1, & \text{if } x \in N_{sq}. \end{cases}$$

2.1. Cyclotomic classes and cyclotomic fields

Let θ be a fixed primitive element of \mathbb{F}_q and $N \geq 2$ be a divisor of $q - 1$. The cyclotomic classes of order N in \mathbb{F}_q are defined by $C_i^{(N,q)} = \theta^i \langle \theta^N \rangle$ for $i = 0, 1, \dots, N - 1$, where $\langle \theta^N \rangle$ stands for the subgroup of \mathbb{F}_q^* generated by θ^N . Obviously, we have $\mathbb{F}_q^* = \bigcup_{i=0}^{N-1} C_i^{(N,q)}$ and every cyclotomic class has the same number of elements, that is $\#C_i^{(N,q)} = \frac{q-1}{N}$.

The p -th cyclotomic field is denoted by $K = \mathbb{Q}(\zeta_p)$, where $\zeta_p = \exp\left(\frac{2\pi\sqrt{-1}}{p}\right)$. The following lemma enunciates useful properties of this field.

Lemma 1. ([9]) *The following assertions hold for $K = \mathbb{Q}(\zeta_p)$.*

(1) *The ring of integers in K is $\mathbb{Z}[\zeta_p]$, and $\{\zeta_p^i : 1 \leq i \leq p - 1\}$ is an integer basis of $\mathbb{Z}[\zeta_p]$.*

(2) *The field extension K/\mathbb{Q} is Galois of degree $p - 1$, and the Galois group $\text{Gal}(K/\mathbb{Q}) = \{\sigma_z : z \in \mathbb{F}_p^*\}$, where the automorphism σ_z of K is defined by $\sigma_z(\zeta_p) = \zeta_p^z$.*

(3) *The cyclotomic field K has a unique quadratic subfield $\mathbb{Q}(\sqrt{p^*})$, where $p^* = \eta(-1)p$. For $z \in \mathbb{F}_p^*$, $\sigma_z(\sqrt{p^*}) = \eta(z)\sqrt{p^*}$.*

It is easily proved from Lemma 1 that $\sigma_z(\sqrt{p^{*m}}) = \eta^m(z)\sqrt{p^{*m}}$ and $\sigma_z(\zeta_p^a) = \zeta_p^{za}$ for all $a \in \mathbb{F}_p$.

2.2. Exponential sums

In this subsection, we briefly sketch the concept of exponential sums. Let η_m denote the quadratic character of \mathbb{F}_q , where $q = p^m$. The quadratic Gauss sum over \mathbb{F}_q is defined by

$$G(\eta_m) = \sum_{x \in \mathbb{F}_q^*} \eta_m(x) \chi_1(x),$$

where $\chi_1(x) = \zeta_p^{\text{Tr}(x)}$ is the canonical additive character of \mathbb{F}_q , and Tr is the absolute trace function. From Theorem 5.15 in [13], $G(\eta_m) = (-1)^{m-1} \sqrt{p^*}^m$ and $G(\eta) = \sqrt{p^*}$.

For $n \in \mathbb{N}$ and $a \in \mathbb{F}_q^*$, the Jacobsthal sum is defined by

$$H_n(a) = \sum_{x \in \mathbb{F}_q} \eta_m(x^{n+1} + ax) = \sum_{x \in \mathbb{F}_q} \eta_m(x) \eta_m(x^n + a).$$

Define

$$I_n(a) = \sum_{x \in \mathbb{F}_q} \eta_m(x^n + a).$$

It is a companion sum related to Jacobsthal sums because $I_{2n}(a) = I_n(a) + H_n(a)$, which is due to Theorem 5.50 in [13]. We can evaluate easily that $I_1(a) = 0$ and $I_2(a) = -1$ for all $a \in \mathbb{F}_q^*$. In general, the sums $I_n(a)$ can be described in terms of Jacobi sums.

Lemma 2 (Theorem 5.51, [13]). *For all $a \in \mathbb{F}_q^*$ and $n \in \mathbb{N}$, we have*

$$I_n(a) = \eta_m(a) \sum_{j=1}^{d-1} \lambda^j(-a) J(\lambda^j, \eta_m),$$

where λ is a multiplicative character of \mathbb{F}_q of order $d = \gcd(n, q-1)$, and $J(\lambda^j, \eta_m)$ is a Jacobi sum in \mathbb{F}_q .

Lemma 3 (Theorem 5.33, [13]). *Let $q = p^m$ be odd and $f(x) = a_2x^2 + a_1x + a_0 \in \mathbb{F}_q[x]$ with $a_2 \neq 0$. Then*

$$\sum_{x \in \mathbb{F}_q} \chi_1(f(x)) = \chi_1(a_0 - a_1^2(4a_2)^{-1}) \eta_m(a_2) G(\eta_m).$$

2.3. Weakly regular plateaued functions

We now introduce weakly regular plateaued functions and review some basic facts about them. Let $f : \mathbb{F}_q \rightarrow \mathbb{F}_p$ be a p -ary function. For $\beta \in \mathbb{F}_q$, the Walsh transform of f is defined as a complex-valued function $\widehat{\chi}_f$ on \mathbb{F}_q ,

$$\widehat{\chi}_f(\beta) = \sum_{x \in \mathbb{F}_q} \zeta_p^{f(x) - \text{Tr}(\beta x)}.$$

A p -ary function f is called to be balanced if it satisfies $\widehat{\chi}_f(0) = 0$; otherwise, it is called unbalanced over \mathbb{F}_q .

As a natural extension of bent functions, Zheng *et al.* firstly set up the concept of plateaued functions in characteristic 2 in [24], and later Mesnager [14] gave a general version of any characteristic p . Several years ago, Mesnager *et al.* presented the notion of (non)-weakly regular plateaued functions in their work [15]. We follow the notation used in [15]. For each $\beta \in \mathbb{F}_q$, a function f is called s -plateaued if $|\widehat{\chi}_f(\beta)|^2 \in \{0, p^{m+s}\}$, where $0 \leq s \leq m$. It is worth noting that every bent function is 0-plateaued. The Walsh support of function f , denoted by \mathcal{S}_f , is defined by

$$\mathcal{S}_f = \{\beta \in \mathbb{F}_q : |\widehat{\chi}_f(\beta)|^2 = p^{m+s}\}.$$

Applying the Parseval identity, one gets the absolute Walsh distribution of plateaued functions.

Lemma 4. (*Lemma 1, [14]*) *Let f be an s -plateaued function. Then for $\beta \in \mathbb{F}_q$, $|\widehat{\chi}_f(\beta)|^2$ takes the value p^{m+s} for p^{m-s} times and the value 0 for $p^m - p^{m-s}$ times.*

From Lemma 4, the cardinality of \mathcal{S}_f is given by $\#\mathcal{S}_f = p^{m-s}$.

Definition 1. ([15]) *Let f be an s -plateaued function, where $0 \leq s \leq m$. Then, f is called weakly regular s -plateaued if there exists a complex number u having unit magnitude such that*

$$\widehat{\chi}_f(\beta) \in \{0, up^{\frac{m+s}{2}} \zeta_p^{g(\beta)}\}$$

for all $\beta \in \mathbb{F}_q$, where g is a p -ary function over \mathbb{F}_q satisfying $g(\beta) = 0$ for all $\beta \in \mathbb{F}_q \setminus \mathcal{S}_f$. Otherwise, if u depends on β with $|u| = 1$, then f is called non-weakly regular s -plateaued. Particularly, a weakly regular plateaued function f is said to be regular plateaued if $u = 1$.

Lemma 5. (*Lemma 5, [15]*) *Let $\beta \in \mathbb{F}_q$ and f be a weakly regular s -plateaued function. For every $\beta \in \mathcal{S}_f$ we have*

$$\widehat{\chi}_f(\beta) = \varepsilon_f \sqrt{p^{*m+s}} \zeta_p^{f^*(\beta)},$$

where $\varepsilon_f \in \{\pm 1\}$ is the sign of $\widehat{\chi}_f$ and f^* is a p -ary function over \mathbb{F}_q with $f^*(\beta) = 0$ for all $\beta \in \mathbb{F}_q \setminus \mathcal{S}_f$. We call f^* the dual function of f .

We now introduce two non-trivial subclasses of weakly regular plateaued functions. Let f be a weakly regular unbalanced (resp. balanced) s -plateaued function with $0 \leq s \leq m$. We denote by WRP (resp. WRPB) the subclass of the unbalanced (resp. balanced) functions f that meet the following homogeneous conditions simultaneously:

1. $f(0) = 0$;
2. There exists a positive integer h , such that $2 \mid h$, $\gcd(h-1, p-1) = 1$ and $f(zx) = z^h f(x)$ for any $z \in \mathbb{F}_p^*$ and $x \in \mathbb{F}_q$.

Remark 1. For every $f \in \text{WRP}$ (resp. $f \in \text{WRPB}$), we have $0 \in \mathcal{S}_f$ (resp. $0 \notin \mathcal{S}_f$).

The following lemmas, due to [16] and [18], play a significant role in calculating the parameters of our constructed codes.

Lemma 6. (Lemma 6, [16]) Let $f \in \text{WRP}$ or $f \in \text{WRPB}$ with $\widehat{\chi}_f(\beta) = \varepsilon_f \sqrt{p^{*m+s}} \zeta_p^{f^*(\beta)}$ for every $\beta \in \mathcal{S}_f$. Then, for every $z \in \mathbb{F}_p^*$, $z\beta \in \mathcal{S}_f$ if $\beta \in \mathcal{S}_f$, and otherwise, $z\beta \in \mathbb{F}_q \setminus \mathcal{S}_f$.

Lemma 7. (Propositions 2 and 3, [16]) Let $f \in \text{WRP}$ or $f \in \text{WRPB}$ with $\widehat{\chi}_f(\beta) = \varepsilon_f \sqrt{p^{*m+s}} \zeta_p^{f^*(\beta)}$ for every $\beta \in \mathcal{S}_f$. Then $f^*(0) = 0$ and $f^*(z\beta) = z^{l_f} f^*(\beta)$ for all $z \in \mathbb{F}_p^*$ and $\beta \in \mathcal{S}_f$, where l_f is an even positive integer with $\gcd(l_f - 1, p - 1) = 1$. We call l_f the index of f .

Lemma 8. (Lemma 10, [16]) Let $f \in \text{WRP}$ or $f \in \text{WRPB}$ with $\widehat{\chi}_f(\beta) = \varepsilon_f \sqrt{p^{*m+s}} \zeta_p^{f^*(\beta)}$ for every $\beta \in \mathcal{S}_f$. For $c \in \mathbb{F}_p$, define

$$\mathcal{N}_f(c) = \#\{\beta \in \mathcal{S}_f : f^*(\beta) = c\}.$$

When $m - s$ is even, we have

$$\mathcal{N}_f(c) = \begin{cases} p^{m-s-1} + (p-1)\eta^{m+1}(-1)\varepsilon_f \sqrt{p^{*m-s-2}}, & \text{if } c = 0, \\ p^{m-s-1} - \eta^{m+1}(-1)\varepsilon_f \sqrt{p^{*m-s-2}}, & \text{if } c \in \mathbb{F}_p^*. \end{cases}$$

Otherwise,

$$\mathcal{N}_f(c) = \begin{cases} p^{m-s-1}, & \text{if } c = 0, \\ p^{m-s-1} + \eta(c)\eta^m(-1)\varepsilon_f\sqrt{p^*}^{m-s-1}, & \text{if } c \in \mathbb{F}_p^*. \end{cases}$$

Lemma 9. (Lemma 3.12, [18]) Let $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$, with $\widehat{\chi}_f(\alpha) = \varepsilon_f\sqrt{p^*}^{m+s}\zeta_p^{f^*(\alpha)}$ and $\widehat{\chi}_g(\beta) = \varepsilon_g\sqrt{p^*}^{m+t}\zeta_p^{g^*(\beta)}$ for $\alpha \in \mathcal{S}_f$ and $\beta \in \mathcal{S}_g$, respectively. Define

$$\mathcal{T}(0) = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) + g^*(b) = 0\},$$

$$\mathcal{T}(c) = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) + g^*(b) = c\}, \text{ where } c \in \mathbb{F}_p^*.$$

Then we have

$$\mathcal{T}(0) = \begin{cases} p^{2m-s-t-1} + (p-1)p^{-1}\varepsilon_f\varepsilon_g\sqrt{p^*}^{2m-s-t}, & \text{if } s+t \text{ is even,} \\ p^{2m-s-t-1}, & \text{if } s+t \text{ is odd,} \end{cases}$$

$$\mathcal{T}(c) = \begin{cases} p^{2m-s-t-1} - p^{-1}\varepsilon_f\varepsilon_g\sqrt{p^*}^{2m-s-t}, & \text{if } s+t \text{ is even,} \\ p^{2m-s-t-1} + \eta(c)\varepsilon_f\varepsilon_g\sqrt{p^*}^{2m-s-t-1}, & \text{if } s+t \text{ is odd.} \end{cases}$$

Lemma 10. (Lemma 3.7, [18]) Let $n = \#D_{f,g}$, where $D_{f,g}$ is defined by (1.3) with f and g be as in Lemma 9. If $f, g \in \text{WRPB}$, then $n = p^{2m-1} - 1$. If $f, g \in \text{WRP}$, then

$$n = \begin{cases} p^{2m-1} - 1, & \text{if } 2 \nmid s+t, \\ p^{2m-1} - 1 + (p-1)p^{-1}\varepsilon_f\varepsilon_g\sqrt{p^*}^{2m+s+t}, & \text{if } 2 \mid s+t. \end{cases}$$

3. Auxiliary results

To get the frequencies of codewords in the constructed codes, we will need several lemmas which are depicted and proved in the sequel.

Lemma 11. Let $p \equiv 1 \pmod{2}$. For the quadratic character η over \mathbb{F}_p , we have

$$\sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \eta(u+v) = -\frac{p-1}{2}(\eta(2) + 1),$$

$$\sum_{u \in N_{sq}} \sum_{\substack{v \in N_{sq} \\ v \neq \pm u}} \eta(u+v) = \frac{p-1}{2}(\eta(2)+1).$$

Proof. Notation that $-1 \in S_q$ when $p \equiv 1 \pmod{4}$, and $-1 \in N_{sq}$ when $p \equiv 3 \pmod{4}$. It follows that

$$\begin{aligned} \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \eta(u+v) &= \sum_{u \in S_q} \eta(u) \sum_{\substack{v \in S_q \\ v \neq \pm u}} \eta\left(1 + \frac{v}{u}\right) \\ &= \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm 1}} \eta(1+v) \\ &= \frac{p-1}{2} \left(\sum_{v \in S_q} \eta(1+v) - \eta(2) \right) \\ &= \frac{p-1}{2} \left(\frac{1}{2} \sum_{x \in \mathbb{F}_p} \eta(1+x^2) - \frac{1}{2} - \eta(2) \right) \\ &= \frac{p-1}{2} \left(\frac{1}{2} I_2(1) - \frac{1}{2} - \eta(2) \right). \end{aligned}$$

The first assertion then follows from the fact that $I_2(1) = -1$. The second one is analogously proved and is omitted here. \square

Lemma 12. *Let $p \equiv 1 \pmod{4}$ and $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$, with $\widehat{\chi}_f(\alpha) = \varepsilon_f \sqrt{p}^{m+s} \zeta_p^{f^*(\alpha)}$ and $\widehat{\chi}_g(\beta) = \varepsilon_g \sqrt{p}^{m+t} \zeta_p^{g^*(\beta)}$ for every $\alpha \in \mathcal{S}_f$ and every $\beta \in \mathcal{S}_g$, respectively. Suppose that $s+t$ is odd. Write*

$$\begin{aligned} B_{S_q} &= \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) + g^*(b) \in S_q, f^*(a) - g^*(b) \in S_q\}, \\ B_{N_{sq}} &= \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) + g^*(b) \in N_{sq}, f^*(a) - g^*(b) \in N_{sq}\}. \end{aligned}$$

Then if $2 \nmid m-s$ and $2 \mid m-t$, we have

$$\begin{aligned} B_{S_q} &= \frac{p-1}{2} \sqrt{p}^{2m-s-t-3} \left(\frac{p-1}{2} \sqrt{p}^{2m-s-t-1} - \eta(2) \varepsilon_f \sqrt{p}^{m-t} \right. \\ &\quad \left. + \frac{p+1}{2} \varepsilon_g \sqrt{p}^{m-s-1} + (\eta(2)+p) \varepsilon_f \varepsilon_g \right), \\ B_{N_{sq}} &= \frac{p-1}{2} \sqrt{p}^{2m-s-t-3} \left(\frac{p-1}{2} \sqrt{p}^{2m-s-t-1} + \eta(2) \varepsilon_f \sqrt{p}^{m-t} \right. \\ &\quad \left. + \frac{p+1}{2} \varepsilon_g \sqrt{p}^{m-s-1} - (\eta(2)+p) \varepsilon_f \varepsilon_g \right). \end{aligned}$$

Otherwise if $2 \mid m - s$ and $2 \nmid m - t$, we have

$$\begin{aligned} B_{S_q} &= \frac{p-1}{2} \sqrt{p}^{2m-s-t-3} \left(\frac{p-1}{2} \sqrt{p}^{2m-s-t-1} - \eta(2) \varepsilon_g \sqrt{p}^{m-s} \right. \\ &\quad \left. + \frac{p+1}{2} \varepsilon_f \sqrt{p}^{m-t-1} + (\eta(2) + p) \varepsilon_f \varepsilon_g \right), \\ B_{N_{s_q}} &= \frac{p-1}{2} \sqrt{p}^{2m-s-t-3} \left(\frac{p-1}{2} \sqrt{p}^{2m-s-t-1} + \eta(2) \varepsilon_g \sqrt{p}^{m-s} \right. \\ &\quad \left. + \frac{p+1}{2} \varepsilon_f \sqrt{p}^{m-t-1} - (\eta(2) + p) \varepsilon_f \varepsilon_g \right). \end{aligned}$$

Proof. We only calculate B_{S_q} and omit the other. Now suppose that $2 \nmid m - s$ and $2 \mid m - t$. Let $f^*(a) + g^*(b) = u$, $f^*(a) - g^*(b) = v$, where $u, v \in \mathbb{F}_p^*$. Then we have $f^*(a) = \frac{u+v}{2}$, $g^*(b) = \frac{u-v}{2}$ and so

$$B_{S_q} = \sum_{u \in S_q} \sum_{v \in S_q} \mathcal{N}_f\left(\frac{u+v}{2}\right) \mathcal{N}_g\left(\frac{u-v}{2}\right),$$

where \mathcal{N}_f and \mathcal{N}_g are computed in Lemma 8. It follows that

$$B_{S_q} = \sum_{u \in S_q} \mathcal{N}_f(u) \mathcal{N}_g(0) + \sum_{u \in S_q} \mathcal{N}_f(0) \mathcal{N}_g(u) + S,$$

where

$$S = \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \mathcal{N}_f\left(\frac{u+v}{2}\right) \mathcal{N}_g\left(\frac{u-v}{2}\right). \quad (3.1)$$

Observe that $\frac{u-v}{2} \neq 0$ in (3.1). If we write $c = \frac{u-v}{2} \neq 0$, then from Lemma 8,

$$\begin{aligned} S &= \mathcal{N}_g(c) \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \mathcal{N}_f\left(\frac{u+v}{2}\right) \\ &= \mathcal{N}_g(c) \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \left(p^{m-s-1} + \eta\left(\frac{u+v}{2}\right) \varepsilon_f \sqrt{p}^{m-s-1} \right) \\ &= \mathcal{N}_g(c) \left(\frac{p-1}{2} \cdot \frac{p-5}{2} p^{m-s-1} + \eta(2) \varepsilon_f \sqrt{p}^{m-s-1} \sum_{u \in S_q} \sum_{\substack{v \in S_q \\ v \neq \pm u}} \eta(u+v) \right). \end{aligned}$$

The desired assertion then follows from Lemmas 8 and 11. \square

4. Main results

In this section, we fix $p \equiv 1 \pmod{4}$. Let $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$. For $\alpha, \beta \in \mathbb{F}_q$, we may assume from Lemma 5 that $\widehat{\chi}_f(\alpha) = \varepsilon_f \sqrt{p}^{m+s} \zeta_p^{f^*(\alpha)}$ and

$\widehat{\chi}_g(\beta) = \varepsilon_g \sqrt{p}^{m+t} \zeta_p^{g^*(\beta)}$, where $\alpha \in \mathcal{S}_f$, $\beta \in \mathcal{S}_g$, $\varepsilon_f, \varepsilon_g \in \{\pm 1\}$ and $0 \leq s, t \leq m$. The dual functions f^* and g^* are given in Lemma 7 with $l_f \in \{2, \frac{p-1}{2}, p-1\}$ and $l_g = \frac{p-1}{2}$. For $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$, we define

$$N_0 = \# \{(x, y) \in \mathbb{F}_q^2 : f(x) + g(y) = 0, \text{Tr}(ax + by) = 0\}. \quad (4.1)$$

4.1. The calculation of N_0

To compute the weights of codewords in our codes, it suffices to determine the values of N_0 in (4.1), which are stated in Lemmas 13, 14 and 15.

Lemma 13. *Let $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$ with $l_g = \frac{p-1}{2}$. Suppose that $s+t$ is odd and $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$. For $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, we have $N_0 = p^{2m-2}$, and for $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$, we have the following assertions. When $l_f = \frac{p-1}{2}$, we have*

$$N_0 = \begin{cases} p^{2m-2}, & \text{if } f^*(a) + g^*(b) = 0, \\ p^{2m-2} + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) + g^*(b) \in S_q, \\ p^{2m-2} - (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) + g^*(b) \in N_{sq}. \end{cases}$$

When $l_f = p-1$, we have

$$N_0 = \begin{cases} p^{2m-2} + \frac{p-1}{2} \varepsilon_f \varepsilon_g \eta(2) \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) \in S_q, g^*(b) = \pm f^*(a), \\ p^{2m-2} - \frac{p-1}{2} \varepsilon_f \varepsilon_g \eta(2) \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) \in N_{sq}, g^*(b) = \pm f^*(a), \\ p^{2m-2} + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) + g^*(b) \in S_q, f^*(a) - g^*(b) \in S_q, \\ p^{2m-2} - (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) + g^*(b) \in N_{sq}, f^*(a) - g^*(b) \in N_{sq}, \\ p^{2m-2}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 1 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) = 0, g^*(b) \in S_q \\ & \text{or } g^*(b) = 0, f^*(a) \in S_q, \\ p^{2m-2} - (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) = 0, g^*(b) \in N_{sq} \\ & \text{or } g^*(b) = 0, f^*(a) \in N_{sq}, \\ p^{2m-2} - 2(p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) \in S_q, g^*(b) \in S_q, \\ p^{2m-2} + 2(p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) \in N_{sq}, g^*(b) \in N_{sq}, \\ p^{2m-2}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 5 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) = 0, g^*(b) \in S_q \\ & \text{or } g^*(b) = 0, f^*(a) \in S_q, \\ p^{2m-2} - (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}, & \text{if } f^*(a) = 0, g^*(b) \in N_{sq} \\ & \text{or } g^*(b) = 0, f^*(a) \in N_{sq}, \\ p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-3}\eta(f^*(a))\left(I_4\left(\frac{g^*(b)}{f^*(a)}\right) - \eta\left(\frac{g^*(b)}{f^*(a)}\right)\right), & \text{otherwise,} \end{cases}$$

where I_4 is a companion sum determined in Lemma 2.

Proof. Let $2 \nmid s+t$. By definition in (4.1) and the orthogonal property of group characters,

$$\begin{aligned} N_0 &= \frac{1}{p^2} \sum_{x,y \in \mathbb{F}_q} \sum_{z \in \mathbb{F}_p} \zeta_p^{z(f(x)+g(y))} \sum_{h \in \mathbb{F}_p} \zeta_p^{h\text{Tr}(ax+by)} \\ &= \frac{1}{p^2} \sum_{x,y \in \mathbb{F}_q} \left(1 + \sum_{z \in \mathbb{F}_p^*} \zeta_p^{z(f(x)+g(y))}\right) \left(1 + \sum_{h \in \mathbb{F}_p^*} \zeta_p^{h\text{Tr}(ax+by)}\right) \\ &= p^{2m-2} + \frac{1}{p^2} \sum_{z \in \mathbb{F}_p^*} \sum_{x,y \in \mathbb{F}_q} \zeta_p^{z(f(x)+g(y))} \\ &\quad + \frac{1}{p^2} \sum_{x,y \in \mathbb{F}_q} \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \zeta_p^{z(f(x)+g(y))+h\text{Tr}(ax+by)} \\ &= p^{2m-2} + p^{-2}(\Delta_1 + \Delta_2), \end{aligned} \tag{4.2}$$

where we write

$$\begin{aligned}\Delta_1 &= \sum_{z \in \mathbb{F}_p^*} \sum_{x, y \in \mathbb{F}_q} \zeta_p^{z(f(x)+g(y))}, \\ \Delta_2 &= \sum_{x, y \in \mathbb{F}_q} \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \zeta_p^{z(f(x)+g(y))+h\text{Tr}(ax+by)}.\end{aligned}$$

It follows that

$$\begin{aligned}\Delta_1 &= \sum_{z \in \mathbb{F}_p^*} \sigma_z(\widehat{\chi}_f(0)\widehat{\chi}_g(0)) \\ &= \begin{cases} 0, & \text{if } f, g \in \text{WRPB}, \\ \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta^{s+t}(z), & \text{if } f, g \in \text{WRP}. \end{cases}\end{aligned}$$

So we always have $\Delta_1 = 0$ when $2 \nmid s+t$. Now it is sufficient to determine Δ_2 .

We observe from its definition that

$$\begin{aligned}\Delta_2 &= \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_q} \zeta_p^{zf(x)-\text{Tr}(hax)} \sum_{y \in \mathbb{F}_q} \zeta_p^{zg(y)-\text{Tr}(hby)} \\ &= \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_q} \zeta_p^{z(f(x)-\text{Tr}(\frac{h}{z}ax))} \sum_{y \in \mathbb{F}_q} \zeta_p^{z(g(y)-\text{Tr}(\frac{h}{z}by))} \\ &= \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \sigma_z(\widehat{\chi}_f(ha)\widehat{\chi}_g(hb)).\end{aligned}\tag{4.3}$$

Obviously, when $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, then from Lemma 6 $(ha, hb) \notin \mathcal{S}_f \times \mathcal{S}_g$ for every $h \in \mathbb{F}_p^*$. Hence $\widehat{\chi}_f(ha) = 0$ or $\widehat{\chi}_g(hb) = 0$, and consequently by (4.3)

$$\Delta_2 = 0.$$

When $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$, then $(ha, hb) \in \mathcal{S}_f \times \mathcal{S}_g$ for every $h \in \mathbb{F}_p^*$. By (4.3), Lemmas 1, 5 and 7, we obtain

$$\begin{aligned}\Delta_2 &= \sum_{z \in \mathbb{F}_p^*} \sigma_z \left(\sum_{h \in \mathbb{F}_p^*} \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \zeta_p^{h^l f^*(a) + h^l g^*(b)} \right) \\ &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta^{s+t}(z) \sigma_z \left(\sum_{h \in \mathbb{F}_p^*} \zeta_p^{h^l f^*(a) + h^l g^*(b)} \right).\end{aligned}\tag{4.4}$$

In the following, we will apply Lemmas 1 and 3 to determine Δ_2 in (4.4) by considering the cases of $l_f = 2$, $l_f = \frac{p-1}{2}$ and $l_f = p-1$, separately.

(1) The first case we consider is $l_f = \frac{p-1}{2}$.

For $h \in \mathbb{F}_p^*$, clearly $h^{\frac{p-1}{2}} = 1$ if $h \in S_q$, and -1 otherwise. So we have from (4.4) that

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{f^*(a)+g^*(b)} + \sum_{h \in N_{sq}} \zeta_p^{-f^*(a)-g^*(b)} \right) \\
&= \frac{p-1}{2} \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \left(\sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{z(f^*(a)+g^*(b))} + \sum_{z \in \mathbb{F}_p^*} \eta(-z) \zeta_p^{-z(f^*(a)+g^*(b))} \right) \\
&= (p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{z(f^*(a)+g^*(b))} \\
&= \begin{cases} 0, & \text{if } f^*(a) + g^*(b) = 0, \\ (p-1) \varepsilon_f \varepsilon_g \eta(f^*(a) + g^*(b)) \sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) + g^*(b) \neq 0. \end{cases}
\end{aligned}$$

(2) The second case is that $l_f = p-1$.

In this case, $h^{p-1} = 1$ for every $h \in \mathbb{F}_p^*$. By (4.4), we have

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{f^*(a)+g^*(b)} + \sum_{h \in N_{sq}} \zeta_p^{f^*(a)-g^*(b)} \right) \\
&= \frac{p-1}{2} \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \left(\sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{z(f^*(a)+g^*(b))} + \sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{z(f^*(a)-g^*(b))} \right) \\
&= \begin{cases} 0, & \text{if } f^*(a) = g^*(b) = 0, \\ \frac{p-1}{2} \varepsilon_f \varepsilon_g \eta(2f^*(a)) \sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) \neq 0, g^*(b) = -f^*(a), \\ \frac{p-1}{2} \varepsilon_f \varepsilon_g \eta(2f^*(a)) \sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) \neq 0, g^*(b) = f^*(a), \\ \frac{p-1}{2} \varepsilon_f \varepsilon_g \left(\eta(f^*(a) + g^*(b)) + \eta(f^*(a) - g^*(b)) \right) \sqrt{p}^{2m+s+t+1}, & \text{otherwise.} \end{cases}
\end{aligned}$$

(3) The last case is that $l_f = 2$ and it is distinguished between two subcases.

Subcase (a): If $p \equiv 1 \pmod{8}$, then $-1 \in C_0^{(4,p)}$. So from (4.4),

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{h^2 f^*(a)+g^*(b)} + \sum_{h \in N_{sq}} \zeta_p^{h^2 f^*(a)-g^*(b)} \right) \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{h^2 f^*(a)+g^*(b)} + \sum_{h \in N_{sq}} \zeta_p^{-(h^2 f^*(a)+g^*(b))} \right) \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \left(\sum_{z \in \mathbb{F}_p^*} \eta(z) \sum_{h \in S_q} \zeta_p^{z(h^2 f^*(a)+g^*(b))} + \sum_{z \in \mathbb{F}_p^*} \eta(-z) \sum_{h \in N_{sq}} \zeta_p^{-z(h^2 f^*(a)+g^*(b))} \right).
\end{aligned}$$

Replacing $-z$ by z in the last double sum above, we obtain

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sum_{h \in \mathbb{F}_p^*} \zeta_p^{z(h^2 f^*(a) + g^*(b))} \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{zg^*(b)} \sum_{h \in \mathbb{F}_p^*} \zeta_p^{zh^2 f^*(a)} \\
&= \begin{cases} 0, & \text{if } f^*(a) = g^*(b) = 0, \\ (p-1)\varepsilon_f \varepsilon_g \eta(g^*(b)) \sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) = 0, g^*(b) \neq 0, \\ (p-1)\varepsilon_f \varepsilon_g \eta(f^*(a)) \sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) \neq 0, g^*(b) = 0, \\ -(p-1)\varepsilon_f \varepsilon_g (\eta(f^*(a)) + \eta(g^*(b))) \sqrt{p}^{2m+s+t+1}, & \text{otherwise.} \end{cases}
\end{aligned}$$

Subcase (b): If $p \equiv 5 \pmod{8}$, then $-1 \in C_2^{(4,p)}$. So from (4.4),

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{h^2 f^*(a) + g^*(b)} + \sum_{h \in N_{sq}} \zeta_p^{h^2 f^*(a) - g^*(b)} \right) \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{h^2 f^*(a) + g^*(b)} + \sum_{h \in S_q} \zeta_p^{-(h^2 f^*(a) + g^*(b))} \right) \\
&= 2\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \eta(z) \sigma_z \left(\sum_{h \in S_q} \zeta_p^{h^2 f^*(a) + g^*(b)} \right) \\
&= 2\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{h \in S_q} \sum_{z \in \mathbb{F}_p^*} \eta(z) \zeta_p^{zh^2 f^*(a) + g^*(b)}.
\end{aligned}$$

Assume that $f^*(a)g^*(b) \neq 0$. If $\frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}$, then the equation $h^2 f^*(a) + g^*(b) = 0$ has exactly two solutions h_1, h_2 in S_q , where $h_2 = -h_1$. Otherwise if $\frac{g^*(b)}{f^*(a)} \notin C_2^{(4,p)}$, then the inequality $h^2 f^*(a) + g^*(b) \neq 0$ holds for all h in S_q . Consequently, if $f^*(a)g^*(b) \neq 0$, then

$$\begin{aligned}
\Delta_2 &= 2\varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t+1} \sum_{h \in S_q} \eta(h^2 f^*(a) + g^*(b)) \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t+1} \sum_{h \in \mathbb{F}_p^*} \eta(h^4 f^*(a) + g^*(b)) \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t+1} \eta(f^*(a)) \left(I_4 \left(\frac{g^*(b)}{f^*(a)} \right) - \eta \left(\frac{g^*(b)}{f^*(a)} \right) \right),
\end{aligned}$$

where I_4 is determined from Lemma 2. Thus we conclude that

$$\Delta_2 = \begin{cases} 0, & \text{if } f^*(a) = g^*(b) = 0, \\ (p-1)\varepsilon_f\varepsilon_g\eta(g^*(b))\sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) = 0, g^*(b) \neq 0, \\ (p-1)\varepsilon_f\varepsilon_g\eta(f^*(a))\sqrt{p}^{2m+s+t+1}, & \text{if } f^*(a) \neq 0, g^*(b) = 0, \\ \varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t+1}\eta(f^*(a))\left(I_4\left(\frac{g^*(b)}{f^*(a)}\right) - \eta\left(\frac{g^*(b)}{f^*(a)}\right)\right), & \text{otherwise.} \end{cases}$$

The desired conclusion then follows from (4.2), completing the proof. \square

Lemma 14. *Let $f, g \in \text{WRP}$ with $l_g = \frac{p-1}{2}$. Suppose that $s+t$ is even and $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$. For $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, we have $N_0 = p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-4}$, and for $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$, we have the following assertions. When $l_f = \frac{p-1}{2}$, we have*

$$N_0 = \begin{cases} p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a) + g^*(b) = 0, \\ p^{2m-2}, & \text{if } f^*(a) + g^*(b) \neq 0. \end{cases}$$

When $l_f = p-1$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + \frac{p-1}{2}\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a) \neq 0, g^*(b) = -f^*(a) \\ & \text{or } f^*(a) \neq 0, g^*(b) = f^*(a), \\ p^{2m-2}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 1 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + 2\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a)g^*(b) \in S_q, \\ p^{2m-2}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 5 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + 4\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t-2}, & \text{if } \frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}, \\ p^{2m-2}, & \text{otherwise.} \end{cases}$$

Proof. The proof is completed in a manner analogous to the previous lemma by noting that $2 \mid s + t$. From (4.2) and (4.3),

$$N_0 = p^{2m-2} + p^{-2}(\Delta_1 + \Delta_2),$$

where

$$\begin{aligned} \Delta_1 &= (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, \\ \Delta_2 &= \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \sigma_z(\widehat{\chi}_f(ha)\widehat{\chi}_g(hb)). \end{aligned}$$

We always have $\Delta_2 = 0$ unless $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$. Now we let $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$. Then the value of Δ_2 in (4.4) is determined by distinguishing the cases of $l_f = 2$, $l_f = \frac{p-1}{2}$ and $l_f = p-1$, respectively.

(1) The first case we consider is $l_f = \frac{p-1}{2}$.

It follows from (4.4) that

$$\begin{aligned} \Delta_2 &= (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \zeta_p^{z(f^*(a)+g^*(b))} \\ &= \begin{cases} (p-1)^2\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, & \text{if } f^*(a) + g^*(b) = 0, \\ -(p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, & \text{if } f^*(a) + g^*(b) \neq 0. \end{cases} \end{aligned}$$

(2) The second case is that $l_f = p-1$.

Again from (4.4), we have

$$\begin{aligned} \Delta_2 &= \frac{p-1}{2}\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t} \left(\sum_{z \in \mathbb{F}_p^*} \zeta_p^{z(f^*(a)+g^*(b))} + \sum_{z \in \mathbb{F}_p^*} \zeta_p^{z(f^*(a)-g^*(b))} \right) \\ &= \begin{cases} (p-1)^2\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, & \text{if } f^*(a) = g^*(b) = 0, \\ \frac{p-1}{2}(p-2)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, & \text{if } f^*(a) \neq 0, g^*(b) = -f^*(a) \\ & \text{or } f^*(a) \neq 0, g^*(b) = f^*(a), \\ -(p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{2m+s+t}, & \text{otherwise.} \end{cases} \end{aligned}$$

(3) The last case is that $l_f = 2$ and we need only consider two different subcases.

Subcase (a): If $p \equiv 1 \pmod{8}$, then from (4.4),

$$\begin{aligned}
\Delta_2 &= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \sum_{h \in \mathbb{F}_p^*} \zeta_p^{z(h^2 f^*(a) + g^*(b))} \\
&= \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{z \in \mathbb{F}_p^*} \zeta_p^{z g^*(b)} \sum_{h \in \mathbb{F}_p^*} \zeta_p^{z h^2 f^*(a)} \\
&= \begin{cases} (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{if } f^*(a) = g^*(b) = 0, \\ (p+1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{if } f^*(a) g^*(b) \in S_q, \\ -(p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{otherwise.} \end{cases}
\end{aligned}$$

Subcase (b): If $p \equiv 5 \pmod{8}$, then from (4.4),

$$\Delta_2 = 2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \sum_{h \in S_q} \sum_{z \in \mathbb{F}_p^*} \zeta_p^{z(h^2 f^*(a) + g^*(b))}.$$

The value of Δ_2 is clear if $f^*(a)g^*(b) = 0$. We now assume that $f^*(a)g^*(b) \neq 0$. If $\frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}$, then the equation $h^2 f^*(a) + g^*(b) = 0$ has exactly two solutions h_1, h_2 in S_q , where $h_2 = -h_1$. Hence,

$$\begin{aligned}
\Delta_2 &= 2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \left(2(p-1) - \left(\frac{p-1}{2} - 2 \right) \right) \\
&= (3p+1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}.
\end{aligned}$$

Otherwise if $\frac{g^*(b)}{f^*(a)} \notin C_2^{(4,p)}$, then the inequality $h^2 f^*(a) + g^*(b) \neq 0$ holds for all h in S_q . Thus

$$\begin{aligned}
\Delta_2 &= 2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t} \times \frac{p-1}{2} \times (-1) \\
&= -(p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}.
\end{aligned}$$

So we conclude that

$$\Delta_2 = \begin{cases} (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{if } f^*(a) = g^*(b) = 0, \\ (3p+1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{if } \frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}, \\ -(p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t}, & \text{otherwise.} \end{cases}$$

The desired conclusion then follows from (4.2), completing the proof. \square

Lemma 15. *Let $f, g \in \text{WRPB}$ with $l_g = \frac{p-1}{2}$. Suppose that $s+t$ is even and $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$. For $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, we have $N_0 = p^{2m-2}$, and for $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$, we have the following assertions. When $l_f = \frac{p-1}{2}$, we have*

$$N_0 = \begin{cases} p^{2m-2} + (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) + g^*(b) = 0, \\ p^{2m-2} - (p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) + g^*(b) \neq 0. \end{cases}$$

When $l_f = p-1$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + \frac{p-1}{2} (p-2) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) \neq 0, g^*(b) = -f^*(a) \\ & \text{or } f^*(a) \neq 0, g^*(b) = f^*(a), \\ p^{2m-2} - (p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 1 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + (p+1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a)g^*(b) \in S_q, \\ p^{2m-2} - (p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{otherwise.} \end{cases}$$

When $l_f = 2$ and $p \equiv 5 \pmod{8}$, we have

$$N_0 = \begin{cases} p^{2m-2} + (p-1)^2 \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } f^*(a) = g^*(b) = 0, \\ p^{2m-2} + (3p+1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{if } \frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}, \\ p^{2m-2} - (p-1) \varepsilon_f \varepsilon_g \sqrt{p}^{2m+s+t-4}, & \text{otherwise.} \end{cases}$$

Proof. Notation that $\Delta_1 = \sum_{z \in \mathbb{F}_p^*} \sigma_z(\widehat{\chi}_f(0)\widehat{\chi}_g(0)) = 0$ for $f, g \in \text{WRPB}$. From (4.2), $N_0 = p^{2m-2} + p^{-2}(\Delta_1 + \Delta_2) = p^{2m-2} + p^{-2}\Delta_2$, where Δ_2 is determined in Lemma 14. This completes the proof. \square

4.2. Weight distributions of $C_{D_{f,g}}$ from two weakly regular plateaued functions

In this subsection, we will deal with the weight distributions of $C_{D_{f,g}}$ defined by (1.1) and (1.3). Its length is denoted by n and is given in Lemma 10. We will show the main results in the following theorems explicitly. For abbreviation, we write $\tau = 2m + s + t$ and $\gamma = 2m - s - t$.

Theorem 1. Suppose that $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$ with $l_g = \frac{p-1}{2}$. Let $s + t$ be odd. If $l_f = \frac{p-1}{2}$, then $C_{D_{f,g}}$ is a three-weight $[p^{2m-1} - 1, 2m]$ linear code with its weight distribution listed in Table 1. If $l_f = p - 1$, then $C_{D_{f,g}}$ is a five-weight $[p^{2m-1} - 1, 2m]$ linear code with its weight distribution listed in Table 2. If $l_f = 2$ and $p \equiv 1 \pmod{8}$, then $C_{D_{f,g}}$ is a five-weight $[p^{2m-1} - 1, 2m]$ linear code with its weight distribution listed in Table 3. If $l_f = 2$ and $p \equiv 5 \pmod{8}$, then $C_{D_{f,g}}$ is a $[p^{2m-1} - 1, 2m]$ linear code with its weight distribution listed in Table 4.

Proof. From Lemma 10, the length is $n = p^{2m-1} - 1$. Let $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$, the weight of nonzero codewords $\mathbf{c}(a, b)$ is denoted by $\text{wt}(\mathbf{c}(a, b))$. It follows that

$$\text{wt}(\mathbf{c}(a, b)) = n + 1 - N_0,$$

where N_0 is given by Lemma 13. To be more precise, for each $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, we have

$$\text{wt}(\mathbf{c}(a, b)) = (p - 1)p^{2m-2}.$$

For each $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g$, there are four different cases when $\text{wt}(\mathbf{c}(a, b)) \neq (p - 1)p^{2m-2}$.

(1) When $l_f = \frac{p-1}{2}$, we have

$$\text{wt}(\mathbf{c}(a, b)) = \begin{cases} (p - 1)(p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & \frac{p-1}{2} \mathcal{T}(i) \text{ times,} \\ (p - 1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & \frac{p-1}{2} \mathcal{T}(j) \text{ times,} \end{cases}$$

where $\mathcal{T}(i)$ and $\mathcal{T}(j)$ are computed in Lemma 9 for $i \in S_q$ and $j \in N_{sq}$. This leads to the weight distribution in Table 1.

(2) When $l_f = p - 1$, we have

$$\text{wt}(\mathbf{c}(a, b)) = \begin{cases} (p - 1)(p^{2m-2} - \frac{1}{2} \varepsilon_f \varepsilon_g \eta(2) \sqrt{p}^{\tau-3}), & E_1 \text{ times,} \\ (p - 1)(p^{2m-2} + \frac{1}{2} \varepsilon_f \varepsilon_g \eta(2) \sqrt{p}^{\tau-3}), & E_2 \text{ times,} \\ (p - 1)(p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & B_{S_q} \text{ times,} \\ (p - 1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & B_{N_{sq}} \text{ times,} \end{cases}$$

where the numbers B_{S_q} and $B_{N_{sq}}$ are computed in Lemma 12, and

$$E_1 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) \in S_q, g^*(b) = \pm f^*(a)\} = (p-1)\mathcal{N}_f(i)\mathcal{N}_g(i),$$

$$E_2 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) \in N_{sq}, g^*(b) = \pm f^*(a)\} = (p-1)\mathcal{N}_f(j)\mathcal{N}_g(j),$$

for $i \in S_q$, $j \in N_{sq}$, and \mathcal{N}_f and \mathcal{N}_g are given in Lemma 8. The weight distribution in Table 2 is then established.

(3) When $l_f = 2$ and $p \equiv 1 \pmod{8}$, we have

$$\text{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)(p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & E_3 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & E_4 \text{ times,} \\ (p-1)(p^{2m-2} + 2\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & \frac{(p-1)^2}{4} \mathcal{N}_f(i)\mathcal{N}_g(i) \text{ times,} \\ (p-1)(p^{2m-2} - 2\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & \frac{(p-1)^2}{4} \mathcal{N}_f(j)\mathcal{N}_g(j) \text{ times,} \end{cases}$$

where

$$E_3 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) = 0, g^*(b) \in S_q \text{ or } g^*(b) = 0, f^*(a) \in S_q\}$$

$$= \frac{p-1}{2}(\mathcal{N}_f(0)\mathcal{N}_g(i) + \mathcal{N}_f(i)\mathcal{N}_g(0)),$$

$$E_4 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) = 0, g^*(b) \in N_{sq} \text{ or } g^*(b) = 0, f^*(a) \in N_{sq}\}$$

$$= \frac{p-1}{2}(\mathcal{N}_f(0)\mathcal{N}_g(j) + \mathcal{N}_f(j)\mathcal{N}_g(0)),$$

for $i \in S_q$ and $j \in N_{sq}$. Thus we get the weight distribution listed in Table 3.

(4) When $l_f = 2$ and $p \equiv 5 \pmod{8}$, we have

$$\text{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)(p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & E_3 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}), & E_4 \text{ times,} \\ (p-1)p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3} \eta(u) \left(I_4\left(\frac{v}{u}\right) - \eta\left(\frac{v}{u}\right) \right) & \mathcal{N}_f(u)\mathcal{N}_g(v) \text{ times.} \\ & \text{for all } u, v \in \mathbb{F}_p^*, \end{cases}$$

The weight distribution of this case is summarized in Table 4. \square

Theorem 2. *Suppose that $f, g \in \text{WRP}$ with $l_g = \frac{p-1}{2}$. Let $s + t$ be even. If $l_f = \frac{p-1}{2}$, then $C_{D_{f,g}}$ is a three-weight $[n, 2m]$ linear code with its weight*

Table 1: The weight distribution of $C_{D_{f,g}}$ in Theorem 1 when $l_f = \frac{p-1}{2}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$p^{2m} - (p-1)p^{\gamma-1} - 1$
$(p-1)(p^{2m-2} - \sqrt{p}^{\tau-3})$	$\frac{p-1}{2}(p^{\gamma-1} + \sqrt{p}^{\gamma-1})$
$(p-1)(p^{2m-2} + \sqrt{p}^{\tau-3})$	$\frac{p-1}{2}(p^{\gamma-1} - \sqrt{p}^{\gamma-1})$

Table 2: The weight distribution of $C_{D_{f,g}}$ in Theorem 1 when $l_f = p-1$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$p^{2m} - 1 - E_1 - E_2 - B_{S_q} - B_{N_{sq}}$
$(p-1)(p^{2m-2} - \frac{1}{2}\varepsilon_f\varepsilon_g\eta(2)\sqrt{p}^{\tau-3})$	E_1
$(p-1)(p^{2m-2} + \frac{1}{2}\varepsilon_f\varepsilon_g\eta(2)\sqrt{p}^{\tau-3})$	E_2
$(p-1)(p^{2m-2} - \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	B_{S_q}
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	$B_{N_{sq}}$

Table 3: The weight distribution of $C_{D_{f,g}}$ in Theorem 1 when $l_f = 2$ and $p \equiv 1 \pmod{8}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$p^{2m} - 1 - E_3 - E_4 - \frac{(p-1)^2}{4}(\mathcal{N}_f(i)\mathcal{N}_g(i) + \mathcal{N}_f(j)\mathcal{N}_g(j))$
$(p-1)(p^{2m-2} - \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	E_3
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	E_4
$(p-1)(p^{2m-2} + 2\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	$\frac{(p-1)^2}{4}\mathcal{N}_f(i)\mathcal{N}_g(i)$
$(p-1)(p^{2m-2} - 2\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-3})$	$\frac{(p-1)^2}{4}\mathcal{N}_f(j)\mathcal{N}_g(j)$

Table 4: The weight distribution of $C_{D_{f,g}}$ in Theorem 1 when $l_f = 2$ and $p \equiv 5 \pmod{8}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$p^{2m} - 1 - E_3 - E_4 - \sum_{u,v \in \mathbb{F}_p^*} \mathcal{N}_f(u)\mathcal{N}_g(v)$
$(p-1)(p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3})$	E_3
$(p-1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3})$	E_4
$(p-1)p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3} \eta(u) \left(I_4\left(\frac{v}{u}\right) - \eta\left(\frac{v}{u}\right) \right)$ for all $u, v \in \mathbb{F}_p^*$	$\mathcal{N}_f(u)\mathcal{N}_g(v)$

distribution listed in Table 5. If $l_f = p-1$, then $C_{D_{f,g}}$ is a four-weight $[n, 2m]$ linear code with its weight distribution listed in Table 6. Otherwise if $l_f = 2$, then $C_{D_{f,g}}$ is a four-weight $[n, 2m]$ linear code with its weight distribution listed in Table 7 when $p \equiv 1 \pmod{8}$, and in Table 8 when $p \equiv 5 \pmod{8}$. Here we set $n = p^{2m-1} - 1 + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-2}$ for brevity.

Proof. The length of the code $C_{D_{f,g}}$ comes from Lemma 10. For $(a, b) \in \mathbb{F}_q^2 \setminus \{(0, 0)\}$, the weight $\mathbf{wt}(\mathbf{c}(a, b)) = n + 1 - N_0$ can be obtained from Lemma 14. To be more explicit, when $(a, b) \notin \mathcal{S}_f \times \mathcal{S}_g$, we have

$$\mathbf{wt}(\mathbf{c}(a, b)) = (p-1)(p^{2m-2} + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-4}).$$

By Lemma 4, the frequency of such codewords equals $p^{2m} - p^\gamma$ since $f, g \in \text{WRP}$. When $(a, b) \in \mathcal{S}_f \times \mathcal{S}_g \setminus \{(0, 0)\}$, we will discuss the following four different cases.

(1) When $l_f = \frac{p-1}{2}$, we have

$$\mathbf{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)p^{2m-2}, & \mathcal{T}(0) - 1 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-2}), & (p-1)\mathcal{T}(c) \text{ times,} \end{cases}$$

where $\mathcal{T}(0)$ and $\mathcal{T}(c)$ are given in Lemma 9 for $c \neq 0$. This gives the weight distribution in Table 5.

(2) When $l_f = p - 1$, we have

$$\mathbf{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)p^{2m-2}, & \mathcal{N}_f(0)\mathcal{N}_g(0) - 1 \text{ times,} \\ (p-1)(p^{2m-2} + \frac{1}{2}\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}), & F_1 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}), & F_2 \text{ times,} \end{cases}$$

where we define

$$F_1 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a) \neq 0, g^*(b) = \pm f^*(a)\} = 2 \sum_{c \in \mathbb{F}_p^*} \mathcal{N}_f(c)\mathcal{N}_g(c),$$

$$F_2 = p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_1.$$

Thus we obtain the weight distribution in Table 6.

(3) When $l_f = 2$ and $p \equiv 1 \pmod{8}$, we have

$$\mathbf{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)p^{2m-2}, & \mathcal{N}_f(0)\mathcal{N}_g(0) - 1 \text{ times,} \\ (p-1)p^{2m-2} + (p-3)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}, & F_3 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}), & F_4 \text{ times,} \end{cases}$$

where

$$F_3 = \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : f^*(a)g^*(b) \in S_q\} = \frac{(p-1)^2}{4}(\mathcal{N}_f(i)\mathcal{N}_g(i) + \mathcal{N}_f(j)\mathcal{N}_g(j)),$$

$$F_4 = p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_3,$$

for $i \in S_q$ and $j \in N_{sq}$. This implies the weight distribution listed in Table 7.

(4) When $l_f = 2$ and $p \equiv 5 \pmod{8}$, we get

$$\mathbf{wt}(\mathbf{c}(a, b)) = \begin{cases} (p-1)p^{2m-2}, & \mathcal{N}_f(0)\mathcal{N}_g(0) - 1 \text{ times,} \\ (p-1)p^{2m-2} + (p-5)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}, & F_5 \text{ times,} \\ (p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}), & F_6 \text{ times,} \end{cases}$$

where we write

$$\begin{aligned} F_5 &= \#\{(a, b) \in \mathcal{S}_f \times \mathcal{S}_g : \frac{g^*(b)}{f^*(a)} \in C_2^{(4,p)}\} \\ &= \frac{(p-1)^2}{8}(\mathcal{N}_f(i)\mathcal{N}_g(i) + \mathcal{N}_f(j)\mathcal{N}_g(j)) = \frac{1}{2}F_3, \end{aligned}$$

$$F_6 = p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - \frac{1}{2}F_3,$$

for $i \in S_q$ and $j \in N_{sq}$. Thus the result in Table 8 is derived. \square

Table 5: The weight distribution of $C_{D_{f,g}}$ in Theorem 2 when $l_f = \frac{p-1}{2}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$p^{\gamma-1} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2} - 1$
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2})$	$(p-1)(p^{\gamma-1} - \varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2})$
$(p-1)(p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^{2m} - p^\gamma$

Table 6: The weight distribution of $C_{D_{f,g}}$ in Theorem 2 when $l_f = p-1$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$\mathcal{N}_f(0)\mathcal{N}_g(0) - 1$
$(p-1)(p^{2m-2} + \frac{1}{2}\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2})$	F_1
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_1$
$(p-1)(p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^{2m} - p^\gamma$

Table 7: The weight distribution of $C_{D_{f,g}}$ in Theorem 2 when $l_f = 2$ and $p \equiv 1 \pmod{8}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$\mathcal{N}_f(0)\mathcal{N}_g(0) - 1$
$(p-1)p^{2m-2} + (p-3)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}$	F_3
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_3$
$(p-1)(p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^{2m} - p^\gamma$

Theorem 3. Suppose that $f, g \in \text{WRPB}$ with $l_g = \frac{p-1}{2}$. Let $s+t$ be even. If $l_f = \frac{p-1}{2}$, then $C_{D_{f,g}}$ is a three-weight $[p^{2m-1}-1, 2m]$ linear code with its weight distribution listed in Table 9. If $l_f = p-1$, then $C_{D_{f,g}}$ is a four-weight $[p^{2m-1}-$

Table 8: The weight distribution of $C_{D_{f,g}}$ in Theorem 2 when $l_f = 2$ and $p \equiv 5 \pmod{8}$.

weight	frequency
0	1
$(p-1)p^{2m-2}$	$\mathcal{N}_f(0)\mathcal{N}_g(0) - 1$
$(p-1)p^{2m-2} + (p-5)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2}$	$\frac{1}{2}F_3$
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-2})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - \frac{1}{2}F_3$
$(p-1)(p^{2m-2} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^{2m} - p^\gamma$

$1, 2m]$ linear code with its weight distribution listed in Table 10. Otherwise if $l_f = 2$, then $C_{D_{f,g}}$ is a four-weight $[p^{2m-1} - 1, 2m]$ linear code with its weight distribution listed in Table 11 when $p \equiv 1 \pmod{8}$, and in Table 12 when $p \equiv 5 \pmod{8}$.

Proof. Notation that $(0, 0)$ is not in $\mathcal{S}_f \times \mathcal{S}_g$ since $f, g \in \text{WRPB}$. This theorem can be derived in the same way as Theorem 2 by using Lemmas 4, 8, 9, 10 and 15. We omit the details here. \square

Table 9: The weight distribution of $C_{D_{f,g}}$ in Theorem 3 when $l_f = \frac{p-1}{2}$.

weight	frequency
0	1
$(p-1)(p^{2m-2} - (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^{\gamma-1} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2}$
$(p-1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$(p-1)(p^{\gamma-1} - \varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2})$
$(p-1)p^{2m-2}$	$p^{2m} - p^\gamma - 1$

Remark 2. In Theorems 1, 2 and 3, we dealt with the code $C_{D_{f,g}}$ for $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$ with $l_f \in \{2, \frac{p-1}{2}, p-1\}$ and $l_g = \frac{p-1}{2}$, where $p \equiv 1 \pmod{4}$. However, some of the results coincide with the known ones in the literature. Specifically, when $f, g \in \text{WRP}$, the weight distributions in Tables 1, 5 and 7 coincide with the results of Tables 3, 4 and 5 in [18], respectively. If we set $t = s$ in Tables 5 and 7, then we get the results of Theorem 4 in [4].

Table 10: The weight distribution of $C_{D_{f,g}}$ in Theorem 3 when $l_f = p - 1$.

weight	frequency
0	1
$(p - 1)(p^{2m-2} - (p - 1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$\mathcal{N}_f(0)\mathcal{N}_g(0)$
$(p - 1)(p^{2m-2} - \frac{p-2}{2}\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	F_1
$(p - 1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_1$
$(p - 1)p^{2m-2}$	$p^{2m} - p^\gamma - 1$

Table 11: The weight distribution of $C_{D_{f,g}}$ in Theorem 3 when $l_f = 2$ and $p \equiv 1 \pmod{8}$.

weight	frequency
0	1
$(p - 1)(p^{2m-2} - (p - 1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$\mathcal{N}_f(0)\mathcal{N}_g(0)$
$(p - 1)p^{2m-2} - (p + 1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4}$	F_3
$(p - 1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - F_3$
$(p - 1)p^{2m-2}$	$p^{2m} - p^\gamma - 1$

Table 12: The weight distribution of $C_{D_{f,g}}$ in Theorem 3 when $l_f = 2$ and $p \equiv 5 \pmod{8}$.

weight	frequency
0	1
$(p - 1)(p^{2m-2} - (p - 1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$\mathcal{N}_f(0)\mathcal{N}_g(0)$
$(p - 1)p^{2m-2} - (3p + 1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4}$	$\frac{1}{2}F_3$
$(p - 1)(p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4})$	$p^\gamma - \mathcal{N}_f(0)\mathcal{N}_g(0) - \frac{1}{2}F_3$
$(p - 1)p^{2m-2}$	$p^{2m} - p^\gamma - 1$

When $f, g \in \text{WRPB}$, the weight distributions in Tables 9 and 11 coincide with the results of Tables 6 and 7 in [18], respectively.

4.3. The punctured code

In the following, we study the punctured code from $C_{D_{f,g}}$ by deleting some coordinates of each codeword. As we can see from Tables 1, 3, 5 and 9, the length and each nonzero Hamming weight have $p - 1$ as a common divisor. This suggests that they can be punctured into shorter ones. Let $f \in \text{WRP}$ or $f \in \text{WRPB}$. For any $x \in \mathbb{F}_q$, we obtain $f(x) = 0$ if and only if $f(zx) = 0$ for all $z \in \mathbb{F}_p^*$, since $f(zx) = z^h f(x)$ for an even integer h with $\gcd(h - 1, p - 1) = 1$. Thus we can select a subset $\overline{D}_{f,g} = \{\overline{(x, y)} : (x, y) \in D_{f,g}\}$ from $D_{f,g}$ in (1.3), such that $\bigcup_{z \in \mathbb{F}_p^*} z\overline{D}_{f,g} = D_{f,g}$ forms a partition of $D_{f,g}$. Hence, we get the punctured code $C_{\overline{D}_{f,g}}$ from $C_{D_{f,g}}$. Moreover, the code $C_{\overline{D}_{f,g}}$ is projective since the minimum distance of its dual $C_{\overline{D}_{f,g}}^\perp$ is at least 3 as checked in [4]. We can also find some optimal codes when they meet certain specific conditions.

The following results related to the weight distributions of $C_{\overline{D}_{f,g}}$ follow directly from Tables 1, 3, 5 and 9, respectively. Remember that $\tau = 2m + s + t$ and $\gamma = 2m - s - t$.

Corollary 1. *Suppose that $f, g \in \text{WRP}$ or $f, g \in \text{WRPB}$ with $l_g = \frac{p-1}{2}$. Let $s + t$ be odd and \bar{d}^\perp be the minimum distance of $C_{\overline{D}_{f,g}}^\perp$. Then $\bar{d}^\perp \geq 3$. Moreover, if $l_f = \frac{p-1}{2}$, then $C_{\overline{D}_{f,g}}$ is a three-weight $[\frac{p^{2m-1}-1}{p-1}, 2m]$ linear code with its weight distribution listed in Table 13, and if $l_f = 2$ and $p \equiv 1 \pmod{8}$, then $C_{\overline{D}_{f,g}}$ is a five-weight $[\frac{p^{2m-1}-1}{p-1}, 2m]$ linear code with its weight distribution listed in Table 14, where E_3 and E_4 are computed in Theorem 1.*

Corollary 2. *Suppose that $f, g \in \text{WRP}$ with $l_f = l_g = \frac{p-1}{2}$ and $s + t$ is even. Let \bar{d}^\perp be the minimum distance of $C_{\overline{D}_{f,g}}^\perp$. Then $\bar{d}^\perp \geq 3$ and $C_{\overline{D}_{f,g}}$ is a three-weight $[\frac{p^{2m-1}-1}{p-1} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-2}, 2m]$ linear code with its weight distribution listed in Table 15. Moreover, the code $C_{\overline{D}_{f,g}}$ achieves Griesmer bound if $\tau = 4$ and $\varepsilon_f \varepsilon_g = -1$.*

Table 13: The weight distribution of $C_{\overline{D}_{f,g}}$ in Corollary 1 when $l_f = \frac{p-1}{2}$.

weight	frequency
0	1
p^{2m-2}	$p^{2m} - (p-1)p^{\gamma-1} - 1$
$p^{2m-2} - \sqrt{p}^{\tau-3}$	$\frac{p-1}{2}(p^{\gamma-1} + \sqrt{p}^{\gamma-1})$
$p^{2m-2} + \sqrt{p}^{\tau-3}$	$\frac{p-1}{2}(p^{\gamma-1} - \sqrt{p}^{\gamma-1})$

Table 14: The weight distribution of $C_{\overline{D}_{f,g}}$ in Corollary 1 when $l_f = 2$ and $p \equiv 1 \pmod{8}$.

weight	frequency
0	1
p^{2m-2}	$p^{2m} - 1 - E_3 - E_4 - \frac{(p-1)^2}{4}(\mathcal{N}_f(i)\mathcal{N}_g(i) + \mathcal{N}_f(j)\mathcal{N}_g(j))$
$p^{2m-2} - \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}$	E_3
$p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}$	E_4
$p^{2m-2} + 2\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}$	$\frac{(p-1)^2}{4}\mathcal{N}_f(i)\mathcal{N}_g(i)$
$p^{2m-2} - 2\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-3}$	$\frac{(p-1)^2}{4}\mathcal{N}_f(j)\mathcal{N}_g(j)$

Table 15: The weight distribution of $C_{\overline{D}_{f,g}}$ in Corollary 2.

weight	frequency
0	1
p^{2m-2}	$p^{\gamma-1} + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{\gamma-2} - 1$
$p^{2m-2} + \varepsilon_f \varepsilon_g \sqrt{p}^{\tau-2}$	$(p-1)(p^{\gamma-1} - \varepsilon_f \varepsilon_g \sqrt{p}^{\gamma-2})$
$p^{2m-2} + (p-1)\varepsilon_f \varepsilon_g \sqrt{p}^{\tau-4}$	$p^{2m} - p^\gamma$

Corollary 3. *Suppose that $f, g \in \text{WRPB}$ with $l_f = l_g = \frac{p-1}{2}$ and $s + t$ is even. Let \bar{d}^\perp be the minimum distance of $C_{\overline{D}_{f,g}}^\perp$. Then $\bar{d}^\perp \geq 3$ and $C_{\overline{D}_{f,g}}$ is a three-weight $[\frac{p^{2m-1}-1}{p-1}, 2m]$ linear code with its weight distribution listed in Table 16.*

Table 16: The weight distribution of $C_{\overline{D}_{f,g}}$ in Corollary 3.

weight	frequency
0	1
$p^{2m-2} - (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4}$	$p^{\gamma-1} + (p-1)\varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2}$
$p^{2m-2} + \varepsilon_f\varepsilon_g\sqrt{p}^{\tau-4}$	$(p-1)(p^{\gamma-1} - \varepsilon_f\varepsilon_g\sqrt{p}^{\gamma-2})$
p^{2m-2}	$p^{2m} - p^\gamma - 1$

Example 1. Let $f, g : \mathbb{F}_{5^4} \rightarrow \mathbb{F}_5$ be defined as $f(x) = \text{Tr}(x^6)$ and $g(y) = \text{Tr}(y^{26} - y^2)$. Then $f, g \in \text{WRP}$ with $s = t = 2$, $\varepsilon_f = -1$, $\varepsilon_g = 1$ and $l_f = l_g = 2$. Their Walsh transforms satisfy $\widehat{\chi}_f(\alpha) \in \{0, -5^3\zeta_5^{f^*(\alpha)}\}$ and $\widehat{\chi}_g(\beta) \in \{0, 5^3\zeta_5^{g^*(\beta)}\}$, where $\alpha, \beta \in \mathbb{F}_{5^4}$ and $f^*(0) = g^*(0) = 0$. Hence $C_{D_{f,g}}$ is a three-weight code with parameters $[65624, 8, 50000]$ and the weight enumerator $1 + 520z^{50000} + 390000z^{52500} + 104z^{62500}$. Its punctured code $C_{\overline{D}_{f,g}}$ has parameters $[16406, 8, 12500]$ and the weight enumerator $1 + 520z^{12500} + 390000z^{13125} + 104z^{15625}$.

Example 2. Let $f, g : \mathbb{F}_{5^3} \rightarrow \mathbb{F}_5$ be defined as $f(x) = \text{Tr}(x^6 + x^2)$ and $g(y) = \text{Tr}(\theta y^6 + \theta^3 y^2)$ for a primitive element θ of $\mathbb{F}_{5^3}^*$. Then $f, g \in \text{WRP}$ with $s = 0$, $t = 1$, $\varepsilon_f = -1$, $\varepsilon_g = 1$, $l_f = l_g = 2$, $\widehat{\chi}_f(\alpha) \in \{-\sqrt{5}^3\zeta_5^{f^*(\alpha)}\}$ and $\widehat{\chi}_g(\beta) \in \{0, 5^2\zeta_5^{g^*(\beta)}\}$, where $\alpha, \beta \in \mathbb{F}_{5^3}$ and $f^*(0) = g^*(0) = 0$. Actually, the function f is quadratic bent and its Walsh transform satisfies $|\widehat{\chi}_f(\alpha)|^2 = 125$. Hence $C_{D_{f,g}}$ is a three-weight code with parameters $[3124, 6, 2400]$ and the weight enumerator $1 + 1300z^{2400} + 13124z^{2500} + 1200z^{2600}$. Its punctured code $C_{\overline{D}_{f,g}}$ has parameters $[781, 6, 600]$ and the weight enumerator $1 + 1300z^{600} + 13124z^{625} + 1200z^{650}$.

Example 3. Let $f, g : \mathbb{F}_{5^2} \rightarrow \mathbb{F}_5$ be defined as $f(x) = \text{Tr}(x^2)$ and $g(y) = \text{Tr}(\theta y^2 - \theta y^6)$ for a primitive element θ of $\mathbb{F}_{5^2}^*$. Then f, g are quadratic bent

functions in the set WRP, with $s = t = 0$, $\varepsilon_f = -1$, $\varepsilon_g = 1$, $l_f = l_g = 2$, $\widehat{\chi}_f(\alpha) \in \{-5\zeta_5^{f^*(\alpha)}\}$ and $\widehat{\chi}_g(\beta) \in \{5\zeta_5^{g^*(\beta)}\}$, where $\alpha, \beta \in \mathbb{F}_{5^2}$ and $f^*(0) = g^*(0) = 0$. Then the code $C_{D_{f,g}}$ is a two-weight code with parameters $[104, 4, 80]$ and the weight enumerator $1 + 520z^{80} + 104z^{100}$. Its punctured code $C_{D_{f,g}}^-$ has parameters $[26, 4, 20]$ and the weight enumerator $1 + 520z^{20} + 104z^{25}$. The punctured code is optimal with respect to the Griesmer bound.

5. Minimality of the codes and their applications

Any linear code can be applied to design secret sharing schemes by considering the access structure. However, the access structure based on a linear code is usually very complicated, and only can be determined exactly in several specific cases. One such case is when the code is minimal.

A linear code C over \mathbb{F}_p is called minimal if every nonzero codeword \mathbf{c} of C solely covers its scalar multiples $z\mathbf{c}$ for $z \in \mathbb{F}_p^*$. In 1998, Ashikhmin and Barg [1] provided a well-known criteria for minimal linear codes.

Lemma 16. (*Ashikhmin-Barg Bound [1]*) *Let C be a linear code over \mathbb{F}_p . Then all nonzero codewords of C are minimal, provided that*

$$\frac{w_{min}}{w_{max}} > \frac{p-1}{p},$$

where w_{min} and w_{max} stand for the minimum and maximum nonzero weights in C , respectively.

Now we will show under what circumstances the constructed linear codes are minimal according to Lemma 16.

Theorem 4. *We have the following bounds on parameters of the code $C_{D_{f,g}}$.*

(1) *The linear codes described in Tables 1, 2 and 3 are minimal provided when $\varepsilon_f \varepsilon_g \in \{\pm 1\}$ and $2m - s - t \geq 5$.*

(2) *The linear codes described in Tables 5, 6 7 and 8 are minimal provided when $\varepsilon_f \varepsilon_g = 1$ and $2m - s - t \geq 4$, or $\varepsilon_f \varepsilon_g = -1$ and $2m - s - t \geq 6$.*

(3) *The linear codes described in Tables 9, 10, 11 and 12 are minimal provided when $\varepsilon_f \varepsilon_g \in \{\pm 1\}$ and $2m - s - t \geq 4$.*

Remark 3. Our punctured codes $C_{\overline{D}_{f,g}}$ are minimal for almost all cases.

It should be noticed that the minimum distance of $C_{\overline{D}_{f,g}}^\perp$ equals 2 since there are two linearly dependent entries in each codeword in $C_{\overline{D}_{f,g}}$. So under the framework stated in [6], the minimal codes described in Theorems 1, 2 and 3 can be employed to construct high democratic secret sharing schemes with new parameters. The punctured codes are projective and minimal, as we have discussed previously. So they are also suitable for secret sharing schemes. The projective three-weight codes in Tables 13, 15 and 16 can be applied to design association schemes [2].

6. Conclusion

The paper studied the construction of linear codes using defining set from two weakly regular plateaued functions with index $\frac{p-1}{2}$ for $p \equiv 1 \pmod{4}$, and hence, this is an extension of the results in [4], [18] and [21]. The punctured codes were also investigated and we found optimal codes among them. Moreover, our codes are suitable for designing association schemes and secret sharing schemes.

References

- [1] A. Ashikhmin and A. Barg, Minimal vectors in linear codes, *IEEE Transactions on Information Theory*, **44** (5) (1998), 2010–2017.
- [2] A. R. Calderbank and J.-M. Goethals, Three-weight codes and association schemes, *Philips Journal of Research*, **39** (1984), 143–152.
- [3] H. Chen, C. Ding, S. Mesnager and C. Tang, A novel application of Boolean functions with high algebraic immunity in minimal codes, *IEEE Transactions on Information Theory*, **67** (10) (2021), 6856–6867.
- [4] Y. Cheng and X. Cao, Linear codes with few weights from weakly regular plateaued functions, *Discrete Mathematics*, **344** (2021), 112597.

- [5] C. Ding and H. Niederreiter, Cyclotomic linear codes of order 3, *IEEE Transactions on Information Theory*, **53** (6) (2007), 2274–2277.
- [6] C. Ding and J. Yuan, Covering and secret sharing with linear codes, in: *Discrete Mathematics and Theoretical Computer Science 2731* (eds. G. Goos, J. Hartmanis and J. V. Leeuwen), Springer Berlin Heidelberg, (2003), 11–25.
- [7] Z. Heng, C. Ding and Z. Zhou, Minimal linear codes over finite fields, *Finite Fields and Their Applications*, **54** (2018), 176–196.
- [8] Z. Heng, D. Li, J. Du and F. Chen, A family of projective two-weight linear codes, *Designs, Codes and Cryptography*, **89** (2021), 1993–2007.
- [9] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, New York, 1990.
- [10] X. Kong and S. Yang, Complete weight enumerators of a class of linear codes with two or three weights, *Discrete Mathematics*, **342** (2019), 3166–3176.
- [11] C. Li, Q. Yue and F. Fu, A construction of several classes of two-weight and three-weight linear codes, *Applicable Algebra in Engineering, Communication and Computing*, **28** (2017), 11–30.
- [12] H. Lu and S. Yang, Two classes of linear codes from Weil sums, *IEEE Access*, **8** (2020), 180471–180480.
- [13] R. Lidl and H. Niederreiter, *Finite Fields*, Cambridge University Press, Cambridge, 1997.
- [14] S. Mesnager, Characterizations of plateaued and bent functions in characteristic p , in: *International Conference on Sequences and their Applications, SETA-2014*, Springer, Cham, 2014, pp. 72–82.

- [15] S. Mesnager, F. Özbudak, A. Snak, Linear codes from weakly regular plateaued functions and their secret sharing schemes, *Designs, Codes and Cryptography*, **87** (2019), 463–480.
- [16] S. Mesnager and A. Snak, Several classes of minimal linear codes with few weights from weakly regular plateaued functions, *IEEE Transactions on Information Theory*, **66** (4) (2020), 2296–2310.
- [17] A. Snak, Minimal linear codes from weakly regular plateaued balanced functions, *Discrete Mathematics*, **344** (2021), 112215.
- [18] A. Snak, Construction of minimal linear codes with few weights from weakly regular plateaued functions, *Turkish Journal of Mathematics*, **46** (2022), 953–972.
- [19] C. Tang, N. Li, Y. Qi, Z. Zhou and T. Helleseht, Linear codes with two or three weights from weakly regular bent functions, *IEEE Transactions on Information Theory*, **62** (3) (2016), 1166–1176.
- [20] C. Tang, Y. Qi and D. Huang, Two-weight and three-weight linear codes from square functions, *IEEE Communications Letters*, **20** (1) (2016), 29–32.
- [21] Y. Wu, N. Li and X. Zeng, Linear codes with few weights from cyclotomic classes and weakly regular bent functions, *Designs, Codes and Cryptography*, **88** (2020), 1255–1272.
- [22] S. Yang, Complete weight enumerators of linear codes based on Weil sums, *IEEE Communications Letters*, **25** (2) (2021), 346–350.
- [23] T. Zhang, H. Lu and S. Yang, Two-weight and three-weight linear codes constructed from Weil sums, *Mathematical Foundations of Computing*, **5** (2022), 129–144.
- [24] Y. Zheng and X. Zhang, Plateaued functions, in *Proc. ICICS*, vol. 99, Springer, 1999, pp. 284–300.

- [25] D. Zheng, Q. Zhao, X. Wang and Y. Zhang, A class of two or three weights linear codes and their complete weight enumerators, *Discrete Mathematics*, **344** (2021), 112355.
- [26] F. Özbudak and R. M. Pelen, Two or three weight linear codes from non-weakly regular bent functions, *IEEE Transactions on Information Theory*, **68** (5) (2022), 3014–3027.