

# THE COHOMOLOGY OF THE CLASSIFYING SPACE OF $PU(4)$

FEIFEI FAN

ABSTRACT. Let  $BPU(n)$  be the classifying space of the projective unitary group  $PU(n)$ . We determine the integral cohomology ring of  $BPU(4)$ , and the Steenrod algebra structure of its mod 2 cohomology.

## 1. INTRODUCTION

Let  $G$  be a topological group. The classifying space  $BG$  of  $G$  is characterized as the base space of a universal  $G$ -bundle  $G \rightarrow EG \rightarrow BG$ . The cohomology of  $BG$  is the main tool in the theory of classifying principal  $G$ -bundles.

Let  $U(n)$  be the group of  $n \times n$  unitary matrices. The projective unitary group  $PU(n)$  is the quotient group  $U(n)/S^1$  of  $U(n)$  by its center subgroup  $S^1 = \{e^{i\theta}I_n : \theta \in [0, 2\pi]\}$ , where  $I_n$  denotes the identity matrix.  $PU(n)$  is homotopy equivalent to  $PGL(n, \mathbb{C})$ , the projective general linear group  $PGL(n, \mathbb{C})$ , which is the automorphism group of the algebra of  $n \times n$  complex matrices. Therefore, for a pointed space  $X$ , the set of pointed homotopy classes of maps  $[X, BPU(n)]$  also classifies bundles of  $n \times n$  complex matrix algebras over  $X$ , known as the topological Azumaya algebras of degree  $n$  over  $X$ . (See [8].)

The cohomology ring of  $BPU(n)$  also plays a significant role in the study of the topological period-index problem, which was introduced by Antieau and Williams [1, 2] as an analogue of period-index problems in algebraic geometry (cf. [3, 6, 19]). Gu [9, 10], Crowley and Grant [5] also investigated this problem for certain topological spaces.

Since  $PU(n)$  can also be viewed as the quotient group of the special unitary group  $SU(n)$  by its center subgroup  $\mathbb{Z}/n$  generated by  $e^{2\pi i/n}I_n$ , there is a induced fibration of classifying spaces:

$$B(\mathbb{Z}/n) \rightarrow BSU(n) \rightarrow BPU(n).$$

Hence, for a commutative ring  $R$ , if  $1/n \in R$ , then

$$H^*(BPU(n); R) \cong H^*(BSU(n); R) \cong R[c_2, c_3, \dots, c_n], \quad \deg(c_i) = 2i.$$

---

2020 *Mathematics Subject Classification.* 55R35, 55R40, 55S10, 55T10, 57T25.

The author is supported by the National Natural Science Foundation of China (Grant no. 12271183) and by the GuangDong Basic and Applied Basic Research Foundation (Grant no. 2023A1515012217).

It follows that if  $x \in H^*(BPU(n); \mathbb{Z})$  is a torsion element, then there exists  $k \geq 0$  such that  $n^k x = 0$ . (In the case of Chow rings, Vezzosi [23] proved the stronger result that all torsion classes in the Chow ring of  $BPGL(n, \mathbb{C})$  are  $n$ -torsion.)

However, if  $n$  is not invertible in the coefficient ring  $R$ , the ring structure of  $H^*(BPU(n); R)$  is very complicated for general  $n$ . Here we list some of the known results on the cohomology of  $BPU(n)$  for special values of  $n$ .

Since  $PU(2) = SO(3)$ , the ring  $H^*(BPU(2); \mathbb{Z})$  is well understood as a special case of the result of Brown [14].

The ring  $H^*(BPU(3); \mathbb{Z}/3)$  is computed in Kono-Mimura-Shimada [16]. The Steenrod algebra structure of  $H^*(BPU(3); \mathbb{Z}/3)$  and the Brown-Peterson cohomology of  $BPU(3)$  are determined in Kono-Yagita [17].

The ring  $H^*(BPU(n); \mathbb{Z}/2)$  is known if  $n \equiv 2 \pmod{4}$  (Kono-Mimura [15] and Toda [21]). In [21], Toda also computed the ring  $H^*(BPU(4); \mathbb{Z}/2)$ .

In [24], Vistoli provided a description of  $H^*(BPU(p); \mathbb{Z})$ , as well as the Chow ring of  $BPGL(p, \mathbb{C})$ , for any odd prime  $p$ . In particular, the ring structure of  $H^*(BPU(3); \mathbb{Z})$  is given in terms of generators and relations there. We will explicitly explain Vistoli's result in the next Section.

More interesting results on the ordinary cohomology ring  $H^*(BPU(p); \mathbb{Z}/p)$  (resp. on the Brown-Peterson cohomology of  $BPU(p)$ ) for an arbitrary prime  $p$  can be found in Vavpetič-Viruel [22] (resp. in Masaki-Yagita [20]).

For an arbitrary integer  $n \geq 3$ , the cohomology of  $BPU(n)$  is only known in certain finite range of dimensions. For instance, Gu [11] determined the ring structure of  $H^*(BPU(n); \mathbb{Z})$  in dimensions less than or equal to 10. Other partial results on the  $p$ -local cohomology groups of  $BPU(n)$  for an odd prime  $p$ ,  $p \mid n$ , can be found in Gu-Zhang-Zhang-Zhong [13] and Zhang-Zhang-Zhong [25].

Summarizing these results, the only examples of  $BPU(n)$ , whose integral cohomology rings can be described via generators and relations, are the cases  $n = 2, 3$ . Also, these are the only cases for which the Steenrod algebra structure of  $H^*(BPU(n); \mathbb{Z}/p)$ ,  $p \mid n$  a prime, are known. The main goal of this paper is to determine the integral cohomology ring of  $BPU(4)$  and the Steenrod algebra structure of its mod 2 cohomology.

## 2. MAIN RESULTS

**Notations.** For an integer  $n > 0$ , let  $\Lambda_n$  be the ring of symmetric polynomials over  $\mathbb{Z}$  in  $n$  variables. It is known that  $\Lambda_n = \mathbb{Z}[\sigma_1, \dots, \sigma_n]$ , where  $\sigma_i$  is the  $i$ th elementary symmetric polynomial in  $n$  variables. Let  $\Lambda_n^d \subset \Lambda_n$  be the  $\mathbb{Z}$ -submodule of symmetric polynomials of degree  $d$ .

Let  $\nabla = \sum_{i=1}^n \partial/\partial v_i$  be the linear differential operator acting on the polynomial algebra  $\mathbb{Z}[v_1, \dots, v_n]$ . It is easy to see that  $\nabla$  on  $\mathbb{Z}[v_1, \dots, v_n]$  preserves symmetric polynomials, so  $\nabla$  can be restricted to  $\Lambda_n$ . Generally, for a commutative ring  $R$  with unit, we use the notation  $\nabla_R$  to denote the map  $\nabla \otimes R$ . Let  $K_n$  be the kernel of  $\nabla$  on  $\Lambda_n$ .

For the graded polynomial ring  $\mathbb{Z}[v_1, \dots, v_n]$  with  $\deg(v_i) = 2$ , define a graded algebraic homomorphism  $\Theta_n : \mathbb{Z}[v_1, \dots, v_n] \rightarrow \mathbb{Z}[\eta]/(n\eta)$ ,  $\deg(\eta) = 2$ , by  $\Theta(v_i) = i\eta$ . In fact, this is the ring homomorphism  $H^*(BT^n; \mathbb{Z}) \rightarrow H^*(B(\mathbb{Z}/n); \mathbb{Z})$  induced by the embedding  $\mathbb{Z}/n \hookrightarrow T^n$ ,  $\omega \mapsto (\omega, \omega^2, \dots, \omega^{n-1}, 1)$ ,  $\omega = e^{2\pi i/n}$ ,  $T^n = (S^1)^n$ . Let  $\Theta'_n$  be the restriction of  $\Theta_n$  to  $K_n$ , and let  $K'_n = \ker \Theta'_n$ . The result of Vistoli [24] can be described as follows.

**Theorem 2.1** (Vistoli [24, Theorem 3.4]). *For any prime  $p > 2$ , the integral cohomology ring of  $BPU(p)$  is given by*

$$H^*(BPU(p); \mathbb{Z}) \cong \frac{K_p \otimes \mathbb{Z}[x_3, x_{2p+2}]}{(x_3^2, px_3, px_{2p+2}, x_3 K'_p, x_{2p+2} K'_p)}, \quad \deg(x_i) = i.$$

For the element  $\delta = \prod_{i \neq j} (v_i - v_j) \in K_p$ , one easily checks that

$$\Theta'_p(\delta) \equiv -\eta^{p^2-p} \neq 0 \pmod{(p\eta)}.$$

It is shown in [24, Proposition 3.1] that the image of  $\Theta'_p$  is generated by  $\Theta'_p(\delta)$ , so for  $p > 2$ , as  $\mathbb{Z}$ -module,

$$H^*(BPU(p); \mathbb{Z}) \cong K_p \oplus ((E_{\mathbb{Z}/p}[x_3] \otimes \mathbb{Z}/p[x_{2p+2}])^+ \otimes \mathbb{Z}[\delta]),$$

where  $E_{\mathbb{Z}/p}[x_3]$  means the exterior algebra over  $\mathbb{Z}/p$  with the generator  $x_3$ . Since the rank of  $K_p$  is the same as the rank of  $H^*(BSU(n); \mathbb{Z}) \cong \mathbb{Z}[c_2, \dots, c_n]$ , the additive structure of  $H^*(BPU(p); \mathbb{Z})$  are determined completely.

One reason for the difficulty of describing  $H^*(BPU(n); \mathbb{Z})$  is that the ring  $K_n$  is hard to compute for large  $n$ . For small values of  $n$ , here are some results.  $K_2 \cong \mathbb{Z}[\alpha_2]$ ,  $\alpha_2 = \sigma_1^2 - 4\sigma_2 \in \Lambda_2^2$ , is obvious. Vezzosi [23, Lemma 3.2] (see also [24, Theorem 14.2]) proved that

$$K_3 \cong \mathbb{Z}[\alpha_2, \alpha_3, \alpha_6]/(27\alpha_6 - 4\alpha_2^3 - \alpha_3^2), \quad \alpha_i \in \Lambda_3^i.$$

Here we give the description of  $K_4$ .

**Theorem 2.2.** *There is a ring isomorphism*

$$K_4 \cong \mathbb{Z}[\alpha_2, \alpha_3, \alpha_4, \alpha_6]/(64\alpha_6 - \alpha_2^3 - 27\alpha_3^2 + 48\alpha_2\alpha_4), \quad \alpha_i \in \Lambda_4^i.$$

For  $n \geq 5$ , the complexity of  $K_n$  grows very quickly as  $n$  grows. To the author's knowledge, there is no one computed  $K_n$  for  $n \geq 5$  by writing down the generators and relations explicitly so far.

As in the case of  $K_p$  for  $p$  a prime,  $K_4$  can be viewed as the quotient ring of  $H^*(BPU(4); \mathbb{Z})$  by torsion elements.

**Theorem 2.3.** *The cohomology ring  $H^*(BPU(4); \mathbb{Z})$  is isomorphic to the graded ring*

$$K_4 \otimes \mathbb{Z}[x_3, x_{10}, x_{15}]/I, \quad \deg(\alpha_i) = 2i, \quad \deg(x_i) = i,$$

where  $I$  is the ideal generated by

$$4x_3, 2x_3^2, \alpha_2x_3, \alpha_3x_3, 2x_{10}, \alpha_2x_{10}, \alpha_3x_{10}, 2x_{15}, \alpha_2x_{15}, \alpha_3x_{15}, \\ x_3^6\alpha_6 + x_3^4x_{10}\alpha_4 + x_3^5x_{15} + x_{10}^3 + x_{15}^2.$$

One of the results in Toda [21] gives  $H^*(BPSO(4m+2); \mathbb{Z}/2)$ ,  $m \geq 1$ . In particular, this gives the mod 2 cohomology of  $BPU(4)$ , since it is known that  $PU(4)$  is homeomorphic to  $PSO(6)$ .

**Theorem 2.4** (Toda [21, (4.10)]). *The mod 2 cohomology of  $BPU(4)$  is given by*

$$H^*(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2[y_2, y_3, y_5, y_8, y_9, y_{12}]/J, \quad \deg(y_i) = i,$$

where  $J$  is the ideal generated by

$$y_2y_3, y_2y_5, y_2y_9, y_9^2 + y_3^2y_{12} + y_5^2y_8.$$

In the notation of Theorem 2.4, we determine the Steenrod algebra structure of  $H^*(BPU(4); \mathbb{Z}/2)$  as follows.

**Theorem 2.5.**  $Sq^1(y_2) = Sq^1(y_3) = 0$ ,  $Sq^1(y_5) = y_3^2$ ,  $Sq^1(y_8) = x_3^3$ ,  $Sq^1(y_9) = y_5^2$ ,  $Sq^1(y_{12}) = y_3y_5^2$ ;  $Sq^2(y_3) = y_5$ ,  $Sq^2(y_5) = 0$ ,  $Sq^2(y_8) = y_5^2$ ,  $Sq^2(y_9) = y_3^2y_5$ ,  $Sq^2(y_{12}) = y_2y_{12} + y_3^2y_8$ ;  $Sq^4(y_5) = y_3^3 + y_9$ ,  $Sq^4(y_8) = y_2^2y_8 + y_{12} + y_3^4$ ,  $Sq^4(y_9) = y_3y_5^2$ ,  $Sq^4(y_{12}) = y_2^2y_{12} + y_3^2y_5^2$ ;  $Sq^8(y_9) = y_3y_5y_9 + y_5y_{12} + y_8y_9$ ,  $Sq^8(y_{12}) = y_3^4y_8 + y_8y_{12}$ .

In the next two sections, we review Gu's work in [11], where he constructs a fibration with  $BPU(n)$  as the total space, and provides an approach to calculate the Serre spectral sequence associated to this fibration.

### 3. SERRE SPECTRAL SEQUENCES IN THE COMPUTATION OF $H^*(BPU(n))$

**3.1. The spectral sequence  ${}^U E$ .** The short exact sequence of Lie groups

$$1 \rightarrow S^1 \rightarrow U(n) \rightarrow PU(n) \rightarrow 1$$

induces a fibration of their classifying spaces

$$BS^1 \rightarrow BU(n) \xrightarrow{\pi} BPU(n). \quad (3.1)$$

Notice that  $BS^1$  is of the homotopy type of the Eilenberg-Mac Lane space  $K(\mathbb{Z}, 2)$ . So we obtain another fibration:

$$U : BU(n) \xrightarrow{\pi} BPU(n) \xrightarrow{\chi} K(\mathbb{Z}, 3). \quad (3.2)$$

(Cf. [7].) Let  ${}^U E$  be the cohomological Serre spectral sequence associated to this fibration. The  $E_2$  page of this spectral sequence has the form

$${}^U E_2^{s,t} = H^s(K(\mathbb{Z}, 3); H^t(BU(n))) \implies H^{s+t}(BPU(n)).$$

Notice that  ${}^U E_2^{*,*}$  is concentrated in even rows, so all  $d_2$  differentials are trivial, and we have  ${}^U E_2^{*,*} = {}^U E_3^{*,*}$ .

**3.2. The spectral sequences  ${}^K E$  and  ${}^T E$ .** Let  $T^n$  and  $PT^n$  be the maximal tori of  $U(n)$  and  $PU(n)$  respectively, and let  $\psi : T^n \rightarrow U(n)$  and  $\psi' : PT^n \rightarrow BU(n)$  be the inclusions. Then there is an exact sequence of Lie groups

$$1 \rightarrow S^1 \rightarrow T^n \rightarrow PT^n \rightarrow 1,$$

which as above induces a fibration

$$T : BT^n \rightarrow BPT^n \rightarrow K(\mathbb{Z}, 3).$$

We also consider the path fibration

$$K : BS^1 \simeq K(\mathbb{Z}, 2) \rightarrow * \rightarrow K(\mathbb{Z}, 3),$$

where  $*$  denotes a contractible space. Let  ${}^T E$  and  ${}^K E$  denote the Serre spectral sequences associated to  $T$  and  $K$  respectively.

Let  $\varphi : S^1 \rightarrow T^n$  be the diagonal map. Then there is the following homotopy commutative diagram of fibrations

$$\begin{array}{ccccc} K : & BS^1 & \longrightarrow & * & \longrightarrow & K(\mathbb{Z}, 3) \\ & \downarrow \Phi & & \downarrow B\varphi & & \downarrow = \\ T : & BT^n & \longrightarrow & BPT^n & \longrightarrow & K(\mathbb{Z}, 3) \\ & \downarrow \Psi & & \downarrow B\psi & & \downarrow = \\ U : & BU(n) & \longrightarrow & BPU(n) & \longrightarrow & K(\mathbb{Z}, 3) \end{array} \quad (3.3)$$

We write  ${}^U d_*^{*,*}$ ,  ${}^T d_*^{*,*}$ ,  ${}^K d_*^{*,*}$  for the differentials of  ${}^U E$ ,  ${}^T E$ , and  ${}^K E$ , respectively. We also use the simplified notation  $d_*^{*,*}$  if there is no ambiguity.

**3.3. Spectral sequence maps on  $E_2$  pages.** Recall that the cohomology rings of the fibers  $BS^1$ ,  $BT^n$  and  $BU(n)$  are

$$\begin{aligned} H^*(BS^1; \mathbb{Z}) &= \mathbb{Z}[v], \quad \deg(v) = 2, \\ H^*(BT^n; \mathbb{Z}) &= \mathbb{Z}[v_1, \dots, v_n], \quad \deg(v_i) = 2, \\ H^*(BU(n); \mathbb{Z}) &= \mathbb{Z}[c_1, \dots, c_n], \quad \deg(c_i) = 2i. \end{aligned}$$

Here  $c_i$  is the  $i$ th universal Chern class of the classifying space  $BU(n)$  for  $n$ -dimensional complex bundles. The induced homomorphisms between these cohomology rings are given by

$$B\varphi^*(v_i) = v, \quad B\psi^*(c_i) = \sigma_i(v_1, \dots, v_n),$$

where  $\sigma_i(v_1, \dots, v_n)$  is the  $i$ th elementary symmetric polynomial in the variables  $v_1, \dots, v_n$ . Hence, from (3.3) we obtain the maps  $\Phi^* : {}^T E_2^{*,*} \rightarrow {}^K E_2^{*,*}$  and  $\Psi^* : {}^U E_2^{*,*} \rightarrow {}^T E_2^{*,*}$  of spectral sequences on  $E_2$  pages.

The strategy in [11] for computing  ${}^U E$  is to compare its differentials with the differentials in  ${}^T E$  and  ${}^K E$ , which we will explain in the next section.

#### 4. THE COHOMOLOGY OF $K(\mathbb{Z}, 3)$ AND DIFFERENTIALS OF ${}^K E$ AND ${}^T E$

Throughout this section, we write  $R[x; k]$ ,  $E_R[y; k]$  for the polynomial algebra and exterior algebra respectively, with coefficients in a commutative ring  $R$ , and with one generator of degree  $k$ . We also use the simplified notation  $E[y; k]$  for  $R = \mathbb{Z}$ .

Let  $C(3)$  be the (differential graded)  $\mathbb{Z}$ -algebra

$$\begin{aligned} C(3) = & \bigotimes_{p \neq 2 \text{ prime}} \left( \bigotimes_{k \geq 0} \mathbb{Z}[y_{p,k}; 2p^{k+1} + 2] \otimes E[x_{p,k}; 2p^{k+1} + 1] \right) \\ & \otimes \bigotimes_{k \geq 0} \mathbb{Z}[x_{2,k}; 2^{k+2} + 1] \otimes \mathbb{Z}[x_1; 3]. \end{aligned}$$

For notational convenience, we set  $y_{2,0} = x_1^2$  and  $y_{2,k} = x_{2,k-1}^2$  for  $k \geq 1$ . The differential of  $C(3)$  is given by

$$d(x_{p,k}) = py_{p,k}, \quad d(y_{p,k}) = d(x_1) = 0, \quad \text{for } p \text{ a prime and } k \geq 0.$$

Gu [11] showed that the integral cohomology ring of  $K(\mathbb{Z}, 3)$  is isomorphic to the cohomology of  $C(3)$ , which is dual to the DGA construction of Cartan in [4].

Let  ${}_p A$  denote the  $p$ -primary component of a commutative ring  $A$ ,  $p$  a prime. For  $R = \bigoplus_{i > 0} R^i$  a graded commutative ring without unit, let  $\hat{R} = R \oplus R^0$  be the (graded commutative) unital ring, where the degree zero summand  $R^0 = \mathbb{Z}$  has a generator acting as the unit of the ring.

**Theorem 4.1** ([11, Proposition 2.14]). *The cohomology ring of  $K(\mathbb{Z}, 3)$  is given by*

$$H^*(K(\mathbb{Z}, 3); \mathbb{Z}) \cong \bigotimes_p \widehat{R}_p \otimes \mathbb{Z}[x_1] / (x_1^2 - y_{2,0}),$$

where  $p$  ranges over all prime numbers, and  $R_p = {}_pH^*(K(\mathbb{Z}, 3); \mathbb{Z})$  is the  $\mathbb{Z}/p$ -subalgebra (without unit) of  $C(3) \otimes \mathbb{Z}/p$  generated by the elements of the form

$$y_{p,I} := \frac{1}{p}d(x_{p,i_1} \cdots x_{p,i_m}) = \sum_{j=1}^m (-1)^{j-1} x_{p,i_1} \cdots y_{p,i_j} \cdots x_{p,i_m},$$

for any ordered sequence  $I = (i_1, \dots, i_m)$  of nonnegative integers  $i_1 > \cdots > i_m$ . For  $I = (i)$ , we simply write  $y_{p,i}$  for  $y_{p,I}$ .

**Remark 4.2.** In [4, Exposé 11] a differential graded algebra  $A(\pi, n)$ , which is dual to  $C(3)$  for  $n = 3$  and  $\pi = \mathbb{Z}$ , is constructed to calculate the integral homology of  $K(\pi, n)$  for any positive integer  $n$  and a finitely generated abelian group  $\pi$ . [11] first gives an explicit formula for  $H^*(K(\mathbb{Z}, 3); \mathbb{Z})$  in terms of generators and relations.

For  $p$  a prime, the mod  $p$  cohomology ring of  $K(\mathbb{Z}, 3)$  is also given by the cohomology of the differential graded  $\mathbb{Z}/p$ -algebra  $(C(3) \otimes \mathbb{Z}/p, d)$ . Since the differential of  $C(3)$  localized at  $p$  is trivial modulo  $p$ ,  $H^*(K(\mathbb{Z}, 3); \mathbb{Z}/p)$  is isomorphic to  $C(3) \otimes \mathbb{Z}/p$ . Furthermore, the Steenrod algebra structure of  $H^*(K(\mathbb{Z}, 3); \mathbb{Z}/p)$  can be determined from the fact that  $x_1$  (resp.  $x_{p,k}$ ) is the the transgression image of  $v$  (resp.  $v^{p^{k+1}} \in H^*(BS^1; \mathbb{Z}/p) \cong \mathbb{Z}/p[v]$ ) in the mod  $p$  Serre spectral sequence  ${}^K E$ , using Kudo's transgression theorem [18]. In particular, for  $p = 2$ , we have the following analogue of [12, Proposition 2.4].

**Proposition 4.3.**  $H^*(K(\mathbb{Z}, 3); \mathbb{Z}/2) \cong \bigotimes_{k \geq 0} \mathbb{Z}/2[x_{2,k}; 2^{k+2} + 1] \otimes \mathbb{Z}/2[x_1; 3]$ . Moreover,  $Sq^1(x_1) = 0$ ,  $Sq^1(x_{2,k+1}) = x_{2,k}^2$ ,  $x_{2,k} = Sq^{2^{k+1}} Sq^{2^k} \cdots Sq^2(x_1)$ ,  $k \geq 0$ .

The differentials of the integral spectral sequence  ${}^K E$  are described in [11] as follows. Here we only list a part of the results in [11, Corollary 2.16], which will be sufficient for our purposes.

**Proposition 4.4** ([11, Corollary 2.16]). *The higher differentials of  ${}^K E_{*}^{*,*}$  satisfy*

$$\begin{aligned} d_3(v) &= x_1, \\ d_{2p^{k+1}-1}(p^k x_1 v^{lp^e-1}) &= v^{lp^e-1-(p^{k+1}-1)} y_{p,k}, \quad e > k \geq 0, \quad \gcd(l, p) = 1, \\ d_r(x_1) &= d_r(y_{p,k}) = 0 \text{ for all } r \geq 2, \quad k \geq 0, \end{aligned}$$

and the Leibniz rule. Here  ${}^K E_{2p^{k+1}-1}^{3,2(lp^e-1)} \subset {}^K E_2^{3,2(lp^e-1)}$  is generated by  $p^k x_1 v^{lp^e-1}$ .

To describe the differential of  ${}^T E_{*}^{*,*}$ , we first rewrite  $H^*(BT^n; \mathbb{Z}) = \mathbb{Z}[v_1, \dots, v_n]$  as  $\mathbb{Z}[v_1 - v_n, \dots, v_{n-1} - v_n, v_n]$ . Then an element in  ${}^T E_{*}^{*,*}$  can be written as  $f(v_n)\xi$ , where  $\xi \in H^*(K(\mathbb{Z}, 3); \mathbb{Z})$  and  $f(v_n) = \sum_i a_i v_n^i$  is a polynomial in  $v_n$  with coefficients  $a_i \in \mathbb{Z}[v_1 - v_n, \dots, v_{n-1} - v_n]$ . Let  $v'_j = v_j - v_n$  for  $1 \leq j \leq n-1$ , and write  $a_i = \sum_{t_1, \dots, t_{n-1}} k_{i, t_1, \dots, t_{n-1}} (v'_1)^{t_1} \cdots (v'_{n-1})^{t_{n-1}}$ ,  $k_{i, t_1, \dots, t_{n-1}} \in \mathbb{Z}$ .

**Proposition 4.5** ([11, Proposition 3.2, 3.3]). *In the above notation, the differential  ${}^T d_r^{*,*}$  is given by*

$${}^T d_r^{*,*}(f(v_n)\xi) = \sum_i \sum_{t_1, \dots, t_{n-1}} k_{i, t_1, \dots, t_{n-1}} {}^K d_r^{*,*}(\xi v_n^i) \cdot (v'_1)^{t_1} \cdots (v'_{n-1})^{t_{n-1}},$$

where  ${}^K d_r^{*,*}(\xi v_n^i)$  is simply  ${}^K d_r^{*,*}(\xi v^i)$  with  $v$  replaced by  $v_n$ . In particular, the differential  ${}^T d_3^{0,t}$  is given by the “formal divergence”

$$\nabla = \sum_{i=1}^n (\partial/\partial v_i) : H^t(BT^n; \mathbb{Z}) \rightarrow H^{t-2}(BT^n; \mathbb{Z})$$

in such a way that  ${}^T d_3^{0,*} = \nabla(-) \cdot x_1$ . Moreover, the spectral sequence  ${}^T E_*^{*,*}$  degenerates at  ${}^T E_4^{0,*}$ , i.e.,

$${}^T E_\infty^{0,*} = {}^T E_4^{0,*} = \ker {}^T d_3^{0,*} = \mathbb{Z}[v'_1, \dots, v'_{n-1}].$$

Since the operator  $\nabla$  preserves symmetric polynomials, for  $f \in H^*(BU(n); \mathbb{Z}) \cong \Lambda_n$  and  $\xi \in H^*(K(\mathbb{Z}, 3); \mathbb{Z})$ , we have the following corollary of Proposition 4.5 by the Leibniz rule.

**Corollary 4.6** ([11, Corollary 3.10]).  ${}^U d_3(f\xi) = \nabla(f)x_1\xi$ .

In particular, the fact that  $\nabla(\sigma_k) = \sum_{i=1}^n \frac{\partial \sigma_k}{\partial v_i} = (n-k+1)\sigma_{k-1}$  shows that for  $c_k \in H^*(BU(n); \mathbb{Z}) \cong {}^U E_3^{0,*}$ ,  $1 \leq k \leq n$ , we have ([11, Corollary 3.4])

$${}^U d_3(c_k) = (n-k+1)c_{k-1}x_1. \quad (4.1)$$

**Corollary 4.7.**  ${}^U E_4^{0,*} \cong K_n$ .

*Proof.* This comes from the fact that  ${}^U E_3^{0,*} = {}^U E_2^{0,*} = H^*(BU(n); \mathbb{Z}) \cong \Lambda_n$ ,  ${}^U E_4^{0,*} = \ker {}^U d_3^{0,*}$ , where  ${}^U d_3^{0,*} = \nabla(-) \cdot x_1$  by Corollary 4.6.  $\square$

## 5. PROOF OF THEOREM 2.2

The proof of Theorem 2.2 is essentially a  $K_4$  analogue of the proof of Lemma 3.2 for  $K_3$  in [23] (also Theorem 14.2 in [24]).

Firstly, we give the explicit expression of  $\alpha_i$  in  $\Lambda_4$ . Let

$$\begin{aligned} \alpha_2 &= 8\sigma_2 - 3\sigma_1^2, \\ \alpha_3 &= 8\sigma_3 - 4\sigma_1\sigma_2 + \sigma_1^3, \\ \alpha_4 &= 12\sigma_4 - 3\sigma_1\sigma_3 + \sigma_2^2, \\ \alpha_6 &= 27\sigma_1^2\sigma_4 + 27\sigma_3^2 - 9\sigma_1\sigma_2\sigma_3 - 72\sigma_2\sigma_4 + 2\sigma_2^3. \end{aligned} \quad (5.1)$$

A direct calculation shows that  $\nabla(\alpha_i) = 0$  for all  $\alpha_i$  above, and

$$64\alpha_6 = \alpha_2^3 + 27\alpha_3^2 - 48\alpha_2\alpha_4.$$

Consider the ring homomorphism

$$\ker \nabla \cong \mathbb{Z}[v_1 - v_4, v_2 - v_4, v_3 - v_4] \hookrightarrow \mathbb{Z}[v_1, \dots, v_4] \rightarrow \mathbb{Z}[v_1, \dots, v_4]/(\sigma_1).$$

It is easy to see that this map is injective. After tensoring with  $\mathbb{Z}[\frac{1}{2}]$ , this map becomes an isomorphism, and the inverse is obtained by sending  $v_i$  to  $v_i - \frac{\sigma_1}{4}$ . The commutative diagram below shows that  $K_4 \otimes \mathbb{Z}[\frac{1}{2}] \hookrightarrow \mathbb{Z}[\frac{1}{2}][\sigma_2, \sigma_3, \sigma_4]$ .

$$\begin{array}{ccc} K_4 \otimes \mathbb{Z}[\frac{1}{2}] & \longrightarrow & \mathbb{Z}[\frac{1}{2}][\sigma_2, \sigma_3, \sigma_4] \\ \downarrow & & \downarrow \\ \ker \nabla \otimes \mathbb{Z}[\frac{1}{2}] & \xlongequal{\quad} & \mathbb{Z}[\frac{1}{2}][v_1, \dots, v_4]/(\sigma_1) \end{array}$$

On the other hand, the inclusion  $\mathbb{Z}[\frac{1}{2}][\alpha_2, \alpha_3, \alpha_4] \hookrightarrow \mathbb{Z}[\frac{1}{2}][\sigma_2, \sigma_3, \sigma_4]$  is an isomorphism, so in fact we have

$$\mathbb{Z}[\frac{1}{2}][\alpha_2, \alpha_3, \alpha_4] \cong K_4 \otimes \mathbb{Z}[\frac{1}{2}] \cong \mathbb{Z}[\frac{1}{2}][\sigma_2, \sigma_3, \sigma_4].$$

Hence if  $f \in K_4$  then  $2^s f \in \mathbb{Z}[\alpha_2, \alpha_3, \alpha_4]$  for sufficient large  $s$ , and the following lemma shows that  $\alpha_2, \alpha_3, \alpha_4, \alpha_6$  generate  $K_4$  by descending induction on  $s$ .

**Lemma 5.1.** *For  $f \in K_4$ , if  $2f \in \mathbb{Z}[\alpha_2, \alpha_3, \alpha_4, \alpha_6]$ , then  $f \in \mathbb{Z}[\alpha_2, \alpha_3, \alpha_4, \alpha_6]$  too.*

*Proof.* The images of  $\alpha_2, \alpha_3, \alpha_4, \alpha_6$  in  $\mathbb{Z}/2[\sigma_1, \dots, \sigma_4]$  is  $\sigma_1^2, \sigma_1^3, \sigma_1\sigma_3 + \sigma_2^2, \sigma_1^2\sigma_4 + \sigma_3^2 + \sigma_1\sigma_2\sigma_3$  respectively. The ideal of relations between these four polynomials is generated by  $(\sigma_1^2)^3 + (\sigma_1^3)^2$ , which is the image of  $\alpha_2^3 + 27\alpha_3^2 - 48\alpha_2\alpha_4 = 64\alpha_6$ . Hence if we write  $2f = p(\alpha_2, \alpha_3, \alpha_4, \alpha_6)$  for some integral polynomial  $p$ , then there are two integral polynomials  $q$  and  $r$  such that

$$p(\alpha_2, \alpha_3, \alpha_4, \alpha_6) = 2q(\alpha_2, \alpha_3, \alpha_4, \alpha_6) + 64\alpha_6 \cdot r(\alpha_2, \alpha_3, \alpha_4, \alpha_6).$$

Dividing by 2 we get  $f \in \mathbb{Z}[\alpha_2, \alpha_3, \alpha_4, \alpha_6]$ . □

Consider the surjective ring homomorphism  $\mathbb{Z}[y_2, y_3, y_4, y_6] \rightarrow K_4$  that sends  $y_i$  to  $\alpha_i$ . Let  $I$  be its kernel. Then the following lemma is an equivalent statement of Theorem 2.2.

**Lemma 5.2.** *In the above notation,  $I = (64y_6 - y_2^3 - 27y_3^2 + 48y_2y_4)$ .*

*Proof.* Let  $\mathbf{y} = 64y_6 - y_2^3 - 27y_3^2 + 48y_2y_4$ . We know that  $\mathbf{y} \in I$ . Note that

$$\mathbb{Z}[\frac{1}{2}][y_2, y_3, y_4, y_6]/(\mathbf{y}) = \mathbb{Z}[\frac{1}{2}][y_2, y_3, y_4],$$

which is isomorphic to  $K_4 \otimes \mathbb{Z}[\frac{1}{2}]$  as we saw above. This implies that if  $f \in I$ , then some multiple  $2^r f$  is in  $(\mathbf{y})$ . However, since 2 dose not divide  $\mathbf{y}$  in the UFD  $\mathbb{Z}[y_2, y_3, y_4, y_6]$ ,  $f$  is actually in  $(\mathbf{y})$ , and we get the desired equality in the lemma. □

We end this section with an algebraic result, which will be used in Section 7.

**Proposition 5.3.**  $\mathbb{Z}/4[\alpha_4, \alpha_6]$  is a subring of  $\text{coker } \nabla$ , where  $\nabla$  acts on  $\Lambda_4$ .

*Proof.* Since  $\alpha_4, \alpha_6 \in K_4$  and  $\nabla(\sigma_1) = 4$ ,  $\nabla(\sigma_1 f) = 4f$  for any  $f \in \mathbb{Z}[\alpha_4, \alpha_6]$ . So the order of  $f$  in  $\text{coker } \nabla$  divides 4. If its order is not 4, there would exist  $g \in \Lambda_4$  such that  $\nabla(g) = 2f$ , and then  $\sigma_1 f - 2g \in K_4$ . Let  $\rho : \Lambda_4 \rightarrow \Lambda_4 \otimes \mathbb{Z}/2$  be the mod 2 map, and write  $\tilde{h} = \rho(h)$  for an element  $h \in \Lambda_4$ . Then  $\rho(\sigma_1 f - 2g) = \sigma_1 \tilde{f} \in K_4 \otimes \mathbb{Z}/2 = \mathbb{Z}/2[\tilde{\alpha}_2, \tilde{\alpha}_3, \tilde{\alpha}_4, \tilde{\alpha}_6]$ , where  $\tilde{\alpha}_2 = \sigma_1^2$ ,  $\tilde{\alpha}_3 = \sigma_1^3$ ,  $\tilde{\alpha}_4 = \sigma_1 \sigma_3 + \sigma_2^2$ ,  $\tilde{\alpha}_6 = \sigma_1^2 \sigma_4 + \sigma_3^2 + \sigma_1 \sigma_2 \sigma_3$ . This means that there is a polynomial  $p$  over  $\mathbb{Z}/2$  such that  $\sigma_1 \tilde{f} = p(\tilde{\alpha}_2, \tilde{\alpha}_3, \tilde{\alpha}_4, \tilde{\alpha}_6)$  in  $\mathbb{Z}/2[\sigma_1, \dots, \sigma_4]$ . Suppose that the leading monomial terms of  $\tilde{f}$  and  $p$  are  $\mathbf{m}$  and  $\mathbf{n}$  respectively, with respect to an arbitrary term order. Then  $\mathbf{n} = \sigma_1 \mathbf{m}$ , and so  $\deg(\mathbf{n}) = \deg(\mathbf{m}) + 1$ , an odd number, it follows that  $\tilde{\alpha}_3$  divides  $\mathbf{n}$ . However, since  $\mathbb{Z}/2[\sigma_1, \dots, \sigma_4]$  is a UFD and  $\sigma_1$  does not divide  $\tilde{\alpha}_4$  and  $\tilde{\alpha}_6$ ,  $\tilde{\alpha}_3 = \sigma_1^3$  can not divide  $\sigma_1 \mathbf{m}$ , where  $\mathbf{m} = \tilde{\alpha}_4^i \tilde{\alpha}_6^j$  for some  $i, j \geq 0$ . This gives a contradiction. So the order of  $f$  must be 4, and the proposition follows immediately.  $\square$

## 6. ON THE HOMOMORPHISM FROM $H^*(K(\mathbb{Z}, 3))$ TO $H^*(BPU(2))$

In this section, we study the homomorphism  $\chi^* : H^*(K(\mathbb{Z}, 3)) \rightarrow H^*(BPU(2))$  induced by the fibration map  $\chi : BPU(2) \rightarrow K(\mathbb{Z}, 3)$  defined in (3.2). Throughout this section, let  $\rho : H^*(-; \mathbb{Z}) \rightarrow H^*(-; \mathbb{Z}/2)$  denote the mod 2 reduction map on cohomology.

Since  $PU(2) = SO(3)$ , the following result is a special case of [14].

**Theorem 6.1.** Let  $w_i \in H^i(BSO(3); \mathbb{Z}/2)$  be the  $i$ th Stiefel-Whitney class of  $BSO(3)$ , and  $p_i \in H^{4i}(BSO(3); \mathbb{Z})$  the  $i$ th Pontryagin class. Then

$$\begin{aligned} H^*(BSO(3); \mathbb{Z}/2) &= \mathbb{Z}/2[w_2, w_3], \quad w_3 = Sq^1(w_2), \quad \text{and} \\ H^*(BSO(3); \mathbb{Z}) &= \mathbb{Z}[p_1, W_3]/(2W_3), \quad \rho(p_1) = w_2^2, \quad \rho(W_3) = w_3. \end{aligned}$$

Here  $W_3$  is the degree-3 integral Stiefel-Whitney class.

**Theorem 6.2.** In the notation of Proposition 4.3, the mod 2 cohomology homomorphism  $\chi^* : H^*(K(\mathbb{Z}, 3); \mathbb{Z}/2) \rightarrow H^*(BPU(2); \mathbb{Z}/2)$  satisfies

$$\begin{aligned} \chi^*(x_1) &= w_3, \quad \chi^*(x_{2,0}) = w_2 w_3, \\ \chi^*(x_{2,k}) &\equiv w_2^{2k+1-1} w_3 \pmod{(w_3^3)}, \quad \text{for } k \geq 1. \end{aligned}$$

*Proof.* First we show that  $\chi^*(x_1) = w_3$ . This follows from the computation in [11] that  $H^3(BPU(n); \mathbb{Z}) = \mathbb{Z}/n$  is generated by  $\chi^*(x_1)$ . Here we give a short explanation to this. Since  ${}^U E_2^{0,*} = {}^U E_3^{0,*} = H^*(BU(n); \mathbb{Z}) \cong \mathbb{Z}[c_1, \dots, c_n]$  and

$U d_3^{0,*}$  is  $\nabla(-) \cdot x_1$ , it follows that  $U d_3^{0,2}(c_1) = n x_1 \in U E_3^{3,0}$ . Therefore, for degree reasons

$$U E_4^{3,0} = U E_\infty^{3,0} = H^3(BPU(n); \mathbb{Z}) = \mathbb{Z}/n\{\chi^*(x_1)\}.$$

Now we give the formula for  $\chi^*(x_{2,k})$  by induction on  $k$ , starting with  $\chi^*(x_{2,0})$ . Since  $Sq^1(x_{2,0}) = x_1^2$  by the differential rule of  $C(3)$ , we have  $\chi^* Sq^1(x_{2,0}) = Sq^1 \chi^*(x_{2,0}) = w_3^2$ . Hence  $\chi^*(x_{2,0}) = w_2 w_3$ , the only nonzero element of degree 5. Inductively, suppose that  $\chi^*(x_{2,k-1})$  is congruent to  $w_2^{2^k-1} w_3$  modulo the ideal generated by  $w_3^3$ . Then since  $Sq^1(x_{2,k}) = x_{2,k-1}^2$ , we have

$$Sq^1 \chi^*(x_{2,k}) = \chi^* Sq^1(x_{2,k}) \equiv w_2^{2^{k+1}-2} w_3^2 \pmod{(w_3^3)}.$$

Since in  $H^*(BSO(3); \mathbb{Z}/2)/(w_3^3)$ ,  $w_2^{2^{k+1}-1} w_3$  is the only degree- $(2^{k+2} + 1)$  element mapped by  $Sq^1$  to  $w_2^{2^{k+1}-2} w_3^2$ , it follows that  $\chi^*(x_{2,k}) \equiv w_2^{2^{k+1}-1} w_3 \pmod{(w_3^3)}$ , and the induction step is finished.  $\square$

**Corollary 6.3.** *In the notation of Theorem 4.1 and Theorem 6.1, the homomorphism  $\chi^* : H^*(K(\mathbb{Z}, 3); \mathbb{Z}) \rightarrow H^*(BPU(2); \mathbb{Z})$  satisfies  $\chi^*(x_1) = W_3$ , and  $\chi^*(y_{2,k}) \equiv p_1^{2^k-1} W_3^2 \pmod{(W_3^6)}$  for  $k \geq 0$ .*

*Proof.* The formula for  $\chi^*(x_1)$  is obvious. From Section 4, we know that  $\rho(y_{2,0}) = x_1^2$  and  $\rho(y_{2,k}) = x_{2,k-1}^2$  for  $k \geq 1$ . Then, by Theorem 6.2 we have  $\rho \chi^*(y_{2,k}) = \chi^* \rho(y_{2,k}) \equiv w_2^{2^{k+1}-2} w_3^2 \pmod{(w_3^6)}$  for  $k \geq 0$ . This implies that  $\chi^*(y_{2,k}) \equiv p_1^{2^k-1} W_3^2 \pmod{(W_3^6)}$  since  $p_1^{2^k-1} W_3^2$  is the only torsion element in  $H^*(BPU(2); \mathbb{Z})/(W_3^6)$  mapped by  $\rho$  to  $w_2^{2^{k+1}-2} w_3^2 \pmod{(w_3^6)}$ .  $\square$

**Proposition 6.4.** *Let  $n > 0$  be an interger such that  $2 \mid n$ . Then for any  $i, j, k \geq 0$ ,*

- (a)  $x_1^i x_{2,k}^j \neq 0$  in  $H^*(BPU(n); \mathbb{Z}/2)$ .
- (b)  $x_1^i y_{2,k}^j \neq 0$  in  $H^*(BPU(n); \mathbb{Z})$ .

*Here we abuse the notations  $x_1, x_{2,k}, y_{2,k}$ , to denote their images under the cohomology ring homomorphism induced by the fibration map  $BPU(n) \rightarrow K(\mathbb{Z}, 3)$ .*

*Proof.* Since  $2 \mid n$ , there is a diagonal map of matrices

$$U(2) \rightarrow U(n), \quad A \mapsto \begin{bmatrix} A & 0 & \cdots & 0 \\ 0 & A & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & \cdots & A \end{bmatrix},$$

which passes to  $PU(2) \rightarrow PU(n)$ . These diagonal maps induce maps  $\Delta : BU(2) \rightarrow BU(n)$  and  $\bar{\Delta} : BPU(2) \rightarrow BPU(n)$ , and a commutative diagram of fibrations

$$\begin{array}{ccccc} BU(2) & \longrightarrow & BPU(2) & \longrightarrow & K(\mathbb{Z}, 3) \\ \downarrow \Delta & & \downarrow \bar{\Delta} & & \downarrow = \\ BU(n) & \longrightarrow & BPU(n) & \longrightarrow & K(\mathbb{Z}, 3) \end{array} \quad (6.1)$$

Since the map  $\bar{\Delta}^*$  takes the elements  $x_1, x_{2,k}, y_{2,k}$ , to themselves, the corollary follows from Theorem 6.2 and Corollary 6.3.  $\square$

## 7. ON THE SPECTRAL SEQUENCE ${}^U E$

**Proposition 7.1.** *In the integral spectral sequence  ${}^U E$  associated to  $BU(4) \xrightarrow{\pi} BPU(4) \xrightarrow{\chi} K(\mathbb{Z}, 3)$ , we have  $K_4 \cong {}^U E_4^{0,*} = {}^U E_\infty^{0,*}$ . Moreover,  $\mathbb{Z}/4[\alpha_4, \alpha_6]$  is a submodule of  ${}^U E_\infty^{3,*}$ .*

*Proof.* By Corollary 4.7,  $K_4 \cong {}^U E_4^{0,*}$ . Since the differentials in the spectral sequence  ${}^U E$  satisfy the Leibniz rule and  $K_4$  is generated by  $\alpha_2, \alpha_3, \alpha_4, \alpha_6$ , it suffices to show that these generators survive to  ${}^U E_\infty$ . For  $\alpha_2, \alpha_3, \alpha_4$ , this was proved in [11], so we only consider  $\alpha_6 \in {}^U E_4^{0,12}$ .

The differentials in  ${}^U E$  have the form  $d_r : {}^U E_r^{s,t} \rightarrow {}^U E_r^{s+r,t-r+1}$ . Hence from Theorem 4.1 we see that the only possible nontrivial differentials  $d_r$ ,  $r \geq 4$ , originating at  ${}^U E_*^{0,12}$  are  $d_9 : {}^U E_9^{0,12} \rightarrow {}^U E_9^{9,4}$ ,  $d_{11} : {}^U E_{11}^{0,12} \rightarrow {}^U E_{11}^{11,2}$  and  $d_{13} : {}^U E_{13}^{0,12} \rightarrow {}^U E_{13}^{13,0}$ . Note that  ${}^U E_2^{9,4} \cong x_1^3 \cdot H^4(BU(4); \mathbb{Z})$ ,  ${}^U E_2^{11,2} \cong \mathbb{Z}/3\{x_1 x_{3,0} c_1\}$  and  ${}^U E_2^{13,0} \cong \mathbb{Z}/2\{x_1 y_{2,1}\}$  by Theorem 4.1.

From Proposition 6.4 we know that  $x_1 y_{2,1}$  is a permanent cycle, so it can not be hit by any differential in  ${}^U E$ . Furthermore, from (4.1) we see that  $d_3(x_{3,0} c_1^2) \equiv -x_1 x_{3,0} c_1 \pmod{3}$ , so  ${}^U E_4^{11,2} = 0$ . Hence the proposition will follow once we prove that  ${}^U E_4^{9,4} = 0$ . This is an easy calculation using the following commutative diagram

$$\begin{array}{ccccc} \Lambda_4^3 \otimes \mathbb{Z}/2 & \xrightarrow{\nabla_{\mathbb{Z}/2}} & \Lambda_4^2 \otimes \mathbb{Z}/2 & \xrightarrow{\nabla_{\mathbb{Z}/2}} & \Lambda_4^1 \otimes \mathbb{Z}/2 \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ {}^U E_3^{6,6} & \xrightarrow{d_3^{6,6}} & {}^U E_3^{9,4} & \xrightarrow{d_3^{9,4}} & {}^U E_3^{12,2} \end{array}$$

The right vertical map is only an injection since besides  $x_1^4 \cdot H^2(BU(4); \mathbb{Z})$ ,  $y_{5,0} \cdot H^2(BU(4); \mathbb{Z})$  also contributes to  ${}^U E_3^{12,2}$ , but this does not effect our computation. It can be shown that  $\ker d_3^{9,4} = \mathbb{Z}/2\{x_1^3 c_1^2\}$  and  $d_3^{6,6}(x_1^2 c_2 c_1) = x_1^3 c_1^2$ . Then we have  ${}^U E_4^{9,4} = \ker d_3^{9,4} / \text{Im } d_3^{6,6} = 0$ .

Using Proposition 5.3, we see that  $x_1 f$  in  ${}^U E_4^{3,*}$  has order 4 for any  $f \in \mathbb{Z}[\alpha_4, \alpha_6] \subset H^*(BU(4); \mathbb{Z})$ . Since  $\alpha_4, \alpha_6$  and  $x_1$  are all permanent cycles, so is  $x_1 f$ , and the second statement follows.  $\square$

Proposition 7.1 says that there exist elements  $\bar{\alpha}_i \in H^{2i}(BPU(4); \mathbb{Z})$ ,  $i = 2, 3, 4, 6$ , such that  $\pi^*(\bar{\alpha}_i) = \alpha_i \in H^{2i}(BU(4); \mathbb{Z})$ . Let  $\Delta : BU(2) \rightarrow BU(4)$ ,  $\bar{\Delta} : BPU(2) \rightarrow BPU(4)$  be the maps defined in the proof of Proposition 6.4. We consider the images of  $\alpha_i$  and  $\bar{\alpha}_i$  under the maps  $\Delta^*$  and  $\bar{\Delta}^*$  respectively.

Let  $c_i$  and  $c'_i$  be the  $i$ th universal Chern classes of  $BU(4)$  and  $BU(2)$  respectively. Since the map  $\Delta$  factors through the diagonal map  $BU(2) \rightarrow BU(2) \times BU(2)$ , using the Whitney sum formula and the functorial property of total Chern classes we immediately get

$$\Delta^*(1 + c_1 + c_2 + c_3 + c_4) = (1 + c'_1 + c'_2)^2.$$

In other words,

$$\begin{aligned} \Delta^*(c_1) &= 2c'_1, & \Delta^*(c_2) &= 2c'_2 c_1'^2, \\ \Delta^*(c_3) &= 2c'_1 c'_2, & \Delta^*(c_4) &= c_2'^2. \end{aligned} \tag{7.1}$$

**Proposition 7.2.** *The elements  $\bar{\alpha}_i \in H^{2i}(BPU(4); \mathbb{Z})$ ,  $i = 2, 3, 4, 6$ , can be chosen such that  $\bar{\Delta}^*(\bar{\alpha}_4) = p_1^2$ , and  $\bar{\Delta}^*(\bar{\alpha}_i) \equiv 0 \pmod{2}$  for  $i = 2, 3, 6$ .*

*Proof.* It follows from (5.1) and (7.1) that  $\Delta^*(\alpha_i) \equiv 0 \pmod{2}$  for  $i = 2, 3, 6$ , and  $\Delta^*(\alpha_4) = (c_1'^2 - 4c_2')^2$ . On the other hand, for the spectral sequence  ${}^U E$  associated to  $BU(2) \xrightarrow{\pi} BPU(2) \xrightarrow{\chi} K(\mathbb{Z}, 3)$ , an easy calculation shows that  ${}^U E_\infty^{0,4} = {}^U E_4^{0,4} \cong K_2 = \mathbb{Z}[\sigma_1^2 - 4\sigma_2] \subset \Lambda_2$ . This means that for  $p_1 \in H^4(BSO(3); \mathbb{Z}) = H^4(BPU(2); \mathbb{Z})$ ,  $\pi^*(p_1) = \pm(c_1'^2 - 4c_2')$ .

Recall from Theorem 6.1 that

$$\begin{aligned} H^4(BPU(2); \mathbb{Z}) &\cong \mathbb{Z}\{p_1\}, & H^6(BPU(2); \mathbb{Z}) &\cong \mathbb{Z}/2\{W_3^3\}, \\ H^8(BPU(2); \mathbb{Z}) &\cong \mathbb{Z}\{p_1^2\}, & H^{12}(BPU(2); \mathbb{Z}) &\cong \mathbb{Z}\{p_1^3\} \oplus \mathbb{Z}/2\{W_3^4\}. \end{aligned}$$

Since  $\pi^* \bar{\Delta}^* = \Delta^* \pi^*$ , combining the above computations, we obtain  $\bar{\Delta}^*(\bar{\alpha}_4) = p_1^2$  and  $\bar{\Delta}^*(\bar{\alpha}_2) \equiv 0 \pmod{2}$ . We have seen that  $\bar{\Delta}^*(x_1) = W_3$ , hence by adding  $x_1^2$  to  $\bar{\alpha}_3$  or  $x_1^4$  to  $\bar{\alpha}_6$  if necessary, we can also obtain  $\bar{\Delta}^*(\bar{\alpha}_3), \bar{\Delta}^*(\bar{\alpha}_6) \equiv 0 \pmod{2}$ .  $\square$

## 8. ON $H^*(BPU(4); \mathbb{Z}/2)$

In this section we prove some facts about the generators  $y_i$  of  $H^*(BPU(4); \mathbb{Z}/2)$  in Theorem 2.4.

**Proposition 8.1.** *In the notation of Proposition 4.3,  $\chi^*(x_1) = y_3$ ,  $\chi^*(x_{2,0}) = y_5$ ,  $\chi^*(x_{2,1}) = y_9$  or  $y_3^3 + y_9$ .*

*Proof.* From Proposition 6.4 we know that  $\chi^*(x_1), \chi^*(x_{2,0}), \chi^*(x_{2,1}) \neq 0$ . Since  $\deg(x_1) = 3$  and  $y_3$  is the only nonzero element in  $H^*(BPU(4); \mathbb{Z}/2)$ ,  $\chi^*(x_1) = y_3$ . Similarly,  $\chi^*(x_{2,0}) = y_5$ .

From Theorem 2.4 one easily sees that  $H^9(BPU(4); \mathbb{Z}/2) = \mathbb{Z}/2\{y_3^3, y_9\}$ . Since  $Sq^1(x_{2,1}) = x_{2,0}^2$  by Proposition 4.3, it follows that  $Sq^1\chi^*(x_{2,1}) = \chi^*(x_{2,0}^2) = y_5^2$ . Hence  $\chi^*(x_{2,1})$  takes one of the forms in the proposition since  $Sq^1(y_3) = Sq^1\chi^*(x_1) = \chi^*Sq^1(x_1) = 0$ .  $\square$

Recall from Proposition 4.3 that  $Sq^1(x_1) = 0$ ,  $Sq^1(x_{2,0}) = x_1^2$ ,  $Sq^1(x_{2,1}) = x_{2,0}^2$ ,  $Sq^2(x_1) = x_{2,0}$ , and  $Sq^4(x_{2,0}) = x_{2,1}$ . Then we have the following corollary of Proposition 8.1.

**Corollary 8.2.** *In  $H^*(BPU(4); \mathbb{Z}/2)$ ,  $Sq^1(y_3) = 0$ ,  $Sq^1(y_5) = y_3^2$ ,  $Sq^1(y_9) = y_5^2$ ,  $Sq^2(y_3) = y_5$ ,  $Sq^4(y_5) = y_9$  or  $y_3^3 + y_9$ .*

Before proceeding further, let us review a result in [21]. Let  $PSO(2n)$  be the quotient group of  $SO(2n)$  by the center  $\mathbb{Z}/2 = \{\pm 1\}$ , and let  $\phi : BSO(2n) \rightarrow BPSO(2n)$  be the induced map of classifying spaces. It is known that  $PU(4)$  is homeomorphic to  $PSO(6)$ , so there is an induced homomorphism

$$\phi^* : H^*(BPU(4); \mathbb{Z}/2) \rightarrow H^*(BSO(6); \mathbb{Z}/2) \cong \mathbb{Z}[w_2, \dots, w_6],$$

where  $w_i$  is the  $i$ th Stiefel-Whitney class of  $BSO(2n)$ .

**Proposition 8.3** ([21, Proposition 3.7 and 4.5]).  $\phi^*(y_2) = 0$ ,  $\phi^*(y_3) = w_3$ ,  $\phi^*(y_5) = \bar{w}_5$ ,  $\phi^*(y_8) = \bar{w}_4^2 + w_3^2 w_2$ ,  $\phi^*(y_{12}) = \bar{w}_6^2 + \bar{w}_5^2 w_2$ , where  $\bar{w}_4 = w_4 + w_2^2$ ,  $\bar{w}_5 = w_5 + w_2 w_3$ ,  $\bar{w}_6 = w_6 + w_2 w_4$ , and  $\phi^*(y_9) \equiv w_3 w_6 + w_4 w_5 \pmod{(w_2)}$ .

(See [21, p. 91] for the formulas of  $\bar{w}_i$ ,  $i = 4, 5, 6$ .)

**Proposition 8.4.** *In Proposition 8.3,  $\phi^*(y_9) = w_3 w_6 + w_4 w_5 + w_2^2 w_5 + w_3^2 w_3$ . Moreover,  $Sq^4(y_5)$ , i.e.  $\chi^*(x_{2,1})$ , is actually  $y_3^3 + y_9$  in Corollary 8.2.*

*Proof.* Applying the Wu formula on Stiefel-Whitney classes:

$$Sq^i(w_k) = \sum_{j=0}^i \binom{k-j-1}{i-j} w_{k+i-j} w_j,$$

and the Cartan formula on Steenrod squares we get

$$\begin{aligned} Sq^4(\bar{w}_5) &= Sq^4(w_5) + Sq^4(w_2 w_3) \\ &= w_3 w_6 + w_4 w_5 + Sq^1(w_2) Sq^3(w_3) + Sq^2(w_2) Sq^2(w_3) \\ &= w_3 w_6 + w_4 w_5 + w_3^3 + w_2^2 w_5 + w_3^2 w_3. \end{aligned}$$

By Corollary 8.2 and Proposition 8.3,  $Sq^4(\bar{w}_5) = Sq^4(\phi^*(y_5)) = \phi^* Sq^4(y_5) = \phi^*(y_9)$  or  $\phi^*(y_9) + w_3^3$ , where  $\phi^*(y_9) \equiv w_3 w_6 + w_4 w_5 \pmod{(w_2)}$ . Combining this with the

above expression for  $Sq^4(\bar{w}_5)$ , we obtain  $\phi^*(y_9) = w_3w_6 + w_4w_5 + w_2^2w_5 + w_2^3w_3$ . This computation also shows that  $Sq^4(y_5) = y_3^3 + y_9$ , which is the second statement.  $\square$

Let  $\pi : BU(4) \rightarrow BPU(4)$  be the map in (3.1). The following result in [21] gives the induced mod 2 cohomology ring homomorphism  $\pi^*$ .

**Proposition 8.5** ([21, (4.11)]). *In  $H^*(BU(4); \mathbb{Z}/2)$ ,  $\pi^*(y_2) = c_1$ ,  $\pi^*(y_8) = c_1c_3 + c_2^2$ ,  $\pi^*(y_{12}) = c_1^2c_4 + c_3^2 + c_1c_2c_3$ .*

Let  $\Delta : BU(2) \rightarrow BU(4)$  and  $\bar{\Delta} : BSO(3) \cong BPU(2) \rightarrow BPU(4)$  be the maps defined in the proof of Proposition 6.4, and to avoid ambiguity of notations, let  $w'_i$  denote the  $i$ th Stiefel-Whitney class of  $BSO(3)$ .

**Proposition 8.6.**  *$Sq^1(y_2) = 0$ ,  $Sq^1(y_8) = y_3^3$ ,  $Sq^1(y_{12}) = y_3y_5^2$ , and  $\bar{\Delta}^*(y_2) = 0$ ,  $\bar{\Delta}^*(y_3) = w'_3$ ,  $\bar{\Delta}^*(y_5) = w'_2w'_3$ ,  $\bar{\Delta}^*(y_8) = w_2^4 + w_2^2w_3^2$ ,  $\bar{\Delta}^*(y_9) = w_2^3w_3$ ,  $\bar{\Delta}^*(y_{12}) = w_2^3w_3^2$ .*

*Proof.* By Theorem 6.2,  $\bar{\Delta}^*(y_3) = \bar{\Delta}^*\chi^*(x_1) = w'_3$ . Then the equation  $Sq^1(y_2) = 0$  follows since  $H^3(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_3\}$  and  $\phi^*Sq^1(y_2) = Sq^1\phi^*(y_2) = 0 \neq w_3 = \phi^*(y_3)$ . Since  $H^2(BPU(2); \mathbb{Z}/2) \cong \mathbb{Z}/2\{w'_2\}$  and  $\pi^*(w'_2) = c_1$ , the formula  $\bar{\Delta}^*(y_2) = 0$  follows from the fact that  $\pi^*\bar{\Delta}^*(y_2) = \Delta^*\pi^*(y_2) = \Delta^*(c_1) = 0$  by (7.1).

$\bar{\Delta}^*(y_5) = \bar{\Delta}^*\chi^*(x_{2,1}) = w'_2w'_3$  also comes from Theorem 6.2, and  $\bar{\Delta}^*(y_9) = \bar{\Delta}^*(Sq^4(y_5) + y_3^3) = w_2^3w_3$  comes from the the second statement of Proposition 8.4 and the fact that  $\bar{\Delta}^*Sq^4(y_5) = Sq^4\bar{\Delta}^*(y_5) = Sq^4(w'_2w'_3) = w_2^3w_3 + w_3^3$ .

We see from Proposition 8.3 that  $Sq^1\phi^*(y_8) = w_3^3 = \phi^*(y_3^3)$ . Hence  $Sq^1(y_8) = y_3^3$  since  $H^9(BPU(4); \mathbb{Z}/2) = \mathbb{Z}/2\{y_3^3, y_9\}$  and  $\phi^*(y_9) \neq 0$ . Recall from Theorem 6.1 that  $H^8(BPU(2); \mathbb{Z}/2) \cong \mathbb{Z}/2\{w_2^4, w_2^2w_3^2\}$ . Hence  $\bar{\Delta}^*(y_8) = w_2^4 + aw_2^2w_3^2$ ,  $a \in \mathbb{Z}/2$ , since  $Sq^1\bar{\Delta}^*(y_8) = \bar{\Delta}^*Sq^1(y_8) = w_3^3$ . Furthermore, by Proposition 8.5,  $\pi^*(y_8) = c_1c_3 + c_2^2 \in H^*(BU(4); \mathbb{Z}/2)$ , and then  $\pi^*\bar{\Delta}^*(y_8) = \Delta^*\pi^*(y_8) = c_1^4$  by (7.1), which implies that the coefficient  $a$  is 1 since  $\pi^*(w_3^3) = 0$ .

Similarly, we have  $Sq^1(y_{12}) = ay_3y_5^2 + by_5y_8 \neq 0$  in  $H^{13}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_3y_5^2, y_5y_8\}$ . Since  $Sq^1(y_3y_5^2) = 0$  and  $Sq^1(y_5y_8) = y_3^2y_8 + y_5y_3^3 \neq 0$ , it follows that  $Sq^1(y_{12}) = y_3y_5^2$ . The relation  $y_9^2 + y_3^2y_{12} + y_5^2y_8 = 0$  in  $H^*(BPU(4); \mathbb{Z}/2)$  shows that

$$\begin{aligned} & \bar{\Delta}^*(y_9^2 + y_3^2y_{12} + y_5^2y_8) \\ &= (w_2^3w_3)^2 + w_3^2\bar{\Delta}^*(y_{12}) + (w_2^2w_3)^2(w_2^4 + w_2^2w_3^2) = 0. \end{aligned}$$

Solving this equation we obtain  $\bar{\Delta}^*(y_{12}) = w_2^3w_3^2$ .  $\square$

Let  $\alpha_i \in H^*(BU(4); \mathbb{Z})$  and  $\bar{\alpha}_i \in H^*(BPU(4); \mathbb{Z})$ ,  $i = 2, 3, 4, 6$ , be as in Section 7, and let  $\rho : H^*(-; \mathbb{Z}) \rightarrow H^*(-; \mathbb{Z}/2)$  denote the mod 2 reduction map.

**Proposition 8.7.**  $\rho(\bar{\alpha}_2) = y_2^2$ ,  $\rho(\bar{\alpha}_3) = y_2^3$ ,  $\rho(\bar{\alpha}_4) = y_8 + y_3y_5$ ,  $\rho(\bar{\alpha}_6) = y_{12} + y_3y_9$ .

*Proof.* By (5.1),  $\alpha_2 \equiv c_1^2$ ,  $\alpha_3 \equiv c_1^3$ ,  $\alpha_4 \equiv c_1c_3 + c_2^2$ ,  $\alpha_6 \equiv c_1^2c_4 + c_3^2 + c_1c_2c_3 \pmod{2}$ . By Proposition 8.5, this means that  $\pi^*\rho(\bar{\alpha}_2) = \rho(\alpha_2) = \pi^*(y_2^2)$ ,  $\pi^*\rho(\bar{\alpha}_3) = \rho(\alpha_3) = \pi^*(y_2^3)$ ,  $\pi^*\rho(\bar{\alpha}_4) = \rho(\alpha_4) = \pi^*(y_8)$ ,  $\pi^*\rho(\bar{\alpha}_6) = \rho(\alpha_6) = \pi^*(y_{12})$  in  $H^*(BU(4); \mathbb{Z}/2)$ .

$\rho(\bar{\alpha}_2) = y_2^2$  is obvious since  $y_2^2$  is the only nonzero element of  $H^4(BPU(4); \mathbb{Z}/2)$ .  $\rho(\bar{\alpha}_3) = y_2^3$  follows from the facts that  $\bar{\Delta}^*\rho(\bar{\alpha}_3) = 0$  by Proposition 7.2 and that  $H^6(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^3, y_3^2\}$ ,  $\bar{\Delta}^*(y_2) = 0$ ,  $\bar{\Delta}^*(y_3^2) = (w_3')^2 \neq 0$  by Proposition 8.6.

Proposition 7.2 also gives  $\bar{\Delta}^*\rho(\bar{\alpha}_4) = w_2'^4$  because  $\rho(p_1) = w_2'^2$ . Notice that  $H^8(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^4, y_3y_5, y_8\}$ . Hence from  $\pi^*(y_2) = c_1$ , and  $\bar{\Delta}^*(y_2) = 0$ ,  $\bar{\Delta}^*(y_3y_5) = w_2'w_3'^2$ ,  $\bar{\Delta}^*(y_8) = w_2'^4 + w_2'w_3'^2$  by Proposition 8.6, it follows that  $\rho(\bar{\alpha}_4) = y_8 + y_3y_5$ .

The fact that  $\pi^*\rho(\bar{\alpha}_6) = \pi^*(y_{12}) = c_1^2c_4 + c_3^2 + c_1c_2c_3$  shows that  $\rho(\bar{\alpha}_6) = y_{12} + ay_3y_9 + by_3^4$ ,  $a, b \in \mathbb{Z}/2$ , since  $H^{12}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^6, y_3^4, y_3y_9, y_{12}\}$  and  $\pi^*(y_2^6) = c_1^6$ . Also, we have  $\bar{\Delta}^*\rho(\bar{\alpha}_6) = 0$  by Proposition 7.2, and  $\bar{\Delta}^*(y_3^4) = w_3'^4$ ,  $\bar{\Delta}^*(y_3y_9) = w_2'w_3'^2$ ,  $\bar{\Delta}^*(y_{12}) = w_2'^3w_3'^2$  by Proposition 8.6. These together give the coefficients  $a = 1$  and  $b = 0$ . The proof is completed.  $\square$

## 9. PROOF OF THEOREM 2.5

The strategy of the proof is to compute the action of Steenrod squares on the images of the generators  $y_i$  under the map  $\pi^* \oplus \phi^* \oplus \bar{\Delta}^*$ :

$$H^*(BPU(4); \mathbb{Z}/2) \rightarrow H^*(BU(4); \mathbb{Z}/2) \oplus H^*(BSO(6); \mathbb{Z}/2) \oplus H^*(BPU(2); \mathbb{Z}/2).$$

As in Section 8, we use  $w_i'$  and  $w_i$  to denote the  $i$ th Stiefel-Whitney class of  $BSO(3)$  and  $BSO(6)$  respectively.

First we consider  $Sq^i$  on  $\text{Im } \pi^*$ .

**Proposition 9.1.**  $Sq^2\pi^*(y_8) = 0$ ,  $Sq^2\pi^*(y_{12}) = \pi^*(y_2y_{12})$ ,  $Sq^4\pi^*(y_8) = \pi^*(y_2^2y_8 + y_{12})$ ,  $Sq^4\pi^*(y_{12}) = \pi^*(y_2^2y_{12})$ ,  $Sq^8\pi^*(y_{12}) = \pi^*(y_8y_{12})$ .

*Proof.* Recall that  $\pi^*(y_2) = c_1$ ,  $\pi^*(y_8) = c_1c_3 + c_2^2$  and  $\pi^*(y_{12}) = c_1^2c_4 + c_3^2 + c_1c_2c_3$ . Then the calculation can be carried out by using the Cartan formula on Steenrod squares and the Wu formula on the mod 2 Chern classes:

$$Sq^{2i}(c_k) = \sum_{j=0}^i \binom{k-j-1}{i-j} c_{k+i-j}c_j.$$

For  $H^*(BU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2[c_1, \dots, c_4]$ , we have  $Sq^2(c_1) = c_1^2$ ,  $Sq^2(c_2) = c_1c_2 + c_3$ ,  $Sq^2(c_3) = c_1c_3$ ,  $Sq^2(c_4) = c_1c_4$ ,  $Sq^4(c_2) = c_2^2$ ,  $Sq^4(c_3) = c_1c_4 + c_2c_3$ ,  $Sq^4(c_4) = c_2c_4$ ,  $Sq^8(c_4) = c_4^2$ . Thus,

$$Sq^2\pi^*(y_8) = Sq^2(c_1)c_3 + c_1Sq^2(c_3) = c_1^2c_3 + c_1^2c_3 = 0,$$

$$\begin{aligned} Sq^2\pi^*(y_{12}) &= c_1^2Sq^2(c_4) + Sq^2(c_1)c_2c_3 + c_1Sq^2(c_2)c_3 + c_1c_2Sq^2(c_3) \\ &= c_1^3c_4 + c_1^2c_2c_3 + c_1(c_1c_2 + c_3)c_3 + c_1^2c_2c_3 = \pi^*(y_2y_{12}), \end{aligned}$$

$$\begin{aligned} Sq^4\pi^*(y_8) &= Sq^2(c_1)Sq^2(c_3) + c_1Sq^4(c_3) + [Sq^2(c_2)]^2 \\ &= c_1^3c_3 + c_1(c_1c_4 + c_2c_3) + (c_1c_2 + c_3)^2 = \pi^*(y_2^2y_8 + y_{12}), \end{aligned}$$

$$\begin{aligned} Sq^4\pi^*(y_{12}) &= Sq^4(c_1^2)c_4 + c_1^2Sq^4(c_4) + [Sq^2(c_3)]^2 + Sq^4(c_1 \cdot c_2c_3) \\ &= c_1^4c_4 + c_1^2c_2c_4 + c_1^2c_3^2 + [Sq^2(c_1)Sq^2(c_2c_3) + c_1Sq^4(c_2c_3)] \\ &= c_1^4c_4 + c_1^2c_2c_4 + c_1^2c_3^2 + [c_1^2c_3^2 + c_1(c_1^2c_2c_3 + c_1c_3^2 + c_1c_2c_4)] \\ &= c_1^4c_4 + c_1^2c_3^2 + c_1^3c_2c_3 = \pi^*(y_2^2y_{12}), \end{aligned}$$

$$\begin{aligned} Sq^8\pi^*(y_{12}) &= c_1^2Sq^8(c_4) + Sq^4(c_1^2)Sq^4(c_4) + [Sq^4(c_3)]^2 + Sq^8(c_1c_2 \cdot c_3) \\ &= c_1^2c_4^2 + c_1^4c_2c_4 + (c_1c_4 + c_2c_3)^2 + Sq^8(c_1c_2 \cdot c_3) = \pi^*(y_8y_{12}). \end{aligned}$$

In the last equation, we use

$$\begin{aligned} Sq^8(c_1c_2 \cdot c_3) &= Sq^6(c_1c_2)Sq^2(c_3) + Sq^2(c_1c_2)Sq^6(c_3) + Sq^4(c_1c_2)Sq^4(c_3) \\ &= c_1^3c_2^2c_3 + c_1c_3^3 + (c_1^3c_2 + c_1^2c_3 + c_1c_2^2)(c_1c_4 + c_2c_3). \end{aligned}$$

□

Next we consider  $Sq^i$  on  $\text{Im } \bar{\Delta}^*$ . The following proposition contains all we need.

**Proposition 9.2.**  $Sq^2\bar{\Delta}^*(y_8) = w_2'^2w_3'^2$ ,  $Sq^2\bar{\Delta}^*(y_9) = w_2'w_3'^3$ ,  $Sq^4\bar{\Delta}^*(y_8) = w_3'^4 + w_2'^3w_3'^2$ ,  $Sq^4\bar{\Delta}^*(y_9) = w_2'^2w_3'^3$ ,  $Sq^4\bar{\Delta}^*(y_{12}) = w_2'^2w_3'^4$ ,

*Proof.* By Proposition 8.6,  $\bar{\Delta}^*(y_8) = w_2'^4 + w_2'w_3'^2$ ,  $\bar{\Delta}^*(y_9) = w_2'^3w_3'$  and  $\bar{\Delta}^*(y_{12}) = w_2'^3w_3'^2$ . Using the Wu formula and the Cartan formula, it is straightforward to verify the formulas in the proposition.  $Sq^2\bar{\Delta}^*(y_8) = w_2'^2w_3'^2$  is obvious.

$$\begin{aligned} Sq^2\bar{\Delta}^*(y_9) &= Sq^2(w_2' \cdot w_2'^2)w_3' + w_2'^3Sq^2(w_3') \\ &= (w_2'^4 + w_2'w_3'^2)w_3' + w_2'^4w_3' = w_2'w_3'^3, \end{aligned}$$

$$Sq^4\bar{\Delta}^*(y_8) = [Sq^1(w_2')]^4 + w_2'[Sq^4(w_3')] = w_3'^4 + w_2'^3w_3'^2,$$

$$\begin{aligned} Sq^4\bar{\Delta}^*(y_9) &= Sq^4(w_2' \cdot w_2'^2)w_3' + Sq^2(w_2' \cdot w_2'^2)Sq^2(w_3') + Sq^1(w_2'^3)w_3'^2 \\ &= (w_2'^5 + w_2'^2w_3'^2)w_3' + (w_2'^4 + w_2'w_3'^2)w_2'w_3' + w_2'^2w_3'^3 = w_2'^2w_3'^3, \end{aligned}$$

$$\begin{aligned} Sq^4\bar{\Delta}^*(y_{12}) &= Sq^4(w_2' \cdot w_2'^2)w_3'^2 + w_2'^3Sq^4(w_3'^2) \\ &= (w_2'^5 + w_2'^2w_3'^2)w_3'^2 + w_2'^5w_3'^2 = w_2'^2w_3'^4. \end{aligned}$$

□

For  $Sq^i$  on  $\text{Im } \phi^*$ , we only need a partial result.

**Proposition 9.3.**  $Sq^2\phi^*(y_{12}) = w_2^4w_3^2 + w_2w_3^4 + w_3^2w_4^2$ , and

$$Sq^8\phi^*(y_{12}) \equiv \phi^*(y_8)\phi^*(y_{12}) + w_3^4w_4^2 \pmod{(w_2)}.$$

*Proof.* Recall from Proposition 8.3 that  $\phi^*(y_{12}) = w_6^2 + w_2^2w_4^2 + w_2w_5^2 + w_2^3w_3^2$ . One easily computes that  $Sq^2(w_6^2) = 0$ ,  $Sq^2(w_2^2w_4^2) = w_2^2w_5^2 + w_3^2w_4^2$ ,  $Sq^2(w_2w_5^2) = w_2^2w_5^2$ ,  $Sq^2(w_2^3w_3^2) = w_2^4w_3^2 + w_2w_3^4$ , hence the formula for  $Sq^2\phi^*(y_{12})$  holds.

Now we compute the action of  $Sq^8$  on each terms in the expression of  $\phi^*(y_{12})$ , modulo the ideal  $(w_2)$ .  $Sq^8(w_2w_5^2) \equiv 0 \pmod{(w_2)}$  is obvious.

$$Sq^8(w_6^2) = [Sq^4(w_6)]^2 = w_4^2w_6^2 \equiv \phi^*(y_8)\phi^*(y_{12}) \pmod{(w_2)}.$$

$$\begin{aligned} Sq^8(w_2^3w_3^2) &= Sq^2(w_2^3)Sq^6(w_3^2) + Sq^4(w_2^3)Sq^4(w_3^2) + Sq^6(w_2^3)Sq^2(w_3^2) \\ &\equiv 0 \pmod{(w_2)}, \end{aligned}$$

since  $Sq^2(w_2^3) = w_2^4 + w_2w_3^2$ ,  $Sq^4(w_2^3) = w_2^5 + w_2^2w_3^2$  and  $Sq^6(w_2^3) = w_2^6$ .

$$\begin{aligned} Sq^8(w_2^2w_4^2) &= [Sq^4(w_2w_4)]^2 = [w_2w_4^2 + w_3Sq^3(w_4) + w_2^2Sq^2(w_4)]^2 \\ &\equiv [w_3Sq^3(w_4)]^2 \equiv w_3^4w_4^2 \pmod{(w_2)}. \end{aligned}$$

Then the formula for  $Sq^8\phi^*(y_{12})$  follows.  $\square$

Now we are ready to prove Theorem 2.5. The action of  $Sq^1$  on all generators  $y_i$ , and  $Sq^2(y_3)$ ,  $Sq^4(y_5)$  were given in Section 8.  $Sq^2(y_5) = 0$  since  $H^7(BPU(4); \mathbb{Z}/2) = 0$ .

Since  $H^{10}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^5, y_2y_8, y_5^2\}$ , Proposition 9.1, 9.2 and 8.6 give  $Sq^2(y_8) = y_5^2$ . Similarly, we have  $Sq^4(y_8) = y_2^2y_8 + y_{12} + y_3^4$ , using the fact that  $H^{12}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^6, y_2^2y_8, y_{12}, y_3^4\}$ .

Since  $H^{11}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_3^2y_5, y_3y_8\}$  and  $\bar{\Delta}^*(y_3^2y_5) = w_2'w_3'^3$ ,  $\bar{\Delta}^*(y_3y_8) = w_3'(w_2'^4 + w_2'w_3'^2)$  by Proposition 8.6, it follows from Proposition 9.2 that  $Sq^2(y_9) = y_3^2y_5$ . Similarly, the formula  $Sq^4(y_9) = y_3y_5^2$  follows from Proposition 9.2, and the fact that  $H^{13}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_3y_5^2, y_5y_8\}$ , using  $\bar{\Delta}^*(y_3y_5^2) = w_2'^2w_3'^3$  and  $\bar{\Delta}^*(y_5y_8) = w_2'w_3'(w_2'^4 + w_2'w_3'^2)$ .

Since  $H^{17}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_3^4y_5, y_3^3y_8, y_3y_5y_9, y_5y_{12}, y_8y_9\}$ , we may assume that  $Sq^8(y_9) = ay_3^4y_5 + by_3^3y_8 + cy_3y_5y_9 + dy_5y_{12} + ey_8y_9$ . Then

$$Sq^1Sq^8(y_9) = ay_3^6 + by_3^6 + c(y_3^3y_9 + y_3y_5^3) + d(y_3^2y_{12} + y_3y_5^3) + e(y_3^3y_9 + y_5^2y_8).$$

Hence, the relation  $Sq^1Sq^8(y_9) = Sq^9(y_9) = y_9^2 = y_3^2y_{12} + y_5^2y_8$  gives  $a = b$  and  $c = d = e = 1$ . On the other hand, if  $a \neq 0$ , then  $\bar{\Delta}^*(y_3^4y_5) = w_2'w_3'^5$  would appear in the expression of  $Sq^8\bar{\Delta}^*(y_9) = Sq^8(w_2'^3w_3')$ , but this is not true by an easy computation, so  $Sq^8(y_9) = y_3y_5y_9 + y_5y_{12} + y_8y_9$ .

Notice that  $H^{14}(BPU(4); \mathbb{Z}/2)$  is spanned by  $y_2^7, y_2^2y_8, y_2y_{12}, y_3^3y_5, y_3^2y_8, y_5y_9$ . Then in the expression  $Sq^2(y_{12}) = ay_2^7 + by_2^2y_8 + cy_2y_{12} + dy_3^3y_5 + ey_3^2y_8 + fy_5y_9$ ,  $a = b = 0$ ,  $c = 1$  by Proposition 9.1, and  $d = f = 0$ ,  $e = 1$  by Proposition 9.3

and the fact that  $\phi^*(y_3^3 y_5) \equiv w_3^3 w_5$ ,  $\phi^*(y_5^2 y_8) \equiv w_3^2 w_4^2$ ,  $\phi^*(y_5 y_9) \equiv w_5(w_3 w_6 + w_4 w_5) \pmod{(w_2)}$  from Proposition 8.3, which proves  $Sq^2(y_{12}) = y_2 y_{12} + y_3^2 y_8$ .

Since  $H^{16}(BPU(4); \mathbb{Z}/2) \cong \mathbb{Z}/2\{y_2^8, y_2^4 y_8, y_2^2 y_{12}, y_8^2, y_3^2 y_5^2, y_3 y_5 y_8\}$ , Proposition 9.1, 9.2 and 8.6 give  $Sq^4(y_{12}) = y_2^2 y_{12} + y_3^2 y_5^2$ .

An easy calculation shows that  $H^{20}(BPU(4); \mathbb{Z}/2)$  is spanned by  $y_3^5 y_5$ ,  $y_3^4 y_8$ ,  $y_3^2 y_5 y_9$ ,  $y_3 y_5 y_{12}$ ,  $y_3 y_8 y_9$ ,  $y_5^4$  and monomials in  $y_2$ ,  $y_8$ ,  $y_{12}$  of degree 20. Proposition 9.1 says that among monomials in  $y_2$ ,  $y_8$ ,  $y_{12}$  of degree 20, only  $y_8 y_{12}$  appears in the expression of  $Sq^8(y_{12})$ . Furthermore, since  $\phi^*(y_3^3 y_5) \equiv w_3^3 w_5$ ,  $\phi^*(y_3^4 y_8) \equiv w_3^4 w_4^2$ ,  $\phi^*(y_3^2 y_5 y_9) \equiv w_3 w_5(w_3 w_6 + w_4 w_5)$ ,  $\phi^*(y_3 y_5 y_{12}) \equiv w_3 w_5 w_6^2$ ,  $\phi^*(y_3 y_8 y_9) \equiv w_3 w_4^2(w_3 w_6 + w_4 w_5)$ ,  $\phi^*(y_5^4) \equiv w_5^4 \pmod{(w_2)}$  by Proposition 8.3, we obtain from Proposition 9.3 that  $Sq^8(y_{12}) = y_3^4 y_8 + y_8 y_{12}$ .

The proof of Theorem 2.5 is completed.

## 10. PROOF OF THEOREM 2.3

Let  $\bar{\alpha}_i \in H^{2i}(BPU(4); \mathbb{Z})$ ,  $i = 2, 3, 4, 6$ , be the elements constructed in Section 7, and for simplicity let  $x_1$ ,  $y_{2,I}$  denote  $\chi^*(x_1)$ ,  $\chi^*(y_{2,I}) \in H^*(BPU(4); \mathbb{Z})$  for  $\chi : BPU(4) \rightarrow K(\mathbb{Z}, 3)$ .

**Proposition 10.1.** *The cohomology ring  $H^*(BPU(4); \mathbb{Z})$  is generated by  $\bar{\alpha}_i$ ,  $i = 2, 3, 4, 6$ , and  $x_1$ ,  $y_{2,1}$ ,  $y_{2,(1,0)}$ .*

Before proving Proposition 10.1, we need a result on the image of  $H^*(BPU(4); \mathbb{Z})$  under the mod 2 reduction.

**Lemma 10.2.** *The image of  $\rho : H^*(BPU(4); \mathbb{Z}) \rightarrow H^*(BPU(4); \mathbb{Z}/2)$  is the subalgebra generated by  $y_2^2$ ,  $y_2^3$ ,  $y_3$ ,  $y_5^2$ ,  $y_8 + y_3 y_5$ ,  $y_{12} + y_3 y_9$ ,  $y_3^2 y_9 + y_5^3$ .*

*Proof.* Let  $S \subset H^*(BPU(4); \mathbb{Z}/2)$  be the subring generated by the elements in the Lemma. First we prove that  $S \subset \text{Im } \rho$ . By Proposition 8.7,  $\rho(\bar{\alpha}_2) = y_2^2$ ,  $\rho(\bar{\alpha}_3) = y_2^3$ ,  $\rho(\bar{\alpha}_4) = y_8 + y_3 y_5$ ,  $\rho(\bar{\alpha}_6) = y_{12} + y_3 y_9$ . Recall from Section 4 that in  $H^*(K(\mathbb{Z}, 3); \mathbb{Z}/2)$  we have  $y_{2,1} = x_{2,0}^2$  and  $y_{2,(1,0)} = y_{2,0} x_{2,1} + x_{2,0} y_{2,1} = x_1^2 x_{2,1} + x_{2,0}^3$ . Hence from Proposition 8.1 and the second part of Proposition 8.4 we have  $\rho(x_1) = y_3$ ,  $\rho(y_{2,1}) = y_5^2$ ,  $\rho(y_{2,(1,0)}) = y_3^5 + y_3^2 y_9 + y_5^3$ . It follows immediately that  $S \subset \text{Im } \rho$ .

For the reverse inclusion  $\text{Im } \rho \subset S$ , we use the fact that  $\text{Im } \rho = \ker \delta \subset \ker Sq^1$ , where  $\delta$  is the connecting homomorphism in the long exact sequence

$$\rightarrow H^i(-; \mathbb{Z}) \xrightarrow{\cdot 2} H^i(-; \mathbb{Z}) \xrightarrow{\rho} H^i(-; \mathbb{Z}/2) \xrightarrow{\delta} H^{i+1}(-; \mathbb{Z}) \rightarrow$$

induce by the sequence  $0 \rightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \rightarrow \mathbb{Z}/2 \rightarrow 0$ , noting that  $Sq^1 = \rho\delta$ . We claim that the kernel of  $Sq^1$  on  $H^*(BPU(4); \mathbb{Z}/2)$  is the subalgebra generated by  $y_2$ ,  $y_3$ ,  $y_5^2$ ,  $y_8 + y_3 y_5$ ,  $y_{12} + y_3 y_9$ ,  $y_3^2 y_9 + y_5^3$ , whose proof is deferred to the last Section for

reader's convenience. Then from the ring structure of  $H^*(BPU(4); \mathbb{Z}/2)$  it is easy to see that there is a  $\mathbb{Z}/2$ -module isomorphism

$$\ker Sq^1 \cong S \oplus \mathbb{Z}\{y_2, y_2y_8, y_2y_{12}\}.$$

Furthermore, we know that  $H^2(BPU(n); \mathbb{Z}) = 0$  and  $H^3(BPU(n); \mathbb{Z}) = \mathbb{Z}/n\{x_1\}$  for any  $n$ . Hence the universal coefficient theorem shows that  $H^2(BPU(4); \mathbb{Z}/4) \cong \mathbb{Z}/4$ . The sequence  $0 \rightarrow \mathbb{Z} \xrightarrow{\cdot 4} \mathbb{Z} \rightarrow \mathbb{Z}/4 \rightarrow 0$  also induces a long exact sequence of cohomology, and there is a commutative diagram

$$\begin{array}{ccccccccc} \longrightarrow & H^i(-; \mathbb{Z}) & \xrightarrow{\cdot 4} & H^i(-; \mathbb{Z}) & \xrightarrow{\rho'} & H^i(-; \mathbb{Z}/4) & \xrightarrow{\delta'} & H^{i+1}(-; \mathbb{Z}) & \longrightarrow \\ & \downarrow \cdot 2 & & \parallel & & \downarrow \theta = \text{mod } 2 & & \downarrow \cdot 2 & \\ \longrightarrow & H^i(-; \mathbb{Z}) & \xrightarrow{\cdot 2} & H^i(-; \mathbb{Z}) & \xrightarrow{\rho} & H^i(-; \mathbb{Z}/2) & \xrightarrow{\delta} & H^{i+1}(-; \mathbb{Z}) & \longrightarrow \end{array}$$

Suppose that  $y'_2$  generates  $H^2(BPU(4); \mathbb{Z}/4)$ . Then  $\delta'(y'_2) = x_1$ , and  $\theta(y'_2) = y_2$ . Hence, the above diagram shows that  $\delta(y_2) = 2x_1 \neq 0$ , and then

$$\begin{aligned} \delta(y_2y_8) &= \delta(y_2(y_8 + y_3y_5)) = \delta(y_2\rho(\bar{\alpha}_4)) = 2x_1\bar{\alpha}_4, \\ \delta(y_2y_{12}) &= \delta(y_2(y_{12} + y_3y_9)) = \delta(y_2\rho(\bar{\alpha}_6)) = 2x_1\bar{\alpha}_6. \end{aligned}$$

Since Proposition 5.3 implies that  $x_1\bar{\alpha}_4$  and  $x_1\bar{\alpha}_6$  have order 4 in  $H^*(BPU(4); \mathbb{Z})$ , we obtain  $\delta(y_2y_8), \delta(y_2y_{12}) \neq 0$ . It follows that  $y_2, y_2y_8, y_2y_{12} \notin \text{Im } \rho$ , and then  $\text{Im } \rho \subset S$ .  $\square$

*Proof of Proposition 10.1.* Let  $R$  be the subring generated by the elements in the proposition, and let  $N = H^*(BPU(4); \mathbb{Z})/R$  be the quotient  $\mathbb{Z}$ -module. Then lemma 10.2 and the first paragraph of its proof show that  $\rho(R) = \rho(H^*(BPU(4); \mathbb{Z}))$ , i.e.  $\rho(N) = 0$ . We need to show that  $N$  itself is zero.

Let  $\mathfrak{t}$  be the ideal consisting of torsion elements of  $H^*(BPU(4); \mathbb{Z})$ , and let  $F = H^*(BPU(4); \mathbb{Z})/\mathfrak{t}$ . Since  $\pi^*(\bar{\alpha}_i) = \alpha_i$ , using Proposition 7.1 and the fact that torsion free elements in  ${}^U E$  concentrated in  ${}^U E_*^{0,*}$ , we see that

$$\pi^*(R) \cong K_4 \cong {}^U E_\infty^{0,*} \cong F.$$

Hence we get a commutative diagram of short exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathfrak{t} \cap R & \longrightarrow & R & \longrightarrow & F & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & \mathfrak{t} & \longrightarrow & H^*(BPU(4); \mathbb{Z}) & \longrightarrow & F & \longrightarrow & 0 \end{array}$$

This implies that  $N \cong \mathfrak{t}/(\mathfrak{t} \cap R)$  as  $\mathbb{Z}$ -modules. Recall that the torsion elements of  $H^*(BPU(4); \mathbb{Z})$  are all 2-primary. So if  $N \neq 0$ , then  $N \otimes \mathbb{Z}/2 \neq 0$ . However, since  $0 = \rho(N) \cong N \otimes \mathbb{Z}/2$ , we must have  $N = 0$ . Hence  $R = H^*(BPU(4); \mathbb{Z})$ .  $\square$

Let  $A = K_4 \otimes \mathbb{Z}[x_3, x_{10}, x_{12}]/I$  be the ring defined in Theorem 2.3.

**Proposition 10.3.** *There is a surjective homomorphism of graded rings:*

$$A \rightarrow H^*(BPU(4); \mathbb{Z}) \quad x_3 \mapsto x_1, \quad x_{10} \mapsto y_{2,1}, \quad x_{15} \mapsto y_{2,(1,0)}, \quad \alpha_i \mapsto \bar{\alpha}_i.$$

*Proof.* By Proposition 10.1, we only need to show that this map is well-defined, in other words, the element  $\alpha = 64\alpha_6 - \alpha_2^3 - 27\alpha_3^2 + 48\alpha_2\alpha_4$  and all generators of the ideal  $I$  map to zero.

We already know that in  $H^*(BPU(4); \mathbb{Z})$ ,  $x_1$  has order 4 by [11, Theorem 1.1], and  $x_1^2 = y_{2,0}$ ,  $y_{2,1}$ ,  $y_{2,(1,0)}$  are all of order 2 by Theorem 4.1. From this one can see that the order of a torsion element of even degree in  $H^*(BPU(4); \mathbb{Z})$  is at most 2.

Let  $\bar{\alpha} = 64\bar{\alpha}_6 - \bar{\alpha}_2^3 - 27\bar{\alpha}_3^2 + 48\bar{\alpha}_2\bar{\alpha}_4$ . Then  $\pi^*(\bar{\alpha}) = \alpha = 0$  for  $\pi : BU(4) \rightarrow BPU(4)$ , since  $\text{Im } \pi^* \cong K_4$  by Proposition 7.1. Hence  $\bar{\alpha}$  is a torsion element. From Proposition 8.7 we know that  $\rho(\bar{\alpha}) = 0$ , and so  $\bar{\alpha} \in 2H^*(BPU(4); \mathbb{Z})$ . Then we get  $\bar{\alpha} = 0$  since  $\bar{\alpha}$  is of even degree.

Since  $y_2y_5 = 0$  and  $\rho(y_{2,1}) = \chi^*(x_{2,0}^2) = x_2^2$ , we have  $\rho(\bar{\alpha}_2y_{2,1}) = \rho(\bar{\alpha}_3y_{2,1}) = 0$ . Hence  $\bar{\alpha}_2y_{2,1}, \bar{\alpha}_3y_{2,1} \in 2H^*(BPU(4); \mathbb{Z})$ , and then  $\bar{\alpha}_2y_{2,1} = \bar{\alpha}_3y_{2,1} = 0$ , since  $\bar{\alpha}_2y_{2,1}, \bar{\alpha}_3y_{2,1}$  are of even degree.

It is easy to see that  $\rho(\bar{\alpha}_2x_1) = \rho(\bar{\alpha}_3x_1) = 0$ , since  $y_2y_3 = 0$  and  $\rho(\bar{\alpha}_2) = y_2^2$ ,  $\rho(\bar{\alpha}_3) = y_2^3$ ,  $\rho(x_1) = y_3$ . Hence  $\bar{\alpha}_2x_1, \bar{\alpha}_3x_1 \in 2H^*(BPU(4); \mathbb{Z})$ . Then the fact that  $\bar{\alpha}_2x_1$  generates  $H^7(BPU(4); \mathbb{Z})$  gives the relation  $\bar{\alpha}_2x_1 = 0$ . Similarly, since  $\bar{\alpha}_2x_1, x_1^3$  generate  $H^9(BPU(4); \mathbb{Z})$  and  $x_1^3$  has order 2, we get  $\bar{\alpha}_3x_1 = 0$ .

Since  $\rho(y_{2,(1,0)}) = \chi^*(x_1^2x_{2,1} + x_{2,0}^3) = y_3^5 + y_3^2y_9 + y_3^3$ , we have  $\rho(\bar{\alpha}_2y_{2,(1,0)}) = 0$ , then  $\bar{\alpha}_2y_{2,(1,0)} = 2\lambda$  for some element  $\lambda$  of degree 19. Checking each torsion element of degree 19, one can show that  $2\lambda = 0$  or  $2\bar{\alpha}_4^2x_1$ . But the later is impossible, since if this is the case, it would imply that  $\bar{\alpha}_4^2x_1$  in  ${}^UE_\infty^{3,16}$  has order 2 because of the Serre filtration construction, which contradicts the second part of Proposition 7.1. Hence we get the relation  $\bar{\alpha}_2y_{2,(1,0)} = 0$ . The proof of  $\bar{\alpha}_3y_{2,(1,0)} = 0$  is easier, using the fact that all torsion elements of degree 21 have order 2.

Let  $y = x_1^6\bar{\alpha}_6 + x_1^4y_{2,1}\bar{\alpha}_4 + x_1^5y_{2,(1,0)} + y_{2,1}^3 + y_{2,(1,0)}^2$ . Then using the computations in the first paragraph of the proof Lemma 10.2 and the relation  $y_9^2 = y_3^2y_{12} + y_5^2y_8$  in  $H^*(BPU(4); \mathbb{Z}/2)$ , one readily checks that  $\rho(y) = 0$ . Since  $y$  has even degree,  $y = 0$  by the same reasoning.

The above calculations verify that  $\alpha$  and all generators of  $I$  map to zero, finishing the proof.  $\square$

The following Proposition completes the proof of Theorem 2.3.

**Proposition 10.4.** *The homomorphism in Proposition 10.3 is injective.*

*Proof.* The proposition is equivalent to saying that if a polynomial  $f$  in the variables  $\alpha_i$ ,  $i = 2, 3, 4, 6$ ,  $x_3$ ,  $x_{10}$ ,  $x_{15}$ , maps to zero in  $H^*(BPU(4); \mathbb{Z})$ , then  $f \in (\alpha) + I$ , where  $\alpha = 64\alpha_6 - \alpha_2^3 - 27\alpha_3^2 + 48\alpha_2\alpha_4$ .

Write  $f = f(\bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4, \bar{\alpha}_6, x_1, y_{2,1}, y_{2,(1,0)})$ . Since  $f$  is zero in  $H^*(BPU(4); \mathbb{Z})$ ,  $\pi^*(f) = 0$ , and then  $f$  can be written as  $f = f_0 + f_1$ , where  $f_0$  belongs to the ideal of  $\mathbb{Z}[\bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4, \bar{\alpha}_6]$  generated by  $\bar{\alpha} = 64\bar{\alpha}_6 - \bar{\alpha}_2^3 - 27\bar{\alpha}_3^2 + 48\bar{\alpha}_2\bar{\alpha}_4$ , and  $f_1 \in (x_1, y_{2,1}, y_{2,(1,0)})$ . Hence it suffices to show that  $f_1 \in I$ . Here we identify  $I$  with the corresponding ideal of  $\mathbb{Z}[\bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_4, \bar{\alpha}_6, x_1, y_{2,1}, y_{2,(1,0)}]$ .

Let  $\phi : BSO(6) \rightarrow BPSO(6) = BPU(4)$  be the map in Section 8, and let

$$\begin{aligned} g_1 &= \phi^* \rho(x_1) = \phi^*(y_3), & g_2 &= \phi^* \rho(y_{2,1}) = \phi^*(y_5^2), \\ g_3 &= \phi^* \rho(\bar{\alpha}_4) = \phi^*(y_8 + y_3 y_5), & g_4 &= \phi^* \rho(\bar{\alpha}_6) = \phi^*(y_{12} + y_3 y_9), \\ g_5 &= \phi^* \rho(y_{2,(1,0)}) = \phi^*(y_3^5 + y_3^2 y_9 + y_5^3). \end{aligned}$$

Then  $\phi^* \rho(f_1)$  belongs to the ideal  $I_1$  of relations between the elements  $g_1, \dots, g_5$  in  $H^*(BSO(6); \mathbb{Z}/2) \cong \mathbb{Z}/2[w_2, \dots, w_6]$ .

Let us compute the ideal  $I_1 \subset \mathbb{Z}/2[g_1, \dots, g_5]$ . By Proposition 8.3, modulo  $(w_2)$ ,

$$\begin{aligned} g_1 &\equiv w_3, & g_2 &\equiv w_5^2, & g_3 &\equiv w_4^2 + w_3 w_5, \\ g_4 &\equiv w_6^2 + w_3^2 w_6 + w_3 w_4 w_5, & g_5 &\equiv w_3^5 + w_3^3 w_6 + w_3^2 w_4 w_5 + w_5^3. \end{aligned}$$

It is easy to see that the four elements  $g_1, \dots, g_4$  are algebraically independent. Hence the Krull dimension of the subalgebra of  $\mathbb{Z}/2[w_2, \dots, w_6]$  generated by  $g_i$ ,  $i = 1, \dots, 5$ , is at least 4, which means that  $I_1$  is a principal ideal of  $\mathbb{Z}/2[g_1, \dots, g_5]$ . To see this, note that  $I_1$  is the kernel of the map  $\mathbb{Z}/2[g_1, \dots, g_5] \rightarrow \mathbb{Z}/2[w_2, \dots, w_6]$  with target a UFD, so  $I_1$  is a prime ideal of  $\mathbb{Z}/2[g_1, \dots, g_5]$ . Since the Krull dimension of  $\mathbb{Z}/2[g_1, \dots, g_5]/I_1$  is at least 4, the height of  $I_1$  is at most 1, and then  $I_1$  is principal since  $\mathbb{Z}/2[g_1, \dots, g_5]$  is a UFD.

Suppose that  $h = h(g_1, \dots, g_5)$  generates  $I_1$ . Since the powers  $w_6^i$  only appear in the expressions of  $g_4$  and  $g_5$ , it follows that if  $h \neq 0$ , then the degrees of  $g_4$  and  $g_5$  in  $h$  are both nonzero. Furthermore, since the degree of  $w_6$  in  $g_4$  (resp. in  $g_5$ ) is 2 (resp. 1), the degree of  $g_5$  in  $h$  is at least 2 when  $h \neq 0$ . We have seen in the proof of Proposition 10.3 that  $\phi^* \rho(y) = g_1^6 g_4 + g_1^4 g_2 g_3 + g_1^5 g_5 + g_2^3 + g_5^2 = 0$  in  $\mathbb{Z}/2[w_2, \dots, w_6]$ , where  $y = x_1^6 \bar{\alpha}_6 + x_1^4 y_{2,1} \bar{\alpha}_4 + x_1^5 y_{2,(1,0)} + y_{2,1}^3 + y_{2,(1,0)}^2$ , hence we must have  $h = g_1^6 g_4 + g_1^4 g_2 g_3 + g_1^5 g_5 + g_2^3 + g_5^2$ .

Since  $f_1 \in (x_1, y_{2,1}, y_{2,(1,0)})$  and  $\phi^* \rho(\bar{\alpha}_2) = \phi^* \rho(\bar{\alpha}_3) = 0$ , the above analysis implies that

$$f_1 \in (y) + (\bar{\alpha}_2, \bar{\alpha}_3) \cdot (x_1, y_{2,1}, y_{2,(1,0)}) + 2(x_1, y_{2,1}, y_{2,(1,0)}).$$

We know that  $f$  and  $f_0$  are zero in  $H^*(BPU(4); \mathbb{Z})$ , so  $f_1 = 0$  in  $H^*(BPU(4); \mathbb{Z})$ . Since  $y$ ,  $2y_{2,1}$ ,  $2y_{2,(1,0)}$  and  $(\bar{\alpha}_2, \bar{\alpha}_3) \cdot (x_1, y_{2,1}, y_{2,(1,0)})$  are all zero in  $H^*(BPU(4); \mathbb{Z})$

and the order of  $x_1\xi$  is 4 for any  $\xi \in \mathbb{Z}[\bar{\alpha}_4, \bar{\alpha}_6]$  by the second part of Proposition 7.1, it follows that  $f_1$  is actually in the ideal

$$(y) + (\bar{\alpha}_2, \bar{\alpha}_3) \cdot (x_1, y_{2,1}, y_{2,(1,0)}) + 4(x_1),$$

which is contained in  $I$ . The proof is completed.  $\square$

### 11. KERNEL OF $Sq^1$ ON $H^*(BPU(4); \mathbb{Z}/2)$

In this section we prove the claim in the proof of Lemma 10.2 that the kernel of the action  $Sq^1$  on  $H^*(BPU(4); \mathbb{Z}/2)$  is the subalgebra generated by  $y_2, y_3, y_5^2, y_8 + y_3y_5, y_{12} + y_3y_9, y_3^2y_9 + y_3^3$ . To make the computation easier, we shall establish a  $\mathbb{Z}/2$ -algebra isomorphism to  $H^*(BPU(4); \mathbb{Z}/2)$ , and define an operation on it corresponding to  $Sq^1$ .

Let  $W$  be the quotient of the polynomial algebra  $\mathbb{Z}/2[x_2, x_3, x_5, x_8, x_9, x_{12}]$  by the ideal generated by  $x_2x_3, x_2x_5, x_2x_9$  and  $x_9^2 + x_3^2x_{12} + x_5^2x_8 + x_3^3x_9 + x_3x_5^3$ . Then it is easy to see that there is an algebra isomorphism  $W \cong H^*(BPU(4); \mathbb{Z}/2)$  given by  $x_i \mapsto y_i$  for  $i = 2, 3, 5, 9, x_8 \mapsto y_8 + y_3y_5, x_{12} \mapsto y_{12} + y_3y_9$ . Define an operation  $\mathcal{D}$  on  $W$  by  $\mathcal{D}(x_i) = 0$  for  $i = 2, 3, 8, 12$ ,  $\mathcal{D}(x_5) = x_3^2$ ,  $\mathcal{D}(x_9) = x_5^2$ , and the Leibniz rule  $\mathcal{D}(ab) = \mathcal{D}(a)b + a\mathcal{D}(b)$ . Then using Theorem 2.5, one easily verifies that there is a commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{\mathcal{D}} & W \\ \cong \downarrow & & \cong \downarrow \\ H^*(BPU(4); \mathbb{Z}/2) & \xrightarrow{Sq^1} & H^*(BPU(4); \mathbb{Z}/2) \end{array}$$

Thus,  $W$  can be viewed as a DGA with differential  $\mathcal{D}$ . We compute its cohomology as follows

**Proposition 11.1.**  $H^*(W, \mathcal{D}) \cong \mathbb{Z}/2[x_2, x_8, x_{12}] \otimes E_{\mathbb{Z}/2}[x_3]$ .

Our desired result on  $\ker Sq^1$  is a direct consequence of the following corollary of Proposition 11.1.

**Corollary 11.2.**  $\ker \mathcal{D}$  is the subalgebra of  $W$  generated by  $x_2, x_8, x_{12}, x_3, x_5^2, x_3^2x_9 + x_5^3$ .

*Proof.* Since  $H^*(W, \mathcal{D}) = \ker \mathcal{D} / \text{Im } \mathcal{D}$ , it follows from Proposition 11.1 that as  $\mathbb{Z}/2$ -modules

$$\ker \mathcal{D} \cong \text{Im } \mathcal{D} \oplus \mathbb{Z}/2[x_2, x_8, x_{12}] \oplus \mathbb{Z}/2[x_2, x_8, x_{12}] \cdot x_3.$$

Now we compute  $\text{Im } \mathcal{D}$ . For a monomial  $\mathbf{m} = \mathbf{n} \cdot x_3^i x_5^j x_9^k \in W$ ,  $\mathbf{n} \in \mathbb{Z}/2[x_2, x_8, x_{12}]$ , by the rule of  $\mathcal{D}$  we have

$$\mathcal{D}(\mathbf{m}) = \mathbf{n} \cdot (\tilde{j} \cdot x_3^{i+2} x_5^{j-1} x_9^k + \tilde{k} \cdot x_3^i x_5^{j+2} x_9^{k-1}).$$

Here  $\tilde{*}$  means the mod 2 number. Hence  $\text{Im } \mathcal{D}$  is a  $\mathbb{Z}/2$ -vector space spanned by elements of the forms:

$$\mathbf{n} \cdot x_3^r x_5^{2s} x_9^{2t}, \quad \mathbf{n} \cdot x_3^i x_5^{2q} x_9^{2t}, \quad \mathbf{n} \cdot x_3^i x_5^{2s} x_9^{2t} (x_3^2 x_9 + x_5^3),$$

where  $i, s, t \geq 0$ ,  $q \geq 1$ ,  $r \geq 2$ . Note that  $x_9^{2t} = [x_3^2 x_{12} + x_5^2 x_8 + x_3(x_3^2 x_9 + x_5^3)]^t$  by definition. Then an easy calculation gives the result.  $\square$

*Proof of Proposition 11.1.* Define a  $\mathbb{Z}/2$ -linear map  $\lambda : W \rightarrow W$  by  $\lambda(\mathbf{m}) = \mathbf{m}$  if  $\mathbf{m} = x_2^a x_8^b x_{12}^c$  or  $x_2^a x_8^b x_{12}^c x_3$ ,  $a, b, c \geq 0$ , and  $\lambda(\mathbf{m}) = 0$  for all other monomials. It is easy to show that  $\lambda$  is a chain map associated to the differential  $\mathcal{D}$ . We claim that  $\lambda$  is chain homotopic to the identity map. Since  $\mathcal{D}$  restricted to  $\mathbb{Z}/2[x_2, x_8, x_{12}, x_3]$  is zero and any element in  $\mathbb{Z}/2[x_2, x_8, x_{12}] \cdot \mathbb{Z}/2\{1, x_3\}$  is clearly not in  $\text{Im } \mathcal{D}$ , the proposition follows immediately from this claim.

Now we construct the chain homotopy  $P$  as follows. For a monomial  $\mathbf{m} = \mathbf{n} \cdot x_3^i x_5^j x_9^k \in W$ ,  $\mathbf{n} \in \mathbb{Z}/2[x_2, x_8, x_{12}]$ , let

$$P(\mathbf{m}) = \begin{cases} \mathbf{n} \cdot x_3^{i-2} x_5^{j+1} x_9^k & \text{if } i \geq 2, \\ \mathbf{n} \cdot x_3^i x_5^{j-2} x_9^{k+1} & \text{if } i \leq 1, j, k \text{ even, } j \neq 0, \\ \mathbf{n} \cdot x_3^i x_9^{k-2} (x_5 x_{12} + x_8 x_9 + x_3 x_5 x_9) & \text{if } i \leq 1, j = 0, k \geq 2 \text{ even,} \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to show that  $PD + DP = \lambda + \text{id}$  on all monomials in  $W$  case by case.  $\square$

## REFERENCES

- [1] B. Antieau and B. Williams, *The period-index problem for twisted topological K-theory*, *Geom. Topol.* **18** (2014), no. 2, 1115–1148.
- [2] ———, *The topological period-index problem over 6-complexes*, *J. Topol.* **7** (2014), no. 3, 617–640.
- [3] M. Artin, *Brauer-Severi varieties*, In: *Brauer groups in ring theory and algebraic geometry*, *Lect. Notes in Math.*, vol. 917, Springer, Berlin, 1982, pp. 194–210.
- [4] H. Cartan, *Algèbres d'Eilenberg-Mac Lane et homotopie*, *Séminaire Henri Cartan*, vol. 7, no. 1, 1954/1955, available at [www.numdam.org](http://www.numdam.org).
- [5] D. Crowley and M. Grant, *The topological period-index conjecture for  $Spin^c$  6-manifolds*, *Ann. K-Theory* **5** (2020), no. 3, 605–620.
- [6] A. J. de Jong, *The period-index problem for the Brauer group of an algebraic surface*, *Duke Math. J.* **123** (2004), no. 1, 71–94.
- [7] T. Ganea, *Induced fibrations and cofibrations*, *Trans. Amer. Math. Soc.* **127** (1967), no. 3, 442–459.
- [8] A. Grothendieck, *Le groupe de Brauer: I. Algèbres d'Azumaya et interprétations diverses*, *Séminaire Bourbaki*, vol. 9, pp. Exp. No. 290, 199–219, 1966.
- [9] X. Gu, *The topological period-index problem over 8-complexes I*, *J. Topol.* **12** (2019), no. 4, 1368–1395.
- [10] ———, *The topological period-index problem over 8-complexes II*, *Proc. Am. Math. Soc.* **148** (2020), no. 10, 4531–4545.

- [11] ———, *On the cohomology of the classifying spaces of projective unitary groups*, J. Topol. Anal. **13** (2021), no. 2, 535–573.
- [12] ———, *Some torsion classes in the Chow ring and cohomology of  $BPGL_n$* , J. London Math. Soc. **103** (2021), no. 1, 127–160.
- [13] X. Gu, Y. Zhang, Z. Zhang, and L. Zhong, *The  $p$ -primary subgroups of the cohomology of  $BPU_n$  in dimensions less than  $2p + 5$* , Proc. Am. Math. Soc. **150** (2022), no. 9, 4099–4111.
- [14] E. H. Brown Jr, *The cohomology of  $BSO_n$  and  $BO_n$  with integer coefficients*, Proc. Amer. Math. Soc. **85** (1982), no. 2, 283–288.
- [15] A. Kono and M. Mimura, *On the cohomology of the classifying spaces of  $PSU(4n + 2)$  and  $PO(4n + 2)$* , Publ. Res. Inst. Math. Sci. **10** (1975), no. 3, 691–720.
- [16] A. Kono, M. Mimura, and N. Shimada, *Cohomology of classifying spaces of certain associative  $H$ -spaces*, J. Math. Kyoto Univ. **15** (1975), no. 3, 607–617.
- [17] A. Kono and N. Yagita, *Brown-Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups*, Trans. Amer. Math. Soc. **339** (1993), no. 2, 781–798.
- [18] T. Kudo, *A transgression theorem*, Mem. Fac. Sci. Kyusyu Univ. **9** (1956), 79–81.
- [19] M. Lieblich, *Twisted sheaves and the period-index problem*, Compos. Math. **144** (2008), no. 1, 1–31.
- [20] K. Masaki and N. Yagita, *The Brown-Peterson cohomology of the classifying spaces of the projective unitary groups  $PU(p)$  and exceptional Lie groups*, Trans. Am. Math. Soc. **360** (2008), no. 5, 2265–2284.
- [21] H. Toda, *Cohomology of classifying spaces*, In: Homotopy theory and related topics (Kyoto, 1984), Adv. Stud. Pure Math., vol. 9, North-Holland, Amsterdam, 1987, pp. 75–108.
- [22] A. Vavpetič and A. Viruel, *On the mod  $p$  cohomology of  $BPU(p)$* , Trans. Am. Math. Soc. **357** (2005), no. 11, 4517–4532.
- [23] G. Vezzosi, *On the Chow ring of the classifying stack of  $PGL_{3,C}$* , J. Reine Angew. Math. **523** (2000), 1–54.
- [24] A. Vistoli, *On the cohomology and the Chow ring of the classifying space of  $PGL_p$* , J. Reine Angew. Math. **610** (2007), 181–227.
- [25] Y. Zhang, Z. Zhang, and L. Zhong, *The  $p$ -primary subgroups of the cohomology of  $BPU_n$  in dimension  $2p + 6$* , Topol. Appl. **3380** (2023), 108642, 12pp.

FEIFEI FAN, SCHOOL OF MATHEMATICAL SCIENCES, SOUTH CHINA NORMAL UNIVERSITY, GUANGZHOU, 510631, CHINA.

*Email address:* fanfeifei@mail.nankai.edu.cn