

Computable entanglement cost

Ludovico Lami,^{1,2,3,*} Francesco Anna Mele,^{4,†} and Bartosz Regula^{5,‡}

¹QuSoft, Science Park 123, 1098 XG Amsterdam, the Netherlands

²Korteweg-de Vries Institute for Mathematics, University of Amsterdam,
Science Park 105-107, 1098 XG Amsterdam, the Netherlands

³Institute for Theoretical Physics, University of Amsterdam,
Science Park 904, 1098 XH Amsterdam, the Netherlands

⁴NEST, Scuola Normale Superiore and Istituto Nanoscienze,
Consiglio Nazionale delle Ricerche, Piazza dei Cavalieri 7, IT-56126 Pisa, Italy

⁵Mathematical Quantum Information RIKEN Hakubi Research Team,
RIKEN Cluster for Pioneering Research (CPR) and RIKEN Center for Quantum Computing (RQC), Wako, Saitama 351-0198, Japan

Quantum information theory is plagued by the problem of regularisations, which require the evaluation of formidable asymptotic quantities. This makes it computationally intractable to gain a precise quantitative understanding of the ultimate efficiency of key operational tasks such as entanglement manipulation. Here we consider the problem of computing the asymptotic entanglement cost of preparing noisy quantum states under quantum operations with positive partial transpose (PPT). A previously claimed solution to this problem is shown to be incorrect. We construct instead an alternative solution in the form of two hierarchies of semi-definite programs that converge to the true asymptotic value of the entanglement cost from above and from below. Our main result establishes that this convergence happens exponentially fast, thus yielding an efficient algorithm that approximates the cost up to an additive error ε in time $\text{poly}(D, \log(1/\varepsilon))$, where D is the underlying Hilbert space dimension. To our knowledge, this is the first time that an asymptotic entanglement measure is shown to be efficiently computable despite no closed-form formula being available.

Introduction.— Quantum Shannon theory studies the fundamental limitations on the manipulation of quantum information in the presence of external noise. Calculating those limits often involves computing certain functions that encapsulate the ultimate capabilities of information carriers. Paradigmatic examples include the various capacities of quantum channels, such as the classical [1, 2], quantum [3–5], private [5, 6], and entanglement-assisted [7, 8] capacities, but also the operational entanglement measures that tell us how much entanglement can be extracted from a given bipartite quantum state, i.e. the distillable entanglement [9–13], and vice versa, how much entanglement must be invested to *create* that state [14, 15]. This latter quantity, the entanglement cost, is the main focus of this work.

With the sole exception of the entanglement-assisted capacity, all of the above functions are expressed by *regularised formulas*, i.e. formulas that involve an explicit limit $n \rightarrow \infty$ over the number of uses of the channel or available copies of the state. For example, by the Lloyd–Shor–Devetak theorem [3–5] the quantum capacity of a channel \mathcal{N} equals $Q(\mathcal{N}) = \lim_{n \rightarrow \infty} \frac{1}{n} I_c(\mathcal{N}^{\otimes n})$, where $I_c(\mathcal{N})$ is the ‘coherent information’ of \mathcal{N} , and $\mathcal{N}^{\otimes n}$ represents n parallel uses of \mathcal{N} . In stark contrast with classical information theory, for quantum channels it holds in general that $I_c(\mathcal{N}^{\otimes n}) \neq n I_c(\mathcal{N})$, meaning that evaluating the limit cannot be avoided. Such non-additivity is a fundamental feature of most settings encountered in quantum information [16–21]. Analogously, the entanglement cost of preparing a state $\rho = \rho_{AB}$ using local operations and

classical communication (LOCC) is given by

$$E_{c,\text{LOCC}}(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} E_f(\rho^{\otimes n}), \quad (1)$$

where E_f is the ‘entanglement of formation’ [11, 22]. The precise nature of these formulas is not so important here; what is important, however, is that the regularisation $n \rightarrow \infty$ makes them analytically hard to control and computationally intractable. Indeed, on the one hand the dimension of the quantum system on which $\mathcal{N}^{\otimes n}$ acts, or to which $\rho^{\otimes n}$ pertains, is exponential in n , quickly rendering numerical calculations infeasible as n grows; on the other, there is no guarantee on the quality of the approximation obtained by stopping at the n^{th} level of any of these hierarchies — for instance, an unbounded n may be required to even check that the quantum capacity is non-zero [23]. The regularisation thus appears to be an omnipresent curse that stifles almost every attempt to quantitatively understand the ultimate limitations of quantum information manipulation.

But is that really so? In this work we show how to overcome this fundamental obstacle, efficiently calculating a type of entanglement cost — expressed as a regularised quantity — on all quantum states. To this end, we look at a problem that has been studied by many authors [24–28], but for which a full solution had not been found prior to our work: namely, zero-error asymptotic entanglement dilution with ‘positive partial transpose’ (PPT) operations. The task is to determine the ultimate rate R at which singlets Φ_2 (i.e. two-qubit maximally entangled states) have to be consumed to prepare an asymptotic number of copies of a given bipartite quan-

tum state $\rho = \rho_{AB}$ using PPT operations only. That is, we declare a rate R achievable if for all sufficiently large n there exists a PPT operation Λ_n with the property that $\Lambda_n(\Phi_2^{\otimes \lfloor Rn \rfloor}) = \rho^{\otimes n}$. The infimum of all achievable rates R is the zero-error PPT entanglement cost of ρ , denoted $E_{c,\text{PPT}}^{\text{exact}}(\rho)$. In a nutshell, the key reason why many authors have been interested in this problem is that it provides a more tractable model for the fundamental problem of entanglement dilution under LOCC operations, whose underlying figure of merit, given by (1), is computationally inaccessible in most cases of interest.

Prior work. — In a pioneering paper by Audenaert, Plenio, and Eisert [24] it was shown that the PPT entanglement cost can be evaluated exactly for all bipartite states $\rho = \rho_{AB}$ that satisfy a condition known as ‘zero bi-negativity’ [24, 29, 30]. Specifically, if $|\rho^\Gamma|^\Gamma \geq 0$, where Γ denotes the partial transpose operation [31] defined as $(X_A \otimes Y_B)^\Gamma = X_A \otimes Y_B^\Gamma$ and $|X| := \sqrt{X^\dagger X}$ is the absolute value, then

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = \log_2 \|\rho^\Gamma\|_1 =: E_N(\rho). \quad (2)$$

The expression on the right, E_N , is a celebrated entanglement measure known as the *entanglement negativity* [32, 33], obtained by simply evaluating the trace norm $\|\cdot\|_1 := \text{Tr}|\cdot|$ of the partially transposed state. This framework thus provides a partial operational interpretation for E_N through its equality with the zero-error PPT cost $E_{c,\text{PPT}}^{\text{exact}}$ for some states. At the same time, Eq. (2) effectively solves the problem of computing $E_{c,\text{PPT}}^{\text{exact}}$ for states with zero bi-negativity, because E_N is efficiently computable via a *semi-definite program* (SDP) [34–36]:

$$E_N(\rho) = \log_2 \min \{ \text{Tr } S : -S \leq \rho^\Gamma \leq S \}. \quad (3)$$

But what to do for those states that have non-zero bi-negativity, i.e. satisfy $|\rho^\Gamma|^\Gamma \not\geq 0$? In their recent works [25, 26], Wang and Wilde claimed to have found a general solution to this problem, in the form of an alternative SDP-computable quantity called E_κ that would coincide with $E_{c,\text{PPT}}^{\text{exact}}$ for all states — and in particular reduce to E_N for states with zero bi-negativity. The validity of the claimed identity between E_κ and $E_{c,\text{PPT}}^{\text{exact}}$ rests crucially on the *additivity* of E_κ : in [25], it is averred that

$$E_\kappa(\rho^{\otimes n}) \stackrel{?}{=} n E_\kappa(\rho) \quad (4)$$

holds true for all bipartite states ρ and all positive integers n , thus completely eliminating the issue of regularisation. However, below we construct a simple counterexample to (4). Its existence shows that in general $E_\kappa \neq E_{c,\text{PPT}}^{\text{exact}}$, thus invalidating the efficiently computable solution claimed in [25]. The connection between the quantity E_κ and the PPT entanglement cost revealed in [25, 26] will still prove useful to us, albeit *a priori* it

is not clear how it could lead to a computable formula for $E_{c,\text{PPT}}^{\text{exact}}$ — the daunting problem of regularisation persists. Further details about the claims of [25, 26] can be found in the erratum [37].

Main results. — In this work, we completely solve the problem of calculating $E_{c,\text{PPT}}^{\text{exact}}$ through an entirely different approach. Namely, we construct two converging hierarchies of semi-definite programs that can be used to calculate $E_{c,\text{PPT}}^{\text{exact}}$ for any given state to any degree of precision *efficiently*, i.e. in time polynomial in the underlying Hilbert space dimension and in $\log(1/\varepsilon)$, with ε being the additive error. The key quantities in our approach are a family of PPT entanglement monotones indexed by an integer $p \in \mathbb{N}$ and given by

$$E_{\chi,p}(\rho) := \log_2 \chi_p(\rho), \quad (5)$$

where

$$\chi_p(\rho) := \min \left\{ \text{Tr } S_p : -S_i \leq S_{i-1}^\Gamma \leq S_i, \ i=0, \dots, p, \ S_{-1} = \rho \right\} \quad (6)$$

is an SDP with variables S_0, \dots, S_p . These quantities are increasing in p for every fixed ρ , and we refer to them as the χ -*hierarchy*. Note also that $E_{\chi,0} = E_N$, hence $E_{\chi,p}$ can be regarded as a generalisation of the entanglement negativity [32, 33].

The functions $E_{\chi,p}$ can be related to the quantity E_κ studied in [25, 26] and its generalisations. In particular, by exploiting the connection between the entanglement cost $E_{c,\text{PPT}}^{\text{exact}}$ and a regularised form of E_κ shown in [25, 26], the χ -hierarchy can be shown to approximate the entanglement cost $E_{c,\text{PPT}}^{\text{exact}}$ from below, in the sense that $E_{c,\text{PPT}}^{\text{exact}}(\rho) \geq E_{\chi,p}(\rho)$ for all states and all p . Our first main result establishes that this approximation becomes increasingly tight as p increases, and the χ -hierarchy gives the value of $E_{c,\text{PPT}}^{\text{exact}}$ *exactly* in the limit $p \rightarrow \infty$. This allows us to replace the limit in the number of copies n , which is what makes $E_{c,\text{PPT}}^{\text{exact}}$ difficult to compute, with a limit in the hierarchy level p . This already provides a single-letter formula for the PPT entanglement cost that no longer suffers from the curse of regularisation. However, because of the limiting procedure $p \rightarrow \infty$, it is still unclear if the expression can be evaluated easily. Crucially, we show that calculating the limit of the χ -hierarchy is indeed significantly easier than evaluating regularised expressions: the convergence to the true value of $E_{c,\text{PPT}}^{\text{exact}}$ is *exponentially fast* uniformly on all states, which opens the way to an accurate calculation of $E_{c,\text{PPT}}^{\text{exact}}$ in practice.

Theorem 1. *For all bipartite states $\rho = \rho_{AB}$ on a system of minimal local dimension $d := \min\{|A|, |B|\} \geq 2$, and all positive integers $p \in \mathbb{N}^+$, it holds that*

$$0 \leq E_{c,\text{PPT}}^{\text{exact}}(\rho) - E_{\chi,p}(\rho) \leq \log_2 \frac{1}{1 - (1 - \frac{2}{d})^p}, \quad (7)$$

entailing that

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = \lim_{p \rightarrow \infty} E_{\chi,p}(\rho). \quad (8)$$

Note that for every fixed value of d and for large p , the approximation error on the right-hand side of (7) can be estimated as $(1 - \frac{2}{d})^p (\log_2 e)$. In other words, the speed of convergence in (8) is exponential in p and furthermore independent of ρ .

The single-letter formula (8) can be used to establish two notable properties of the zero-error PPT entanglement cost $E_{c,\text{PPT}}^{\text{exact}}$, namely additivity and continuity [38, § IV D]. Also, Theorem 1 yields immediately a simple solution in the qubit-qudit case ($d = 2$), generalising Ishizaka's result that $E_{c,\text{PPT}}^{\text{exact}} = E_N$ for all two-qubit states [30]:

Corollary 2. *For all states $\rho = \rho_{AB}$ on a $2 \times n$ bipartite quantum system,*

$$\begin{aligned} E_{c,\text{PPT}}^{\text{exact}}(\rho) &= E_{\chi,1}(\rho) \\ &= \log_2 \min \{ \|S_0^\Gamma\|_1 : -S_0 \leq \rho^\Gamma \leq S_0 \}. \end{aligned} \quad (9)$$

But the most important implication of Theorem 1 is that it allows us to construct an efficient algorithm that calculates $E_{c,\text{PPT}}^{\text{exact}}$ accurately.

Theorem 3 (Efficient algorithm to compute the cost). *There exists an algorithm that computes $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ up to an additive error ε in time*

$$\mathcal{O}\left((dD)^{6+o(1)} \text{polylog}(1/\varepsilon)\right) \quad (10)$$

for an arbitrary bipartite state ρ on a system of total dimension D and minimal local dimension d .

This result is a first of its kind for two distinct reasons:

(a) First, because it establishes the efficient computability of an operationally meaningful asymptotic entanglement measure (i.e. a distillable entanglement or an entanglement cost). There is no known algorithm to estimate any other such measure, not even under the simplifying zero-error assumption.

(b) Second, because efficient computability is shown without exhibiting a closed-form single-letter formula, but rather by describing converging SDP hierarchies. To the extent of our knowledge, the only other case in quantum information theory where a similar situation arises is in [39, Theorem 5.1]. However, unlike ours, the algorithm described there is computationally extremely expensive, featuring an exponential dependence on d^3/ε . More generally, expressing difficult-to-compute quantities through converging SDP hierarchies is a technical tool that has found various uses in quantum information [40–45], but such results typically do not result in efficiently computable algorithms or do not yield exact operational results.

Our algorithm is simple to describe: the first step is to use (7) to find p large enough so that $E_{\chi,p}(\rho)$ approximates $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ up to an additive error $\varepsilon/2$. It turns out that $p = \mathcal{O}(d \log(d/\varepsilon))$ suffices. We then solve the SDP that computes $E_{\chi,p}(\rho)$ with a precision $\varepsilon/2$ and estimate the corresponding time complexity. Doing so yields an ε -approximation of $E_{c,\text{PPT}}^{\text{exact}}(\rho)$. The key observation in the analysis is that climbing the χ -hierarchy up to level p introduces only polynomially many more constraints in the SDP and is thus relatively inexpensive. Indeed, the time complexity of the above algorithm, reported in (10), is only marginally larger than that of computing the entanglement negativity itself, which is $\mathcal{O}(D^{6+o(1)} \text{polylog}(1/\varepsilon))$.

Methods. — *The task.* We start by defining $E_{c,\text{PPT}}^{\text{exact}}$ in rigorous terms. A (quantum) channel $\Lambda : X \rightarrow Y$ is a completely positive and trace preserving map taking as input states of a quantum system X and outputting states of Y . The set of completely positive maps (respectively, quantum channels) from X to Y will be denoted as $\text{CP}(X \rightarrow Y)$ (respectively, $\text{CPTP}(X \rightarrow Y)$). If $X = AB$ and $Y = A'B'$ are both bipartite systems and $\Lambda \in \text{CP}(AB \rightarrow A'B')$, we say that Λ is **PPT** if $\Gamma_{B'} \circ \Lambda \circ \Gamma_B$ is still completely positive, where Γ_B denotes the partial transpose on B , and analogously for $\Gamma_{B'}$. We can then define the **zero-error PPT entanglement cost** of any bipartite state $\rho = \rho_{AB}$ as

$$\begin{aligned} E_{c,\text{PPT}}^{\text{exact}}(\rho) &:= \inf \left\{ R : \text{for all sufficiently large } n \in \mathbb{N} \right. \\ &\quad \left. \exists \Lambda_n \in \text{PPT} \cap \text{CPTP} : \Lambda_n(\Phi_2^{\otimes \lfloor Rn \rfloor}) = \rho^{\otimes n} \right\}. \end{aligned} \quad (11)$$

Here, $\Phi_2 := |\Phi_2\rangle\langle\Phi_2|$, where $|\Phi_2\rangle := \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ is the two-qubit maximally entangled state, i.e. the **ebit**, and Λ_n is required to be a PPT channel.

The above task is of interest because it constitutes a simplification of fully fledged entanglement dilution (cf. (1)). In particular: (I) The mathematically hard-to-characterise set of LOCC [46] is replaced by its outer approximation given by the set of PPT operations. Because of this connection to LOCC, PPT operations have been studied extensively in entanglement theory [47–50]. (II) Dilution with asymptotically vanishing error (which leads to entropic measures such as that in (1)) is replaced with exact, zero-error dilution. However, in the SM [38] we show that no substantial change occurs if we require that the error, instead of being exactly zero, decay to zero sufficiently fast — the relevant figure of merit is then still $E_{c,\text{PPT}}^{\text{exact}}$.

The quantifier E_κ . Wang and Wilde [25, 26, 37] introduced and studied the SDP-computable quantity

$$E_\kappa(\rho) := \log_2 \min \{ \text{Tr } S : -S \leq \rho^\Gamma \leq S, S^\Gamma \geq 0 \}. \quad (12)$$

Among other things, they showed that: (i) E_κ is monotonically non-increasing under PPT channels; (ii) $E_\kappa(\rho) \geq$

$E_N(\rho)$ for all states ρ , with equality when ρ has zero bi-negativity; (iii) E_κ is sub-additive, meaning that

$$E_\kappa(\rho \otimes \rho') \leq E_\kappa(\rho) + E_\kappa(\rho') \quad (13)$$

for all pairs of states $\rho = \rho_{AB}$ and $\rho' = \rho'_{A'B'}$; and (iv) its regularisation yields the zero-error PPT entanglement cost, i.e.

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_\kappa^\infty(\rho) := \lim_{n \rightarrow \infty} \frac{1}{n} E_\kappa(\rho^{\otimes n}). \quad (14)$$

As said, it was claimed in [25] that E_κ is additive, meaning that equality holds in (13). However, this claim is incorrect (see also [37]). To disprove it, it is useful to first note that additivity indeed holds when both ρ and ρ' have zero bi-negativity, simply because in that case E_κ coincides with the entanglement negativity by property (ii), and this latter measure is additive. Hence, our search for a counterexample must start with the construction of states with non-zero bi-negativity. That such states do exist was reported already in [29, 30] based on numerical evidence. However, here we present a simpler, analytical construction.

Punch card states. Let $A \geq 0$ be a positive semi-definite $d \times d$ matrix, and let Q be another $d \times d$ symmetric matrix with only 0/1 entries. The associated **punch card state** is the bipartite quantum state on $\mathbb{C}^d \otimes \mathbb{C}^d$ defined by

$$\pi_{A,Q} := \frac{1}{N_{A,Q}} \left(\sum_{i,j} A_{ij} |ii\rangle\langle jj| + \sum_{i \neq j} Q_{ij} |A_{ij}\rangle\langle ij| \right), \quad (15)$$

where $N_{A,Q}$ is chosen so that $\text{Tr } \pi_{A,Q} = 1$. It is straightforward to verify that

$$|\pi_{A,Q}^\Gamma|^\Gamma = \frac{1}{N_{A,Q}} \left(\sum_{i,j} Q_{ij} A_{ij} |ii\rangle\langle jj| + \sum_{i \neq j} |A_{ij}\rangle\langle ij| \right). \quad (16)$$

Therefore, if A and Q are chosen such that $Q \circ A \not\geq 0$, where \circ denotes the Hadamard (i.e. entry-wise) product between matrices, $\pi_{A,Q}$ will have non-zero bi-negativity. It is easy to come up with examples of A and Q that meet the above criteria, the simplest one being

$$A_0 := \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad Q_0 := \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}. \quad (17)$$

Having constructed a state with non-zero bi-negativity, we can wonder whether two copies of it already violate the additivity of E_κ . And sure enough, they do:

Lemma 4. *The punch card state $\pi_0 := \pi_{A_0,Q_0}$ defined by (15) with the substitution (17) satisfies $|\pi_0^\Gamma|^\Gamma \not\geq 0$, and moreover*

$$1.001 \approx E_\kappa(\pi_0^{\otimes 2}) < 2E_\kappa(\pi_0) \approx 1.029. \quad (18)$$

In particular, the sub-additivity of E_κ and Lemma 4 imply that $E_{c,\text{PPT}}^{\text{exact}}(\pi_0) \leq \frac{1}{2} E_\kappa(\pi_0^{\otimes 2}) < E_\kappa(\pi_0)$ and therefore that $E_{c,\text{PPT}}^{\text{exact}}(\pi_0) \neq E_\kappa(\pi_0)$, thus invalidating the main claims of [25, 26].

Two SDP hierarchies. In (5)–(6) we have already introduced the χ -hierarchy as a generalisation of the entanglement negativity. We now introduce another complementary hierarchy of SDPs, the **κ -hierarchy**, defined for $q \in \mathbb{N}^+$ (note that $q \geq 1$) by

$$E_{\kappa,q}(\rho) := \log_2 \kappa_q(\rho), \quad (19)$$

with

$$\kappa_q(\rho) := \min \left\{ \text{Tr } S_{q-1}: -S_i \leq S_{i-1}^\Gamma \leq S_i, \ i=0, \dots, q-1, \right. \\ \left. S_{-1} = \rho, \ S_{q-1}^\Gamma \geq 0 \right\}. \quad (20)$$

Observe the resemblance to the definition of χ_p in (6): the only difference between the two optimisations is the condition $S_{p-1}^\Gamma \geq 0$; adding that to (6) yields immediately (20) with $q \mapsto p$, and indeed in that case the optimal S_p would automatically be $S_p = S_{p-1}^\Gamma$. Furthermore, $E_{\kappa,1} = E_\kappa$ coincides with the quantity (12) introduced by Wang and Wilde, of which the κ -hierarchy thus constitutes a generalisation.

In the Supplemental Material [38] we explore the properties of the quantities $E_{\chi,p}$ and $E_{\kappa,q}$, showing them to be legitimate entanglement measures. In particular, the functions are all suitably normalised, continuous, faithful on PPT states, and strongly monotonic under PPT operations. The pivotal property that distinguishes $E_{\chi,p}$ from $E_{\kappa,q}$ is that, while the quantities $E_{\kappa,q}$ are only sub-additive, the χ -quantities are fully additive under tensor products, meaning that regularisation can always be avoided.

Two key insights lead to the proof of our main results. First, that the two hierarchies provide complementary bounds on $E_{c,\text{PPT}}^{\text{exact}}$, the fundamental quantity we want to estimate: namely, the χ -hierarchy gives increasing lower bounds on $E_{c,\text{PPT}}^{\text{exact}}$, while the κ -hierarchy gives decreasing upper bounds on it. In other words, on any fixed state ρ

$$E_{\chi,0} \leq E_{\chi,1} \leq \dots \leq E_{c,\text{PPT}}^{\text{exact}} = E_\kappa^\infty \leq \dots \leq E_{\kappa,2} \leq E_{\kappa,1}. \quad (21)$$

In particular, $E_{\chi,p}$ is increasing in p , while $E_{\kappa,q}$ is decreasing in q . Eq. (21) immediately shows the remarkable connection between two very different limits: one in the number of copies n , which is needed to compute E_κ^∞ (see Eq. (14)), and one in the hierarchy levels p and q .

The second insight is that there is a connection between the χ - and κ -hierarchies, as expressed by the following key technical result, proven in the SM [38].

Proposition 5. *For all states $\rho = \rho_{AB}$ on a system of minimal local dimension $d := \min\{|A|, |B|\} \geq 2$, and all $p \in \mathbb{N}^+$,*

$$\kappa_p(\rho) \leq \frac{d}{2} \chi_p(\rho) - \left(\frac{d}{2} - 1 \right) \chi_{p-1}(\rho). \quad (22)$$

With Proposition 5 at hand, we can now see how it implies our two main results, Theorems 1 and 3.

Proof sketch of Theorem 1. Combining (21) and (22) shows that

$$2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \leq \kappa_p(\rho) \leq \frac{d}{2} \chi_p(\rho) - \left(\frac{d}{2} - 1\right) \chi_{p-1}(\rho). \quad (23)$$

The quantity that we are really interested in, however, is the normalised difference between $\chi_p(\rho)$ and its claimed limiting value $2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)}$. To see what the above inequality tells us in this respect, we can define the quantity

$$\varepsilon_p(\rho) := 1 - \frac{\chi_p(\rho)}{2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)}}, \quad (24)$$

by means of which (23) can be cast as

$$\varepsilon_p(\rho) \leq \left(1 - \frac{2}{d}\right) \varepsilon_{p-1}(\rho). \quad (25)$$

Iterating the above relation gives

$$\varepsilon_p(\rho) \leq \left(1 - \frac{2}{d}\right)^p \varepsilon_0(\rho) \leq \left(1 - \frac{2}{d}\right)^p, \quad (26)$$

which entails (7) after elementary algebraic manipulations. Taking the limit $p \rightarrow \infty$ in (7) proves also (8). \square

Proof sketch of Theorem 3. It suffices to formalise the qualitative argument provided below the statement of the theorem. For $d > 2$, we first choose

$$p_d := \left\lceil \frac{\log_2(2d/\varepsilon)}{-\log_2(1 - \frac{2}{d})} \right\rceil = \mathcal{O}(d \log(d/\varepsilon)), \quad (27)$$

so that, with the notation in (24), $\varepsilon_{p_d}(\rho) \leq \frac{\varepsilon}{2d}$. Using $\ln(2)|a - b| \leq |2^a - 2^b|$, valid for $a, b \geq 0$, we get

$$0 \leq E_{c,\text{PPT}}^{\text{exact}}(\rho) - E_{\chi,p_d}(\rho) \leq \frac{2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \varepsilon}{2d \ln 2} \leq \frac{\varepsilon}{2 \ln 2}, \quad (28)$$

where the last inequality is a consequence of the fact that every state can be created via a quantum teleportation protocol — and hence with PPT operations — from a maximally entangled state, which entails that $E_{c,\text{PPT}}^{\text{exact}}(\rho) \leq \log_2 d$ for all ρ . We then solve the SDP for $\chi_{p_d}(\rho)$ up to an additive error $(\ln 2 - 1/2) \varepsilon$ by running an optimised SDP solver [51, 52]. Doing so yields an approximation of $E_{\chi,p_d}(\rho)$ up to an additive error $(1 - \frac{1}{2 \ln 2}) \varepsilon$. Adding this up with the error in (28) yields a total error of ε . The time complexity in (10) can be calculated using known theoretical bounds on the complexity of SDPs, e.g. those found in [51]. \square

Discussion and conclusions. In this paper, we have provided a computationally complete, single-letter solution to the problem of calculating the zero-error PPT entanglement cost of arbitrary (finite-dimensional) quantum states. To the best of our knowledge, it is the first time that any operational asymptotic entanglement measure is shown to be efficiently computable. A particularly interesting feature of our construction is that it does not rely on a closed-form formula, but rather on two converging hierarchies of semi-definite programs that approximate the cost from above and below with controllable error.

An open question in our analysis is whether the χ -hierarchy collapses at any finite level for some — or even all — states. We found examples of states ρ such that $E_N(\rho) = E_{\chi,0}(\rho) < E_{\chi,1}(\rho)$ and also $E_{\chi,1}(\rho) < E_{\chi,2}(\rho)$, but we were not able to ascertain whether there exists in general a gap between $E_{\chi,2}$ and $E_{\chi,3}$. If $E_{\chi,2} = E_{\chi,3}$ holds in general, then this would mean that $E_{c,\text{PPT}}^{\text{exact}} = E_{\chi,2}$, and thus the cost could be computed with a simple single-letter formula. While this would be a considerable simplification from the analytical standpoint, we stress that it will only entail a $\text{poly}(d)$ improvement in the time complexity of evaluating it numerically.

Acknowledgements. L.L. thanks Andreas Winter for enlightening correspondence on zero-error information theory, as well as the organisers and attendees of the workshop ‘Quantum information’ (Les Diablerets, Switzerland, 25 February–1 March 2024), where this work was first presented, for stimulating and entertaining discussions. L.L. and B.R. thank Xin Wang and Mark M. Wilde for feedback on the first draft of this paper.

Note. The issue with the original argument by Wang and Wilde that leads to the additivity violation for E_{κ} (Lemma 4) is discussed in detail in the erratum [37].

* ludovico.lami@gmail.com

† francesco.mele@sns.it

‡ bartosz.regula@gmail.com

- [1] A. S. Holevo, *The capacity of the quantum channel with general signal states*, *IEEE Trans. Inf. Theory* **44**, 269 (1998). 1
- [2] B. Schumacher and M. D. Westmoreland, *Sending classical information via noisy quantum channels*, *Phys. Rev. A* **56**, 131 (1997). 1
- [3] S. Lloyd, *Capacity of the noisy quantum channel*, *Phys. Rev. A* **55**, 1613 (1997). 1
- [4] P. Shor, *The quantum channel capacity and coherent information*, Lecture notes, MSRI Workshop on Quantum Computation (2002).
- [5] I. Devetak, *The private classical capacity and quantum capacity of a quantum channel*, *IEEE Trans. Inf. Theory* **51**, 44 (2005). 1
- [6] N. Cai, A. Winter, and R. W. Yeung, *Quantum privacy and quantum wiretap channels*, *Probl. Inf. Transm.* **40**, 318 (2004). 1

- [7] C. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal, *Entanglement-assisted classical capacity of noisy quantum channels*, *Phys. Rev. Lett.* **83**, 3081 (1999). 1
- [8] C. H. Bennett, P. W. Shor, J. A. Smolin, and A. V. Thapliyal, *Entanglement-assisted capacity of a quantum channel and the reverse Shannon theorem*, *IEEE Trans. Inf. Theory* **48**, 2637 (2002). 1
- [9] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, *Concentrating partial entanglement by local operations*, *Phys. Rev. A* **53**, 3824 (1996). 1
- [10] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, *Purification of noisy entanglement and faithful teleportation via noisy channels*, *Phys. Rev. Lett.* **76**, 722 (1996).
- [11] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, *Mixed-state entanglement and quantum error correction*, *Phys. Rev. A* **54**, 3824 (1996). 1
- [12] I. Devetak and A. Winter, *Distillation of secret key and entanglement from quantum states*, *Proc. Royal Soc. A* **461**, 207 (2005).
- [13] L. Lami and B. Regula, *Distillable entanglement under dually non-entangling operations*, *Preprint arXiv:2307.11008* (2023). 1
- [14] P. M. Hayden, M. Horodecki, and B. M. Terhal, *The asymptotic entanglement cost of preparing a quantum state*, *J. Phys. A* **34**, 6891 (2001). 1
- [15] H. Yamasaki, K. Kuroiwa, P. Hayden, and L. Lami, *Entanglement cost for infinite-dimensional physical systems*, *Preprint arXiv:2401.09554* (2024). 1
- [16] K. G. H. Vollbrecht and R. F. Werner, *Entanglement measures under symmetry*, *Phys. Rev. A* **64**, 062307 (2001). 1
- [17] P. W. Shor, *Equivalence of additivity questions in quantum information theory*, *Commun. Math. Phys.* **246**, 473 (2004).
- [18] P. Hayden and A. Winter, *Counterexamples to the maximal p -norm multiplicativity conjecture for all $p > 1$* , *Commun. Math. Phys.* **284**, 263 (2008).
- [19] T. Cubitt, A. W. Harrow, D. Leung, A. Montanaro, and A. Winter, *Counterexamples to additivity of minimum output p -Rényi entropy for p close to 0*, *Commun. Math. Phys.* **284**, 281 (2008).
- [20] M. B. Hastings, *Superadditivity of communication capacity using entangled inputs*, *Nat. Phys.* **5**, 255 (2009).
- [21] G. Smith and J. Yard, *Quantum communication with zero-capacity channels*, *Science* **321**, 1812 (2008). 1
- [22] W. K. Wootters, *Entanglement of formation of an arbitrary state of two qubits*, *Phys. Rev. Lett.* **80**, 2245 (1998). 1
- [23] T. Cubitt, D. Elkouss, W. Matthews, M. Ozols, D. Pérez-García, and S. Strelchuk, *Unbounded number of channel uses may be required to detect quantum capacity*, *Nat. Commun.* **6**, 6739 (2015). 1
- [24] K. Audenaert, M. B. Plenio, and J. Eisert, *Entanglement cost under positive-partial-transpose-preserving operations*, *Phys. Rev. Lett.* **90**, 027901 (2003). 1, 2, 3
- [25] X. Wang and M. M. Wilde, *Cost of quantum entanglement simplified*, *Phys. Rev. Lett.* **125**, 040502 (2020). 2, 3, 4, 5, 6, 7
- [26] X. Wang and M. M. Wilde, *Exact entanglement cost of quantum states and channels under positive-partial-transpose-preserving operations*, *Phys. Rev. A* **107**, 012429 (2023). 2, 3, 4
- [27] G. Gour and C. M. Scandolo, *Dynamical entanglement*, *Phys. Rev. Lett.* **125**, 180505 (2020).
- [28] G. Gour and C. M. Scandolo, *Entanglement of a bipartite channel*, *Phys. Rev. A* **103**, 062422 (2021). 1
- [29] K. Audenaert, B. De Moor, K. G. H. Vollbrecht, and R. F. Werner, *Asymptotic relative entropy of entanglement for orthogonally invariant states*, *Phys. Rev. A* **66**, 032310 (2002). 2, 4
- [30] S. Ishizaka, *Binegativity and geometry of entangled states in two qubits*, *Phys. Rev. A* **69**, 020301 (2004). 2, 3, 4, 18
- [31] A. Peres, *Separability criterion for density matrices*, *Phys. Rev. Lett.* **77**, 1413 (1996). 2
- [32] G. Vidal and R. F. Werner, *Computable measure of entanglement*, *Phys. Rev. A* **65**, 032314 (2002). 2, 6, 7, 13
- [33] M. B. Plenio, *Logarithmic negativity: A full entanglement monotone that is not convex*, *Phys. Rev. Lett.* **95**, 090503 (2005). 2, 6
- [34] J. Watrous, *The Theory of Quantum Information* (Cambridge University Press, 2018). 2, 9
- [35] P. Skrzypczyk and D. Cavalcanti, *Semidefinite Programming in Quantum Information Science* (IOP Publishing, 2023). 4
- [36] L. Vandenberghe and S. Boyd, *Semidefinite programming*, *SIAM Rev.* **38**, 49 (1996). 2, 4, 9
- [37] X. Wang and M. M. Wilde, *Errata for “Cost of quantum entanglement simplified” and “Exact entanglement cost of quantum states and channels under PPT-preserving operations”*, preprint <https://doi.org/10.5281/zenodo.11061607> (2024). 2, 3, 4, 5
- [38] See the Supplemental Material. 3, 4
- [39] H. Fawzi and O. Fawzi, *Defining quantum divergences via convex optimization*, *Quantum* **5**, 387 (2021). 3
- [40] A. C. Doherty, P. A. Parrilo, and F. M. Spedalieri, *Complete family of separability criteria*, *Phys. Rev. A* **69**, 022308 (2004). 3
- [41] M. Navascués, S. Pironio, and A. Acín, *A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations*, *New J. Phys.* **10**, 073013 (2008).
- [42] A. W. Harrow, A. Natarajan, and X. Wu, *An improved semidefinite programming hierarchy for testing entanglement*, *Commun. Math. Phys.* **352**, 881 (2017).
- [43] K. Fang and H. Fawzi, *Geometric Rényi divergence and its applications in quantum channel capacities*, *Commun. Math. Phys.* **384**, 1615 (2021).
- [44] O. Fawzi, A. Shayeghi, and H. Ta, *A hierarchy of efficient bounds on quantum capacities exploiting symmetry*, *IEEE Trans. Inf. Theory* **68**, 7346 (2022).
- [45] M. Berta, F. Borderi, O. Fawzi, and V. B. Scholz, *Semidefinite programming hierarchies for constrained bilinear optimization*, *Math. Program.* **194**, 781 (2022). 3
- [46] E. Chitambar, D. Leung, L. Mančinska, M. Ozols, and A. Winter, *Everything you always wanted to know about LOCC (but were afraid to ask)*, *Commun. Math. Phys.* **328**, 303 (2014). 3
- [47] E. M. Rains, *Bound on distillable entanglement*, *Phys. Rev. A* **60**, 179 (1999). 3
- [48] E. M. Rains, *A semidefinite program for distillable entanglement*, *IEEE Trans. Inf. Theory* **47**, 2921 (2001).
- [49] T. Eggeling, K. G. H. Vollbrecht, R. F. Werner, and M. M. Wolf, *Distillability via protocols respecting the positivity of partial transpose*, *Phys. Rev. Lett.* **87**, 257902 (2001).
- [50] X. Wang and R. Duan, *Irreversibility of asymptotic entanglement manipulation under quantum operations completely preserving positivity of partial transpose*, *Phys. Rev. Lett.* **119**, 180506 (2017). 3
- [51] Y. T. Lee, A. Sidford, and S. C.-W. Wong, *A faster cutting plane method and its implications for combinatorial and convex optimization*, in *Proc. 56th Annual Symp. Found. Computer Science* (2015) pp. 1049–1065. 5, 20
- [52] J. van Apeldoorn, A. Gilyén, S. Gribling, and R. de Wolf,

- Quantum SDP-Solvers: Better upper and lower bounds*, [Quantum](#) **4**, 230 (2020). 5, 20
- [53] R. F. Werner, *Quantum states with Einstein-Podolsky-Rosen correlations admitting a hidden-variable model*, [Phys. Rev. A](#) **40**, 4277 (1989). 2
 - [54] M. Horodecki, P. Horodecki, and R. Horodecki, *Mixed-state entanglement and distillation: Is there a “bound” entanglement in nature?* [Phys. Rev. Lett.](#) **80**, 5239 (1998). 2
 - [55] D. Bruß and A. Peres, *Construction of quantum states with bound entanglement*, [Phys. Rev. A](#) **61**, 030301 (2000). 2
 - [56] M. Fekete, *Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten*, [Math. Z.](#) **17**, 228 (1923). 3
 - [57] M. A. Nielsen, *Conditions for a class of entanglement transformations*, [Phys. Rev. Lett.](#) **83**, 436 (1999). 9
 - [58] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, *Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels*, [Phys. Rev. Lett.](#) **70**, 1895 (1993). 9
 - [59] S. Khatry and M. M. Wilde, *Principles of quantum communication theory: A modern approach*, preprint [arXiv:2011.04672](#) (2020). 10
 - [60] M. Horodecki, *Entanglement measures*, [Quantum Inf. Comput.](#) **1**, 3 (2001). 13
 - [61] M. J. Donald, M. Horodecki, and O. Rudolph, *The uniqueness theorem for entanglement measures*, [J. Math. Phys.](#) **43**, 4252 (2002). 13
 - [62] L. Lami and B. Regula, *No second law of entanglement manipulation after all*, [Nat. Phys.](#) **19**, 184 (2023). 13
 - [63] G. Vidal and R. Tarrach, *Robustness of entanglement*, [Phys. Rev. A](#) **59**, 141 (1999). 15
 - [64] C. Shannon, *The zero error capacity of a noisy channel*, [IRE Trans. Inf. Theory](#) **2**, 8 (1956). 22

Supplemental Material

CONTENTS

I. Introduction	1
A. Quantum states and channels	1
B. Positive partial transpose	2
C. Zero-error PPT entanglement cost	3
II. Violations of additivity of E_κ	3
III. New entanglement monotones: $E_{\chi,p}$ and $E_{\kappa,q}$	4
A. Rationale of our construction	4
B. Relation between the χ - and κ -hierarchies	6
C. $E_{\chi,p}$ and $E_{\kappa,q}$ as PPT monotones	8
D. Dual expressions	9
E. Additivity properties	10
F. Continuity	12
IV. Main results	13
A. Enter the regularisation	13
B. A key technical result: proof of Proposition 5	15
C. What have we achieved so far?	17
Convergence	17
The local qubit case	17
D. Exponential convergence: proof of Theorem 1	18
E. An explicit algorithm to compute the zero-error PPT cost: proof of Theorem 3	19
F. Open problem: hierarchy collapse	22
V. From zero error to very small error	22

I. INTRODUCTION

A. Quantum states and channels

The quantum systems we consider in this work are all represented by finite-dimensional Hilbert spaces \mathcal{H} . We will denote the set of linear operators on \mathcal{H} by $\mathcal{L}(\mathcal{H})$. We occasionally refer to these operators also as matrices, to underline the fact that they are understood to be finite dimensional.

Quantum states pertaining to a quantum system with Hilbert space \mathcal{H} are represented by *density operators*, i.e. positive semi-definite operators on \mathcal{H} with unit trace. We will denote the set of positive semi-definite operators on \mathcal{H} by $\mathcal{L}_+(\mathcal{H})$. Given two Hermitian operators $X, Y \in \mathcal{L}_+(\mathcal{H})$ we write $X \leq Y$ if $Y - X \in \mathcal{L}_+(\mathcal{H})$, and $X < Y$ if $Y - X$ is strictly positive definite. The relations \geq and $>$ are defined similarly.

The *trace norm* of an arbitrary operator $X \in \mathcal{L}(\mathcal{H})$ is given by

$$\|X\|_1 := \text{Tr} \sqrt{X^\dagger X}, \quad (\text{S1})$$

where for a positive semi-definite operator A its square root is constructed as the unique semi-definite operator \sqrt{A} such that $\sqrt{A}^2 = A$. If $X = X^\dagger$ is Hermitian, then it is possible to give the following simple characterisation of its trace norm. We include a proof for the sake of completeness, as the lemma below will be used a few times throughout this paper.

Lemma S1 (Variational characterisation of the trace norm). *Let $X = X^\dagger \in \mathcal{L}(\mathcal{H})$ be a Hermitian matrix. Then*

$$\|X\|_1 = \min \{ \text{Tr } Y : -Y \leq X \leq Y \}. \quad (\text{S2})$$

Proof. Let $X = \sum_i x_i |i\rangle\langle i|$ be the spectral decomposition of X . Then by inspection we find $\|X\|_1 = \sum_i |x_i|$. On the one hand, setting $Y = |X| := \sum_i |x_i| |i\rangle\langle i|$ we find that $\min \{ \text{Tr } Y : -Y \leq X \leq Y \} \leq \text{Tr } \sum_i |x_i| |i\rangle\langle i| = \|X\|_1$. On the other, for an arbitrary Y such that $-Y \leq X \leq Y$ we have that $-\langle i|Y|i\rangle \leq \langle i|X|i\rangle = x_i \leq \langle i|Y|i\rangle$, implying that $\langle i|Y|i\rangle \geq |x_i|$. Therefore, $\text{Tr } Y = \sum_i \langle i|Y|i\rangle \geq \sum_i |x_i| = \|X\|_1$; since Y was arbitrary, this concludes the proof. \square

Given two quantum systems with Hilbert spaces \mathcal{H} and \mathcal{H}' , a linear map $\Lambda : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H}')$ is called **positive** if $\Lambda(\mathcal{L}_+(\mathcal{H})) \subseteq \mathcal{L}_+(\mathcal{H}')$, and **completely positive** if $I_k \otimes \Lambda$ is positive for all $k \in \mathbb{N}$, where I_k is the identity map on $\mathcal{L}(\mathbb{C}^k)$. It is well known that complete positivity can be formulated as a positive semi-definite constraint using the formalism of Choi states. Namely, introducing the maximally entangled state

$$\Phi_d := |\Phi_d\rangle\langle\Phi_d|, \quad |\Phi_d\rangle := \frac{1}{\sqrt{d}} \sum_{i=1}^d |ii\rangle, \quad (\text{S3})$$

a linear map $\Lambda : \mathcal{L}(\mathbb{C}^d) \rightarrow \mathcal{L}(\mathbb{C}^{d'})$ is completely positive if and only if

$$(I_d \otimes \Lambda)(\Phi_d) \geq 0 \quad (\text{S4})$$

is a positive semi-definite operator.

A completely positive map and trace-preserving map $\Lambda : \mathcal{L}(\mathcal{H}) \rightarrow \mathcal{L}(\mathcal{H}')$ is called a **quantum channel**. In what follows, $\text{CP}(X \rightarrow Y)$ and $\text{CPTP}(X \rightarrow Y)$ will denote respectively the set of completely positive maps and the set of quantum channels between two quantum systems X and Y .

B. Positive partial transpose

Composite quantum systems are represented by the tensor product of the local Hilbert spaces; for the simplest case of a bipartite system, this is expressed in formula as $\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B$. In what follows, we will often need to consider the minimal local dimension of a bipartite quantum system AB , i.e. the number $d = \min\{|A|, |B|\}$, where we denote by $|A|$ (respectively, $|B|$) the dimension of the local Hilbert space \mathcal{H}_A (respectively, \mathcal{H}_B).

Given an arbitrary bipartite quantum system AB , the **partial transpose** on B is the linear map $\Gamma_B : \mathcal{L}(\mathcal{H}_{AB}) \rightarrow (\mathcal{H}_{AB})$ uniquely defined by

$$\Gamma_B(X_A \otimes Y_B) = (X_A \otimes Y_B)^\Gamma := X_A \otimes Y_B^\top \quad \forall X_A \in \mathcal{L}(\mathcal{H}_A), Y_B \in \mathcal{L}(\mathcal{H}_B). \quad (\text{S5})$$

Here, \top denotes the transposition with respect to a fixed basis of \mathcal{H}_B . Note that

$$\text{Tr } Z_{AB} W_{AB} = \text{Tr } Z_{AB}^\Gamma W_{AB}^\Gamma \quad \forall Z, W \in \mathcal{L}(\mathcal{H}_{AB}), \quad (\text{S6})$$

and in particular

$$\text{Tr } Z_{AB}^\Gamma = \text{Tr } Z_{AB} \quad \forall Z \in \mathcal{L}(\mathcal{H}_{AB}). \quad (\text{S7})$$

The definition of Γ_B does depend on which basis we choose, but all choices are unitarily equivalent to each other, in the sense that for any two different partial transpositions $\Gamma, \tilde{\Gamma}$ there exists a unitary U_B acting on B such that

$$\tilde{\Gamma}_B(Z_{AB}) = U_B \Gamma_B(Z_{AB}) U_B^\dagger. \quad (\text{S8})$$

A state σ_{AB} on a bipartite quantum system AB is said to be a **PPT state** if $\sigma_{AB}^\Gamma = \Gamma_B(\sigma_{AB}) \geq 0$. The PPT condition is important in quantum information because it provides a relaxation of separability: any **separable** (i.e. unentangled) state σ_{AB} , which can thus be decomposed as $\sigma_{AB} = \sum_x p_x \alpha_x^A \otimes \beta_x^B$ for some probability distribution p and some sets of local states α_x^A and β_x^B [53], is PPT, but the converse is famously not true [54, 55].

A completely positive map $\Lambda \in \text{CP}(AB \rightarrow A'B')$ with bipartite quantum systems AB and $A'B'$ as input and output is called **PPT** if

$$\Gamma_{B'} \circ \Lambda \circ \Gamma_B \in \text{CP}(AB \rightarrow A'B'). \quad (\text{S9})$$

If Λ is in addition also trace preserving, then we call it a **PPT channel** (or a **PPT operation**). Such maps are sometimes also referred to as ‘completely PPT preserving’. The importance of the set of PPT channels is that it provides an outer approximation to the set of quantum channels that can be implemented with local operations and classical communication (LOCC) on a bipartite system AB . The usefulness of the PPT approximation rests on the key observation that condition (S9) amounts to a positive semi-definite constraint, as can be seen by employing (S4).

C. Zero-error PPT entanglement cost

The **zero-error PPT entanglement cost** of a bipartite state $\rho = \rho_{AB}$ is defined as the minimum rate of **ebits** (or singlets)

$$\Phi_2 := |\Phi_2\rangle\langle\Phi_2|, \quad |\Phi_2\rangle := \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle), \quad (\text{S10})$$

that need to be consumed in order to create copies of ρ with PPT operations and zero error. In formula,

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) := \inf \left\{ R : \text{for all sufficiently large } n \in \mathbb{N} \exists \Lambda_n \in \text{PPT} \cap \text{CPTP} : \Lambda_n(\Phi_2^{\otimes \lfloor Rn \rfloor}) = \rho^{\otimes n} \right\}. \quad (\text{S11})$$

Wang and Wilde proved [25, 37] that $E_{c,\text{PPT}}^{\text{exact}}$ can be alternatively computed as

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\kappa}^{\infty}(\rho) := \lim_{n \rightarrow \infty} \frac{1}{n} E_{\kappa}(\rho^{\otimes n}), \quad (\text{S12})$$

$$E_{\kappa}(\rho) := \log_2 \min \{ \text{Tr } S : -S \leq \rho^{\Gamma} \leq S, S^{\Gamma} \geq 0 \}. \quad (\text{S13})$$

he first equality in (S12) is an asymptotic consequence of the more general one-shot statement that the PPT entanglement cost of generating a single copy of a state ρ with PPT operations, denoted by $E_{c,\text{PPT}}^{(1)}(\rho)$, can be bounded as $\log_2(2^{E_{\kappa}(\rho)} - 1) \leq E_{c,\text{PPT}}^{(1)}(\rho) \leq \log_2(2^{E_{\kappa}(\rho)} + 2)$ [25, Proposition 2]. It is easy to see that the function E_{κ} is sub-additive, so that by Fekete’s lemma [56] we can alternatively write the limit in (S12) as an infimum over n , i.e.

$$E_{\kappa}^{\infty}(\rho) = \inf_{n \in \mathbb{N}^+} \frac{1}{n} E_{\kappa}(\rho^{\otimes n}). \quad (\text{S14})$$

Unfortunately, due to an error in the derivation of the super-additivity inequality, which can be traced back to [25, Eq. (S59)], E_{κ} is no longer known to be additive. Hence, the regularisation in the above identity is needed. And in fact, we will show in the forthcoming Section II that it *must* be included, because E_{κ} is indeed *not* additive in general.

We refer the interested reader to the erratum [37] for a detailed explanation of this issue and of its consequences. There, Wang and Wilde also present other examples of additivity violations for E_{κ} . In the next section, we will construct our own examples of additivity violation.

II. VIOLATIONS OF ADDITIVITY OF E_{κ}

To construct an example of additivity violation, it is useful to start by asking the opposite question: when is it that we know $E_{\kappa}(\rho)$ to be additive? There is a simple answer to this question that is already discussed in [25] based on the pioneering work [24]: E_{κ} is easily seen to be additive on many copies of ρ whenever ρ has zero bi-negativity, i.e. whenever $|\rho^{\Gamma}|^{\Gamma} \geq 0$. When this happens, then

$$E_{\kappa}(\rho^{\otimes n}) = \log_2 \left\| (\rho^{\otimes n})^{\Gamma} \right\|_1 = n \log_2 \left\| \rho^{\Gamma} \right\|_1 = n E_{\kappa}(\rho) = n E_{\kappa}(\rho), \quad (\text{S15})$$

for all n . This observation tells us that to look for additivity violations of E_{κ} we need to construct states with non-zero bi-negativity. After a bit of trial and error, we homed in on the following construction.

Definition S2 (Punch card states). *Let $A \geq 0$ be a $d \times d$ positive semi-definite matrix. Also, let Q be a $d \times d$ symmetric matrix with $Q_{ij} \in \{0, 1\}$ for all $i, j = 1, \dots, d$. Then the associated punch card state is a bipartite quantum state acting on $\mathbb{C}^d \otimes \mathbb{C}^d$ and defined by*

$$\pi_{A,Q} \propto \sum_{i,j} A_{ij} |ii\rangle\langle jj| + \sum_{i \neq j} Q_{ij} |A_{ij}\rangle\langle ij|, \quad (\text{S16})$$

with the normalisation $\text{Tr } \pi_{A,Q} = 1$.

Note that

$$\begin{aligned} \pi_{A,Q}^\Gamma &\propto \sum_i A_{ii} |ii\rangle\langle ii| + \sum_{i<j} \left(Q_{ij} |A_{ij}| (|ij\rangle\langle ij| + |ji\rangle\langle ji|) + A_{ij} |ij\rangle\langle ji| + A_{ij}^* |ji\rangle\langle ij| \right) \\ &\simeq \left(\sum_i A_{ii} |ii\rangle\langle ii| \right) \oplus \bigoplus_{i<j} \begin{pmatrix} Q_{ij}|A_{ij}| & A_{ij} \\ A_{ij}^* & Q_{ij}|A_{ij}| \end{pmatrix}, \end{aligned} \quad (\text{S17})$$

from which, using the elementary observation that

$$\left| \begin{pmatrix} Q_{ij}|A_{ij}| & A_{ij} \\ A_{ij}^* & Q_{ij}|A_{ij}| \end{pmatrix} \right| = \begin{pmatrix} |A_{ij}| & Q_{ij}A_{ij} \\ Q_{ij}A_{ij}^* & |A_{ij}| \end{pmatrix}, \quad (\text{S18})$$

valid because $Q_{ij} = 0, 1$, we obtain that

$$\left| \pi_{A,Q}^\Gamma \right|^\Gamma \propto \sum_{i,j} Q_{ij} A_{ij} |ii\rangle\langle jj| + \sum_{i \neq j} |A_{ij}| |ij\rangle\langle ij|, \quad (\text{S19})$$

where \propto hides a positive constant. We therefore record the following observation:

Lemma S3. *Let $A \geq 0$ be a $d \times d$ positive semi-definite matrix, and let Q be a $d \times d$ symmetric matrix with $Q_{ij} \in \{0, 1\}$ for all $i, j = 1, \dots, d$. If $Q \circ A \not\geq 0$, where \circ denotes the Hadamard product, then the punch card state $\pi_{A,Q}$ does not have zero bi-negativity, meaning that*

$$\left| \pi_{A,Q}^\Gamma \right|^\Gamma \not\geq 0. \quad (\text{S20})$$

In particular, the state

$$\pi_0 := \pi_{A_0, Q_0}, \quad A_0 := \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad Q_0 := \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \not\geq 0 \quad (\text{S21})$$

does not have zero bi-negativity.

And sure enough, the above punch card state is already enough to exhibit a small but meaningful violation of additivity for E_κ , already at the 2-copy level:

Proposition S4. *For the state π_0 defined by (S21), it holds that*

$$2E_{c, \text{PPT}}^{\text{exact}}(\pi_0) \leq E_\kappa(\pi_0^{\otimes 2}) < 2E_\kappa(\pi_0). \quad (\text{S22})$$

In particular, $E_\kappa(\pi_0^{\otimes 2}) \approx 1.001$ and $2E_\kappa(\pi_0) \approx 1.029$.

Proof. The first inequality follows by combining (S12) and (S14). The numerical calculations of $E_\kappa(\pi_0^{\otimes 2})$ and $E_\kappa(\pi_0)$ can be verified with an SDP solver. \square

III. NEW ENTANGLEMENT MONOTONES: $E_{\chi,p}$ AND $E_{\kappa,q}$

A. Rationale of our construction

Let us investigate in greater depth the fundamental reason why E_κ fails to be additive. To this end, it is useful to look at the dual SDP program [35, 36] for E_κ , which takes the form [25, Eq. (S25)]

$$E_\kappa(\rho) = \log_2 \max \left\{ \text{Tr } \rho (V - W)^\Gamma : V, W \geq 0, V^\Gamma + W^\Gamma \leq \mathbb{1} \right\}. \quad (\text{S23})$$

Since E_κ is sub-additive, to prove additivity we would only need to prove super-additivity. This is most naturally done with the dual program, which involves a maximisation. The proof of $E_\kappa(\rho \otimes \rho') \stackrel{?}{\geq} E_\kappa(\rho) + E_\kappa(\rho')$ would proceed as follows: first we consider optimal feasible points V, W and V', W' that achieve the maxima in the dual SDPs (S23) for ρ and ρ' ; then we attempt to construct a feasible point V'', W'' for the dual SDP for $\rho \otimes \rho'$; such an ansatz would need to yield a value of the objective function equal to $E_\kappa(\rho) + E_\kappa(\rho')$ (upon taking the logarithm). A quick inspection reveals that this requires that

$$\text{Tr } \rho (V'' - W'')^\Gamma \stackrel{?}{=} \left(\text{Tr } \rho (V - W)^\Gamma \right) \left(\text{Tr } \rho (V' - W')^\Gamma \right). \quad (\text{S24})$$

A natural ansatz at this point is the one found by Wang and Wilde [25], namely,

$$V'' = V \otimes V' + W \otimes W', \quad W'' = V \otimes W' + W \otimes V'. \quad (\text{S25})$$

Note that $V'', W'' \geq 0$; furthermore, this ansatz clearly satisfies (S24). The problem is that

$$(V'' + W'')^\Gamma = (V + W)^\Gamma \otimes (V' + W')^\Gamma \not\leq \mathbb{1} \quad (\text{S26})$$

in general. Indeed, while $(V + W)^\Gamma \leq \mathbb{1}$ and $(V' + W')^\Gamma \leq \mathbb{1}$ holds by hypothesis, from this it does not follow that the tensor product of these operators is also upper bounded by the identity. The reason is that both of these operators could have large negative eigenvalues, which would become large and positive when multiplied.

The first idea we have is to modify the definition of E_κ so as to make sure that this does not happen. To this end, it suffices to add the further constraint $(V + W)^\Gamma \geq -\mathbb{1}$. We can therefore define

$$\begin{aligned} E_\chi(\rho) &:= \log_2 \max \left\{ \text{Tr } \rho (V - W)^\Gamma : V, W \geq 0, -\mathbb{1} \leq V^\Gamma + W^\Gamma \leq \mathbb{1} \right\} \\ &= \log_2 \min \left\{ \text{Tr } T : -S \leq \rho^\Gamma \leq S, -T \leq S^\Gamma \leq T \right\} \\ &= \log_2 \min \left\{ \|S^\Gamma\|_1 : -S \leq \rho^\Gamma \leq S \right\}, \end{aligned} \quad (\text{S27})$$

where in the second line we derived the primal program corresponding to the dual program in the first line, and in the third line we used Lemma S1.

We will see shortly that our approach fixes the additivity problem, in the sense that the new function E_χ we have just constructed is indeed additive, unlike E_κ . Of course, what we are really interested in is the regularisation of E_κ ; in general, E_χ can only provide a lower bound to that quantity. Indeed, for all states it holds that $E_\chi \leq E_\kappa$, because taking the optimisation on the third line of (S27) and restricting to operators S that satisfy $S^\Gamma \geq 0$ yields precisely E_κ , as can be seen by observing that in those cases $\|S^\Gamma\|_1 = \text{Tr } S$. Assuming the additivity of E_χ , which we will prove shortly, we immediately deduce that

$$E_\kappa \geq E_\kappa^\infty \geq E_\chi. \quad (\text{S28})$$

While this is already interesting in itself, we can do better. In fact, the definition of E_χ in the second line of (S27) lends itself naturally to some generalisations.

Definition S5 (χ -hierarchy). Let $\rho = \rho_{AB}$ be any bipartite quantum state. For $p \in \mathbb{N}$, we define the corresponding χ -quantity $E_{\chi,p}$ as

$$E_{\chi,p}(\rho) := \log_2 \chi_p(\rho), \quad (\text{S29})$$

$$\chi_p(\rho) := \min \left\{ \text{Tr } S_p : -S_i \leq S_{i-1}^\Gamma \leq S_i, i = 0, \dots, p, S_{-1} = \rho \right\} \quad (\text{S30})$$

$$= \min \left\{ \text{Tr } S_p : -S_0 \leq \rho^\Gamma \leq S_0, -S_1 \leq S_0^\Gamma \leq S_1, \dots, -S_p \leq S_{p-1}^\Gamma \leq S_p \right\}. \quad (\text{S31})$$

We will refer to the sequence of functions $E_{\chi,p}$, $p \in \mathbb{N}$, as the χ -hierarchy.

Remark S6. The minima in (S30)–(S31) are always achieved, essentially because of the compactness of the set of positive semi-definite operators with trace bounded by a fixed number.

Remark S7. For $p > 0$, due to Lemma S1 we can equivalently write

$$\chi_p(\rho) = \min \left\{ \|S_{p-1}^\Gamma\| : -S_0 \leq \rho^\Gamma \leq S_0, -S_1 \leq S_0^\Gamma \leq S_1, \dots, -S_{p-1} \leq S_{p-2}^\Gamma \leq S_{p-1} \right\}. \quad (\text{S32})$$

When $p = 0$, clearly we have that

$$E_{\chi,0}(\rho) = \log_2 \|\rho^\Gamma\|_1 = E_N(\rho) \quad (\text{S33})$$

yields the celebrated entanglement negativity [32, 33]. For $p = 1$, instead,

$$E_{\chi,1}(\rho) = E_\chi(\rho) \quad (\text{S34})$$

reproduces (S27).

While this is quite interesting, we can now observe that the above functions $E_{\chi,p}$ are of a somewhat different nature than the E_κ monotone introduced by Wang and Wilde [25], whose definition is reported in (S13). We can however try to generalise that construction as well. This is done as follows: for $q \geq 1$, one replaces p with q in (S30), restricts the minimisation there to operators S_{q-1} satisfying $S_{q-1}^\Gamma \geq 0$, and sets $S_q = S_{q-1}^\Gamma$ accordingly; this gives a ‘ κ -hierarchy’ defined as follows.

Definition S8 (κ -hierarchy). *Let $\rho = \rho_{AB}$ be any bipartite quantum state. For an arbitrary positive integer $q \in \mathbb{N}^+$, we define the corresponding κ -quantity $E_{\kappa,q}$ as*

$$E_{\kappa,q}(\rho) := \log_2 \kappa_q(\rho), \quad (\text{S35})$$

$$\kappa_q(\rho) := \min \left\{ \text{Tr } S_{q-1} : -S_i \leq S_{i-1}^\Gamma \leq S_i, \ i = 0, \dots, q-1, \ S_{-1} = \rho, \ S_{q-1}^\Gamma \geq 0 \right\} \quad (\text{S36})$$

$$= \min \left\{ \text{Tr } S_{q-1} : -S_0 \leq \rho^\Gamma \leq S_0, \ \dots, \ -S_{q-1} \leq S_{q-2}^\Gamma \leq S_{q-1}, \ S_{q-1}^\Gamma \geq 0 \right\}. \quad (\text{S37})$$

We will refer to the sequence of functions $E_{\chi,q}$, $q \in \mathbb{N}^+$, as the κ -hierarchy.

For reasons analogos to those explained in Remark S6, the minima in (S36)–(S37) are always achieved. Furthermore, note that

$$E_{\kappa,1}(\rho) = E_\kappa(\rho) \quad (\text{S38})$$

reproduces the function in (S13). We shall now explore the properties of the χ - and κ -hierarchies in detail.

B. Relation between the χ - and κ -hierarchies

We now begin our investigation of the properties of the χ - and κ -hierarchies. First, we will study the relation between the two. A very simple yet very useful observation that we record as a preliminary lemma is as follows.

Lemma S9 (Increasing trace). *Let the operators S_0, \dots, S_p be a feasible point for the SDP (S30) that defines $\chi_p(\rho)$, for some bipartite state $\rho = \rho_{AB}$ and some $p \in \mathbb{N}$. Then $S_0, \dots, S_p \geq 0$ are positive semi-definite, and moreover*

$$1 \leq \text{Tr } S_0 \leq \text{Tr } S_1 \leq \dots \leq \text{Tr } S_p. \quad (\text{S39})$$

Similarly, if S'_0, \dots, S'_{q-1} constitute a feasible point for the SDP (S36) that defines $\kappa_q(\rho)$ ($q \in \mathbb{N}^+$), then $S'_0, \dots, S'_{q-1} \geq 0$, and

$$1 \leq \text{Tr } S'_0 \leq \text{Tr } S'_1 \leq \dots \leq \text{Tr } S'_{q-1}. \quad (\text{S40})$$

Proof. From the inequalities $-S_i \leq S_{i-1}^\Gamma \leq S_i$ ($i = 0, \dots, p$, with $S_{-1} = \rho$) it follows immediately that $-S_i \leq S_i$, i.e. $S_i \geq 0$. To prove (S39), it suffices to take the trace of the inequalities $S_0^\Gamma \leq S_1$, $S_1^\Gamma \leq S_2$, \dots , and $S_{p-1}^\Gamma \leq S_p$. The proof of $S'_0, \dots, S'_{q-1} \geq 0$ and of (S40) is entirely analogous. \square

Remark S10. In particular, for all $p \in \mathbb{N}$, $q \in \mathbb{N}^+$, and bipartite states ρ , it holds that $\chi_p(\rho) \geq 1$ and $\kappa_q(\rho) \geq 1$.

We are now ready to prove the following result.

Proposition S11. *For all non-negative integers $p \in \mathbb{N}$, positive integers $q \in \mathbb{N}^+$, and for all bipartite states $\rho = \rho_{AB}$, it holds that*

$$E_N = E_{\chi,0} \leq \dots \leq E_{\chi,p} \leq E_{\chi,p+1} \leq \dots \leq E_{\kappa,q+1} \leq E_{\kappa,q} \leq \dots \leq E_\kappa, \quad (\text{S41})$$

where we forwent the dependence on ρ . In other words, the function $\mathbb{N} \ni p \mapsto E_{\chi,p}(\rho)$ is increasing¹, while $\mathbb{N}^+ \ni q \mapsto E_{\kappa,q}(\rho)$ is decreasing; moreover, $E_{\chi,p}(\rho) \leq E_{\kappa,q}(\rho)$ holds for all $p \in \mathbb{N}$ and $q \in \mathbb{N}^+$.

Proof. We prove the claims one by one.

- $\mathbb{N} \ni p \mapsto E_{\chi,p}$ is increasing (for fixed ρ). Let S_0, \dots, S_{p+1} be optimal feasible points for the SDP that defines χ_{p+1} . Then the operators S_0, \dots, S_p clearly constitute a feasible point for the SDP that defines χ_p , because all the inequalities that have to be obeyed are also present in the SDP that defines χ_{p+1} . Hence, $\chi_p \leq \text{Tr } S_p \leq \text{Tr } S_{p+1} = \chi_{p+1}$, where for the second inequality we leveraged Lemma S9 (and, more precisely, Eq. (S39)). Taking the logarithm completes the argument.
- $\mathbb{N}^+ \ni q \mapsto E_{\kappa,q}$ is decreasing (for fixed ρ). The proof is somewhat similar, but reversed. If S_0, \dots, S_{q-1} represent the optimal feasible point for the SDP (S36) that defines κ_q , then we can construct a corresponding feasible point $S_0, \dots, S_{q-1}, S_q := S_{q-1}^\Gamma$ for the SDP that defines κ_{q+1} . Indeed, the inequalities $-S_i \leq S_{i-1}^\Gamma \leq S_i$ are automatically satisfied for $i = 0, \dots, q-1$ (here, $S_{-1} = \rho$). For $i = q$, we obviously have that $-S_q = -S_{q-1}^\Gamma \leq S_{q-1}^\Gamma = S_q$ due to the fact that $S_{q-1}^\Gamma \geq 0$. Finally, the last constraint in the SDP for κ_{q+1} is $S_q^\Gamma \geq 0$. This is also obeyed, because $S_q^\Gamma = S_{q-1}$, which is positive definite due to Lemma S9. Using (S7) we thus obtain that $\kappa_{q+1} \leq \text{Tr } S_q = \text{Tr } S_{q-1}^\Gamma = \text{Tr } S_{q-1} = \kappa_q$. Taking the logarithm completes the proof of the claim.
- $E_{\chi,p} \leq E_{\kappa,q}$ for all $p \in \mathbb{N}$ and $q \in \mathbb{N}^+$. Due to the fact that $\mathbb{N} \ni p \mapsto E_{\chi,p}$ is increasing and $\mathbb{N} \ni q \mapsto E_{\kappa,q}$ is decreasing, the claim is equivalent to the statement that

$$\sup_{p \in \mathbb{N}} E_{\chi,p} = \lim_{p \rightarrow \infty} E_{\chi,p} \stackrel{?}{\leq} \lim_{q \rightarrow \infty} E_{\kappa,q} = \inf_{q \in \mathbb{N}^+} E_{\kappa,q}. \quad (\text{S42})$$

Now, let S_0, \dots, S_{q-1} be optimal feasible points for the SDP (S36) that represents κ_q , for some $q \in \mathbb{N}^+$. By direct inspection of (S30), we see that these are also feasible points for the SDP that represents χ_{q-1} . Therefore,

$$\chi_{q-1} \leq \text{Tr } S_{q-1} = \kappa_q. \quad (\text{S43})$$

Since this holds for an arbitrary $q \in \mathbb{N}^+$, we can now take the limit $q \rightarrow \infty$ and subsequently the logarithm of both sides, thus concluding the proof. □

To conclude this subsection, we observe that both of our hierarchies collapse into one single quantity in the simple case where the underlying state ρ has zero bi-negativity.

Lemma S12. *If $\rho = \rho_{AB}$ has zero bi-negativity, in the sense that $|\rho^\Gamma|^\Gamma \geq 0$, then*

$$E_{\chi,p}(\rho) = E_{\kappa,q}(\rho) = E_N(\rho) = \log_2 \|\rho^\Gamma\|_1 \quad (\text{S44})$$

for all $p \in \mathbb{N}$ and $q \in \mathbb{N}^+$. In particular, if Φ_d is the maximally entangled state defined by (S3), then

$$E_{\chi,p}(\Phi_d) = E_{\kappa,q}(\Phi_d) = \log_2 d. \quad (\text{S45})$$

Proof. Wang and Wilde [25, Proposition 8] have already established that $E_\kappa(\rho) = E_N(\rho)$ if ρ has zero bi-negativity. Due to Proposition S11, this implies a complete collapse of both hierarchies. As $\Phi_d^\Gamma = \frac{1}{d}F$ where F is the swap operator, it is elementary to observe that $|\Phi_d^\Gamma| = \frac{1}{d}\mathbb{1}$ and so the maximally entangled state Φ_d indeed has zero bi-negativity; moreover, one immediately recovers the well-known result that $E_N(\Phi_d) = \log_2 \|\Phi_d^\Gamma\|_1 = \log_2 d$ [32]. □

¹ Not necessarily *strictly* increasing. The meaning of ‘decreasing’ should similarly be intended as ‘non-increasing’.

C. $E_{\chi,p}$ and $E_{\kappa,q}$ as PPT monotones

We will now prove that our functions χ_p, κ_q , as well as their logarithmic versions $E_{\chi,p}$ and $E_{\kappa,q}$, represent novel entanglement monotones under PPT operations.

Proposition S13 (Strong monotonicity). *For two bipartite quantum systems AB and $A'B'$, let $(\mathcal{N}_x)_x$ be a collection of PPT maps with input system AB and output system $A'B'$ such that $\sum_x \mathcal{N}_x$ is trace preserving. Then for any bipartite quantum state ρ on AB it holds that*

$$F(\rho) \geq \sum_{x: \text{Tr}[\mathcal{N}_x(\rho)] > 0} \text{Tr}[\mathcal{N}_x(\rho)] F\left(\frac{\mathcal{N}_x(\rho)}{\text{Tr}[\mathcal{N}_x(\rho)]}\right), \quad F = \chi_p, \kappa_q, E_{\chi,p}, E_{\kappa,q}, \quad \forall p \in \mathbb{N}, \forall q \in \mathbb{N}^+. \quad (\text{S46})$$

In particular, for any PPT channel \mathcal{N} it holds that

$$\begin{aligned} E_{\chi,p}(\rho) &\geq E_{\chi,p}(\mathcal{N}(\rho)), \\ E_{\kappa,q}(\rho) &\geq E_{\kappa,q}(\mathcal{N}(\rho)). \end{aligned} \quad (\text{S47})$$

Proof. The cases $F = E_{\chi,p}$ and $F = E_{\kappa,q}$ of (S46) follow by taking the logarithm of the same inequality written for the case $F = \chi_p$ and $F = \kappa_q$ while exploiting the concavity of the function $x \mapsto \log_2(x)$.

We thus focus on the cases $F = \chi_p, \kappa_q$, starting from the former. Let S_{-1}, S_0, \dots, S_p be optimisers of the minimisation problem (S30) that defines $\chi_p(\rho)$. In particular,

$$S_{-1} = \rho, \quad -S_i \leq S_{i-1}^\Gamma \leq S_i \quad \forall i = 0, 1, \dots, p. \quad (\text{S48})$$

Since \mathcal{N}_x is PPT, the map $\Gamma \circ \mathcal{N}_x \circ \Gamma$ is (completely) positive. Eq. (S48) then implies that

$$-[\mathcal{N}_x(S_i^\Gamma)]^\Gamma \leq [\mathcal{N}_x(S_{i-1})]^\Gamma \leq [\mathcal{N}_x(S_i^\Gamma)]^\Gamma \quad \forall i = 0, 1, \dots, p. \quad (\text{S49})$$

Moreover, Eq. (S48) and the fact that \mathcal{N}_x is completely positive imply that

$$-\mathcal{N}_x(S_i) \leq \mathcal{N}_x(S_{i-1}^\Gamma) \leq \mathcal{N}_x(S_i) \quad \forall i = 0, 1, \dots, p-1. \quad (\text{S50})$$

Consequently, for any $i = -1, 0, \dots, p$ we can introduce the operators

$$S_i^{(x)} := \begin{cases} [\mathcal{N}_x(S_i^\Gamma)]^\Gamma & \text{if } i \text{ is even,} \\ \mathcal{N}_x(S_i) & \text{if } i \text{ is odd} \end{cases} \quad (\text{S51})$$

(here, -1 is considered odd), in terms of which we can rewrite (S49) as

$$-S_i^{(x)} \leq (S_{i-1}^{(x)})^\Gamma \leq S_i^{(x)} \quad (i \text{ even}), \quad (\text{S52})$$

and (S50) as

$$-S_i^{(x)} \leq (S_{i-1}^{(x)})^\Gamma \leq S_i^{(x)} \quad (i \text{ odd}). \quad (\text{S53})$$

It follows that for any x such that $\text{Tr}[\mathcal{N}_x(\rho)] > 0$ the operators $S_i^{(x)} / \text{Tr}[\mathcal{N}_x(\rho)]$, $i = -1, 0, \dots, p$ are valid ansatzes in the minimisation problem (S30) that defines $\chi_p\left(\frac{\mathcal{N}_x(\rho)}{\text{Tr}[\mathcal{N}_x(\rho)]}\right)$. Therefore,

$$\begin{aligned} \chi_p(\rho) &= \text{Tr}[S_p] \\ &\stackrel{(i)}{=} \sum_x \text{Tr}[S_p^{(x)}] \\ &\stackrel{(ii)}{\geq} \sum_{x: \text{Tr}[\mathcal{N}_x(\rho)] > 0} \text{Tr}[\mathcal{N}_x(\rho)] \text{Tr}\left[\frac{S_p^{(x)}}{\text{Tr}[\mathcal{N}_x(\rho)]}\right] \\ &\stackrel{(iii)}{\geq} \sum_{x: \text{Tr}[\mathcal{N}_x(\rho)] > 0} \text{Tr}[\mathcal{N}_x(\rho)] \chi_p\left(\frac{\mathcal{N}_x(\rho)}{\text{Tr}[\mathcal{N}_x(\rho)]}\right). \end{aligned} \quad (\text{S54})$$

Here, in (i) we used the fact that the maps $\sum_x \mathcal{N}_x$ and $\sum_x \Gamma \circ \mathcal{N}_x \circ \Gamma$ are trace preserving; in (ii) we observed that $S_p^{(x)} \geq 0$, which follows from (S49) and (S50); finally, in (iii) we leveraged the fact that the operators $S_i^{(x)} / \text{Tr} \mathcal{N}_x(\rho)$, $i = -1, 0, \dots, p$ are valid ansatzes in the minimisation problem (S30) that defines $\chi_p \left(\frac{\mathcal{N}_x(\rho)}{\text{Tr}[\mathcal{N}_x(\rho)]} \right)$, as argued above. This proves (S46) for $F = \chi_p$.

The proof of the strong monotonicity property (S46) for the case where $F = \kappa_q$ is completely analogous. The only additional observation we need is that $(S_{q-1}^{(x)})^\Gamma \geq 0$ if $S_{q-1}^\Gamma \geq 0$. Indeed, if q is odd then $(S_{q-1}^{(x)})^\Gamma = \mathcal{N}_x(S_{q-1}^\Gamma)$, which is positive because \mathcal{N}_x is a positive map; if q is even then $(S_{q-1}^{(x)})^\Gamma = \mathcal{N}_x(S_{q-1}^\Gamma)^\Gamma = (\Gamma \circ \mathcal{N}_x \circ \Gamma)(S_{q-1}^\Gamma)$, which is positive definite because $\Gamma \circ \mathcal{N}_x \circ \Gamma$ is a positive map. This concludes the proof. \square

As an immediate consequence of Lemma S12 and Proposition S13, we deduce the following.

Corollary S14. *Let $\rho = \rho_{AB}$ be an arbitrary quantum state of a bipartite system AB with minimal local dimension $d := \min\{|A|, |B|\}$. Then for all $p \in \mathbb{N}$ and $q \in \mathbb{N}^+$ it holds that*

$$\chi_p(\rho), \kappa_q(\rho) \leq d, \quad E_{\chi,p}(\rho), E_{\kappa,q}(\rho) \leq \log_2 d. \quad (\text{S55})$$

Proof. An application of Nielsen's theorem [57] guarantees that there exists an LOCC transformation Λ such that $\Lambda(\Phi_d) = \rho_{AB}$, where Φ_d is the maximally entangled state with local dimension d . Since LOCC is a subset of PPT, the result then follows by combining Lemma S12 and Proposition S13. Incidentally, an alternative, direct way of constructing an LOCC with the above property is to let the party with the maximum local dimension prepare ρ_{AB} locally and then apply the quantum teleportation protocol [58] to teleport the smaller system to the other party. \square

D. Dual expressions

In what follows, we derive the SDPs dual to those in (S30) and (S36). We refer the reader to [34–36] for an introduction to the general theory of semi-definite programs. We start with the χ -quantities, whose dual expressions will play an important role in establishing their additivity.

Proposition S15 (Dual SDP for the χ -quantity). *For an arbitrary bipartite state $\rho = \rho_{AB}$ and some $p \in \mathbb{N}$, the associated χ -quantity $\chi_p(\rho)$ defined by (S30) can be equivalently expressed as*

$$\begin{aligned} \chi_p(\rho) = \max. \quad & \text{Tr} [\rho (V_0^\Gamma - W_0^\Gamma)] \\ & V_0, \dots, V_p, W_0, \dots, W_p \geq 0, \\ \text{s.t.} \quad & V_i + W_i = V_{i+1}^\Gamma - W_{i+1}^\Gamma \quad \forall i = 0, \dots, p-1, \quad V_p + W_p = \mathbb{1}, \end{aligned} \quad (\text{S56})$$

or else, if $p \geq 1$, as

$$\begin{aligned} \chi_p(\rho) = \max. \quad & \text{Tr} [\rho (V_0^\Gamma - W_0^\Gamma)] \\ & V_0, \dots, V_{p-1}, W_0, \dots, W_{p-1} \geq 0, \\ \text{s.t.} \quad & V_i + W_i = V_{i+1}^\Gamma - W_{i+1}^\Gamma \quad \forall i = 0, \dots, p-2, \quad -\mathbb{1} \leq (V_{p-1} + W_{p-1})^\Gamma \leq \mathbb{1}. \end{aligned} \quad (\text{S57})$$

Remark S16. If $p = 0$, then the only constraints in (S56) are $V_0, W_0 \geq 0$ and $V_0 + W_0 = \mathbb{1}$. Similarly, if $p = 1$ then the only constraints in (S57) are $V_0, W_0 \geq 0$ and $-\mathbb{1} \leq (V_0 + W_0)^\Gamma \leq \mathbb{1}$. Note that for $p = 1$ the SDP in (S57) reproduces (up to the logarithm) the expression on the first line of (S27), as it should.

Proof of Proposition S15. The $p+1$ (effectively free) variables S_0, \dots, S_p in (S30) obey $2(p+1)$ semi-definite constraints, namely $S_i \pm S_{i-1}^\Gamma \geq 0$ for $i = 0, \dots, p$, with the convention that $S_{-1} = \rho$. We therefore introduce $p+1$ positive semi-definite dual variables $V_0, \dots, V_p \geq 0$ for the inequalities with the minus and another $p+1$, namely $W_0, \dots, W_p \geq 0$, for the inequalities with the plus. The resulting Lagrangian is

$$\begin{aligned} \mathcal{L} &= \text{Tr} S_p - \sum_{i=0}^p \text{Tr} V_i (S_i - S_{i-1}^\Gamma) - \sum_{i=0}^p \text{Tr} W_i (S_i + S_{i-1}^\Gamma) \\ &= \text{Tr} \rho (V_0^\Gamma - W_0^\Gamma) - \sum_{i=0}^{p-1} \text{Tr} S_i (V_i + W_i - V_{i+1}^\Gamma + W_{i+1}^\Gamma) + \text{Tr} S_p (\mathbb{1} - V_p - W_p), \end{aligned} \quad (\text{S58})$$

where in the second line we employed (S6) repeatedly. Minimising the above expression over S_0, \dots, S_p yields $-\infty$ unless $V_i + W_i - V_{i+1}^\Gamma + W_{i+1}^\Gamma = 0$ for all $i = 0, \dots, p-1$, and moreover $V_p + W_p = \mathbb{1}$. This proves that the SDP dual to that in (S30) is the one given by (S56).

To see that strong duality holds (and the dual is achieved), it suffices to check that Slater's condition is obeyed, and that both the primal and the dual program are in fact strictly feasible. This simply means that all inequalities can be satisfied strictly. In the primal program (S30), one can take $S_i = (i+1)\mathbb{1}$ ($i = 0, \dots, p$); in the dual program (S56), instead, it suffices to set $V_i = 2 \cdot 3^{-(p-i+1)}\mathbb{1}$, $W_i = 3^{-(p-i+1)}\mathbb{1}$ ($i = 0, \dots, p$).

Finally, it remains to see that (S57) is equivalent to (S56) if $p \geq 1$. This can be done as follows. In (S56), the only constraints on $V_p, W_p \geq 0$ are that $V_p - W_p = (V_{p-1} - W_{p-1})^\Gamma$ and $V_p + W_p = \mathbb{1}$. It is possible to satisfy these constraints if and only if $-\mathbb{1} \leq (V_{p-1} - W_{p-1})^\Gamma \leq \mathbb{1}$. Indeed, this condition is necessary, because on the one hand $(V_{p-1} - W_{p-1})^\Gamma \leq V_p \leq V_p + W_p = \mathbb{1}$, and on the other $(V_{p-1} - W_{p-1})^\Gamma \geq -W_p \geq -(V_p + W_p) = -\mathbb{1}$. It is also sufficient, since if it is obeyed we can set $V_p = \frac{1}{2} \left(\mathbb{1} + (V_{p-1} - W_{p-1})^\Gamma \right)$ and $W_p = \frac{1}{2} \left(\mathbb{1} - (V_{p-1} - W_{p-1})^\Gamma \right)$. \square

Although it is not strictly needed in what follows, for completeness we derive also the dual program for the κ -quantities.

Proposition S17 (Dual SDP for the κ -quantity). *For an arbitrary bipartite state $\rho = \rho_{AB}$ and some $q \in \mathbb{N}^+$, the κ -quantity $\kappa_q(\rho)$ defined by (S36) can be equivalently expressed as*

$$\begin{aligned} \kappa_q(\rho) = \max. \quad & \text{Tr} \left[\rho (V_0^\Gamma - W_0^\Gamma) \right] \\ & V_0, \dots, V_{q-1}, W_0, \dots, W_{q-1} \geq 0, \\ \text{s.t.} \quad & V_i + W_i = V_{i+1}^\Gamma - W_{i+1}^\Gamma \quad \forall i = 0, \dots, q-2, \quad (V_{q-1} + W_{q-1})^\Gamma \leq \mathbb{1}. \end{aligned} \quad (\text{S59})$$

Remark S18. If $q = 1$, then the only constraints in (S59) are $V_0, W_0 \geq 0$ and $(V_0 + W_0)^\Gamma \leq \mathbb{1}$. Note that for $q = 1$ the SDP in (S59) reproduces (up to the logarithm) the expression in (S23).

Proof of Proposition S17. The argument is very similar to that employed to prove Proposition S15. This time around we have $2q + 1$ constraints, namely $S_i \pm S_{i-1}^\Gamma \geq 0$ for $i = 0, \dots, q-1$ (with the convention that $S_{-1} = \rho$), as well as $S_{q-1}^\Gamma \geq 0$. We thus introduce q positive semi-definite dual variables $V_0, \dots, V_{q-1} \geq 0$ for the first inequalities with the minus, another q , namely $W_0, \dots, W_{q-1} \geq 0$, for the first inequalities with the plus, and finally one more variable $Z \geq 0$ for the last inequality. The resulting Lagrangian is

$$\begin{aligned} \mathcal{L} &= \text{Tr} S_{q-1} - \sum_{i=0}^{q-1} \text{Tr} V_i (S_i - S_{i-1}^\Gamma) - \sum_{i=0}^{q-1} \text{Tr} W_i (S_i + S_{i-1}^\Gamma) - \text{Tr} Z S_{q-1}^\Gamma \\ &= \text{Tr} \rho (V_0^\Gamma - W_0^\Gamma) - \sum_{i=0}^{q-2} \text{Tr} S_i (V_i + W_i - V_{i+1}^\Gamma + W_{i+1}^\Gamma) + \text{Tr} S_{q-1} (\mathbb{1} - Z^\Gamma - V_{q-1} - W_{q-1}). \end{aligned} \quad (\text{S60})$$

Minimising the above expression over S_0, \dots, S_{q-1} gives $-\infty$ unless $V_i + W_i - V_{i+1}^\Gamma + W_{i+1}^\Gamma = 0$ for all $i = 0, \dots, q-2$, and moreover $V_{q-1} - W_{q-1} = \mathbb{1} - Z^\Gamma$. The maximisation over Z can be eliminated by observing that $V_{q-1} - W_{q-1} = \mathbb{1} - Z^\Gamma$ for some $Z \geq 0$ if and only if $(V_{q-1} - W_{q-1})^\Gamma \leq \mathbb{1}$. This proves that the SDP in (S59) is indeed the dual of that in (S36). To see that they yield the same value, i.e. that the duality gap is zero, it suffices to check Slater's condition. This can be done with a reasoning entirely analogous to that presented in the proof of Proposition S15. \square

E. Additivity properties

We will now look at the additivity properties of our new monotones $E_{\chi,p}$ and $E_{\kappa,q}$ constructed in Definitions S5 and S8, respectively. We will prove that the former are fully additive (Proposition S20) while the latter are only sub-additive (Proposition S21). Before we begin, we need a little preliminary result that is well known in the quantum information literature (see e.g. [59, Lemma 12.35]). We include a self-contained proof for completeness.

Lemma S19. *For $i = 1, 2$, let A_i and B_i be linear operators on the Hilbert space \mathcal{H}_i . Assume that*

$$-A_i \leq B_i \leq A_i \quad \forall i = 1, 2. \quad (\text{S61})$$

Then

$$-A_1 \otimes A_2 \leq B_1 \otimes B_2 \leq A_1 \otimes A_2. \quad (\text{S62})$$

In particular, if $-A \leq B \leq A$ then for all $n \in \mathbb{N}$ it holds that $-A^{\otimes n} \leq B^{\otimes n} \leq A^{\otimes n}$.

Proof. By adding the two operator inequalities

$$\begin{aligned} 0 &\leq (A_1 + B_1) \otimes (A_2 + B_2) = A_1 \otimes A_2 + B_1 \otimes B_2 + A_1 \otimes B_2 + B_1 \otimes A_2, \\ 0 &\leq (A_1 - B_1) \otimes (A_2 - B_2) = A_1 \otimes A_2 + B_1 \otimes B_2 - A_1 \otimes B_2 - B_1 \otimes A_2, \end{aligned} \quad (\text{S63})$$

we infer that $B_1 \otimes B_2 \geq -A_1 \otimes A_2$. Analogously, by adding the two operator inequalities

$$\begin{aligned} 0 &\leq (A_1 - B_1) \otimes (A_2 + B_2) = A_1 \otimes A_2 - B_1 \otimes B_2 + A_1 \otimes B_2 - B_1 \otimes A_2, \\ 0 &\leq (A_1 + B_1) \otimes (A_2 - B_2) = A_1 \otimes A_2 - B_1 \otimes B_2 - A_1 \otimes B_2 + B_1 \otimes A_2, \end{aligned} \quad (\text{S64})$$

one obtains that $B_1 \otimes B_2 \leq A_1 \otimes A_2$. The last claim is applying (S62) iteratively $n - 1$ times, indexed $j = 1, \dots, n - 1$, with $A_1 = A^{\otimes j}$, $B_1 = B^{\otimes j}$, $A_2 = A$, and $B_2 = B$. \square

We are now ready to state and prove the additivity of $E_{\chi,p}$.

Proposition S20 (Additivity of $E_{\chi,p}$). *For all pairs of bipartite states ρ_{AB} , $\omega_{A'B'}$ and for all $p \in \mathbb{N}$, we have that*

$$\chi_p(\rho_{AB} \otimes \omega_{A'B'}) = \chi_p(\rho_{AB}) \chi_p(\omega_{A'B'}), \quad (\text{S65})$$

$$E_{\chi,p}(\rho_{AB} \otimes \omega_{A'B'}) = E_{\chi,p}(\rho_{AB}) + E_{\chi,p}(\omega_{A'B'}). \quad (\text{S66})$$

Proof. Clearly, Eq. (S66) follows from (S65) upon taking the logarithm. We therefore focus on the latter identity. To prove it, we will show that χ_p is both sub-multiplicative and super-multiplicative. Let us start by verifying the former property, i.e. the inequality

$$\chi_p(\rho \otimes \omega) \stackrel{?}{\leq} \chi_p(\rho) \chi_p(\omega), \quad (\text{S67})$$

where we omitted the system labels for simplicity. Let S_0, \dots, S_p be optimisers of the minimisation problem (S30) that defines $\chi_p(\rho)$. Analogously, let S'_0, \dots, S'_p be optimisers of the minimisation problem that defines $\chi_p(\omega)$. By exploiting Lemma S19, one can easily check that $S_0 \otimes S'_0, \dots, S_p \otimes S'_p$ are valid ansatzes in the minimisation problem that defines $\chi_p(\rho \otimes \omega)$. Consequently,

$$\chi_p(\rho \otimes \omega) \leq \text{Tr}[S_p \otimes S'_p] = \chi_p(\rho) \chi_p(\omega). \quad (\text{S68})$$

Now, let us prove the super-multiplicativity of χ_p . We thus need to verify that

$$\chi_p(\rho \otimes \omega) \stackrel{?}{\geq} \chi_p(\rho) \chi_p(\omega). \quad (\text{S69})$$

To this end, we need to employ the dual SDP for χ_p that we derived in Proposition S15. Let $V_0, \dots, V_p, W_0, \dots, W_p \geq 0$ be optimisers of the SDP in (S56) for $\chi_p(\rho)$. Analogously, let $V'_0, \dots, V'_p, W'_0, \dots, W'_p \geq 0$ be optimisers of the SDP as in (S56) but for the state ω . For any $i = 0, \dots, p$, define

$$\begin{aligned} V''_i &:= V_i \otimes V'_i + W_i \otimes W'_i, \\ W''_i &:= V_i \otimes W'_i + W_i \otimes V'_i. \end{aligned} \quad (\text{S70})$$

Note that

$$\begin{aligned} V''_i + W''_i &= (V_i + W_i) \otimes (V'_i + W'_i), \\ V''_i - W''_i &= (V_i - W_i) \otimes (V'_i - W'_i), \end{aligned} \quad (\text{S71})$$

one can easily check that $V''_0, \dots, V''_p, W''_0, \dots, W''_p$ are valid ansatzes for the optimisation as in (S56) but for the state $\rho \otimes \omega$. Indeed, for $i = 0, \dots, p$ we have that $V''_i, W''_i \geq 0$; moreover, if $i \leq p - 1$ it also holds that

$$\begin{aligned} V''_i + W''_i &= (V_i + W_i) \otimes (V'_i + W'_i) \\ &= (V_{i+1} - W_{i+1})^\Gamma \otimes (V'_{i+1} - W'_{i+1})^\Gamma \\ &= ((V_{i+1} - W_{i+1}) \otimes (V'_{i+1} - W'_{i+1}))^\Gamma \\ &= (V''_{i+1} - W''_{i+1})^\Gamma; \end{aligned} \quad (\text{S72})$$

finally, for $i = p$ we see that

$$V_p'' + W_p'' = (V_p + W_p) \otimes (V_p' + W_p') = \mathbb{1} \otimes \mathbb{1} = \mathbb{1}. \quad (\text{S73})$$

Consequently, it holds that

$$\begin{aligned} \chi_p(\rho \otimes \omega) &\geq \text{Tr} \left[(\rho \otimes \omega) (V_0'' - W_0'')^\Gamma \right] \\ &= \text{Tr} \left[\rho (V_0 - W_0)^\Gamma \right] \text{Tr} \left[\omega (V_0' - W_0')^\Gamma \right] \\ &= \chi_p(\rho) \chi_p(\omega). \end{aligned} \quad (\text{S74})$$

This concludes the proof. \square

Unlike the χ -quantities, the corresponding κ -quantities are only sub-additive but in general not fully additive (as we have seen in Section II). Although we will not make use of this property in what follows, we state and prove it for completeness.

Proposition S21 (Sub-additivity of $E_{\kappa,q}$). *For all pairs of bipartite states ρ_{AB} , $\omega_{A'B'}$ and for all $q \in \mathbb{N}^+$, we have that*

$$\kappa_q(\rho_{AB} \otimes \omega_{A'B'}) \leq \kappa_q(\rho_{AB}) \kappa_q(\omega_{A'B'}), \quad (\text{S75})$$

$$E_{\kappa,q}(\rho_{AB} \otimes \omega_{A'B'}) \leq E_{\kappa,q}(\rho_{AB}) + E_{\kappa,q}(\omega_{A'B'}). \quad (\text{S76})$$

Proof. The argument is totally analogous to that in the first part of the proof of the above Proposition S20. Let S_0, \dots, S_{q-1} and S'_0, \dots, S'_{q-1} , respectively, be optimisers of the SDPs that define $\kappa_q(\rho)$ and $\kappa_q(\omega)$ as in (S36). Due once more to Lemma S19, one sees that $S_0 \otimes S'_0, \dots, S_{q-1} \otimes S'_{q-1}$ constitute a feasible point for the SDP problem that defines $\kappa_q(\rho \otimes \omega)$. Consequently,

$$\kappa_q(\rho \otimes \omega) \leq \text{Tr} \left[S_{q-1} \otimes S'_{q-1} \right] = \kappa_q(\rho) \kappa_q(\omega). \quad (\text{S77})$$

This concludes the proof. \square

F. Continuity

To close off this section, we explore the continuity properties of our new monotones. We need a couple of simple lemmas first. To simplify the exposition, it is useful to note that the functions χ_p and κ_q ($p \in \mathbb{N}$, $q \in \mathbb{N}^+$) defined by (S30) and (S36) are perfectly well defined and non-negative not only for states, but also for arbitrary Hermitian operators. In fact, looking at the SDPs in (S30) and (S36), one sees that $\chi_p(X) \geq 0$ and $\kappa_q(X) \geq 0$ must hold for all Hermitian X , because in both cases any feasible point must satisfy $S_i \geq 0$ for all i .

It is very easy to verify that

$$F(X + Y) \leq F(X) + F(Y), \quad F = \chi_p, \kappa_q, \quad p \in \mathbb{N}, q \in \mathbb{N}^+. \quad (\text{S78})$$

and moreover

$$F(-X) = F(X), \quad F = \chi_p, \kappa_q, \quad p \in \mathbb{N}, q \in \mathbb{N}^+. \quad (\text{S79})$$

since X appears with both signs in the definition of both $\chi_p(X)$ and $\kappa_q(X)$. We can now extend Lemma S12 to the non-positive case.

Lemma S22. *For all Hermitian operators $X = X_{AB}$ on a bipartite quantum system of local dimension $d := \min\{|A|, |B|\}$, it holds that*

$$F(X) \leq d \|X\|_1, \quad F = \chi_p, \kappa_q, \quad p \in \mathbb{N}, q \in \mathbb{N}^+. \quad (\text{S80})$$

Proof. Due to the homogeneity of F and to Lemma S12, we have that $F(R) \leq d \text{Tr } R$ for all $R \geq 0$. But then for an arbitrary Hermitian operator X with positive and negative parts X_+ and X_- , respectively,

$$F(X) = F(X_+ - X_-) \leq F(X_+) + F(-X_-) = F(X_+) + F(X_-) \leq d \text{Tr } X_+ + d \text{Tr } X_- = d \|X\|_1, \quad (\text{S81})$$

where we have used first (S78), then (S79), and finally Lemma S12. \square

We are now ready to state the continuity properties of our monotones.

Proposition S23 (Continuity). *Let ρ, ρ' be two quantum states on a bipartite system AB with minimal local dimension $d := \min\{|A|, |B|\}$. Setting $\varepsilon := \frac{1}{2} \|\rho - \rho'\|_1$, for all $p \in \mathbb{N}$ and $q \in \mathbb{N}^+$ we have that*

$$|F(\rho) - F(\rho')| \leq 2d\varepsilon, \quad F = \chi_p, \kappa_q. \quad (\text{S82})$$

Consequently,

$$|f(\rho) - f(\rho')| \leq \log_2(1 + 2d\varepsilon) \leq 2(\log_2 e) d\varepsilon, \quad f = E_{\chi,p}, E_{\kappa,q}. \quad (\text{S83})$$

Proof. Let $X := \rho' - \rho$. Since $\|X\|_1 = 2\varepsilon$, we can write

$$F(\rho') = F(\rho + X) \leq F(\rho) + F(X) \leq F(\rho) + 2d\varepsilon, \quad (\text{S84})$$

where we have used (S78) and Lemma S22. Applying this reasoning with ρ and ρ' exchanged yields also the reverse inequality and thus completes the proof of (S82). As for (S83), it suffices to note that

$$F(\rho') \leq F(\rho) + 2d\varepsilon \leq F(\rho)(1 + 2d\varepsilon), \quad (\text{S85})$$

where in the last line we used that $F(\rho) \geq 1$ by Remark S10. The claim follows by taking logarithms in (S85), and subsequently exchanging the roles of ρ and ρ' . \square

Remark S24. The type of continuity stated in Proposition S23 is substantially weaker than *asymptotic continuity*, a key property of entanglement monotones [60, 61]. The main point is that the Lipschitz constant appearing on the right hand side of (S82)–(S83) is proportional to d instead of $\log_2 d$, as would be needed to have asymptotic continuity. This is not surprising, because the entanglement negativity itself is known to be *not* asymptotically continuous [32], and our monotones can be thought of generalisations of the negativity. It is worth remarking, however, that even the weaker form of continuity established in Proposition S23 can be extremely useful. The same property for the negativity, for example, which can be re-derived by setting $F = \chi_0$ in (S82), is key to proving that it upper bounds the distillable entanglement [32, Section IV]. This observation also underpins the whole approach of [62].

IV. MAIN RESULTS

A. Enter the regularisation

In the previous section we have investigated two families of new PPT monotones, the χ - and the κ -hierarchies. In Proposition S11 we have established a precise hierarchical relation between the two hierarchies, with the increasing functions $E_{\chi,p}(\rho)$ ($p \in \mathbb{N}$) lying below the decreasing functions $E_{\kappa,q}(\rho)$ ($q \in \mathbb{N}^+$). However, our ultimate goal is to understand the zero-error PPT entanglement cost of an arbitrary state ρ , which, by the results of Wang and Wilde (Eq. (S12)), coincides with the regularised E_{κ} , in formula $E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\kappa}^{\infty}(\rho)$. The purpose of this subsection, therefore, is to understand how E_{κ}^{∞} fits in the hierarchy delineated by Proposition S11. A first insight can be deduced by leveraging the results we have obtained in the previous section.

Lemma S25. *Let $\rho = \rho_{AB}$ be an arbitrary bipartite state. Then for all $p \in \mathbb{N}$ it holds that*

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\kappa}^{\infty}(\rho) \geq E_{\chi,p}(\rho). \quad (\text{S86})$$

In particular,

$$E_{\kappa}^{\infty}(\rho) \geq \lim_{p \rightarrow \infty} E_{\chi,p}(\rho) = \sup_{p \in \mathbb{N}} E_{\chi,p}(\rho). \quad (\text{S87})$$

Proof. For all positive integers $n \in \mathbb{N}^+$ we can write

$$\frac{1}{n} E_{\kappa}(\rho^{\otimes n}) \stackrel{(i)}{\geq} \frac{1}{n} E_{\chi,p}(\rho^{\otimes n}) \stackrel{(ii)}{=} E_{\chi,p}(\rho), \quad (\text{S88})$$

where (i) holds due to Proposition S11, and in (ii) we exploited the additivity of $E_{\chi,p}$ (Proposition S20). Taking the limit $n \rightarrow \infty$ and leveraging (S12) concludes the proof of (S86). Upon taking the limit $p \rightarrow \infty$, we obtain also (S87). \square

We have thus established that the function E_κ^∞ lies above the entire χ -hierarchy. We shall now investigate its relationship with the κ -hierarchy. The main result of this subsection is the following.

Theorem S26. *Let $q \in \mathbb{N}^+$ be a positive integer. For any bipartite state $\rho = \rho_{AB}$, it holds that*

$$E_{c, \text{PPT}}^{\text{exact}}(\rho) = E_\kappa^\infty(\rho) \leq E_{\kappa, q}(\rho). \quad (\text{S89})$$

In particular, for every ρ we have that

$$E_N = E_{\chi, 0} \leq E_{\chi, 1} \leq \dots \leq \lim_{p \rightarrow \infty} E_{\chi, p} \leq E_{c, \text{PPT}}^{\text{exact}} = E_\kappa^\infty \leq \lim_{q \rightarrow \infty} E_{\kappa, q} \leq \dots \leq E_{\kappa, 2} \leq E_{\kappa, 1} = E_\kappa. \quad (\text{S90})$$

Proof. Clearly, it suffices to prove (S89), as (S90) would then follow by combining that with the results of Lemma S25 and Proposition S11. In light of these considerations, we thus set out to prove (S89).

Let $S_{-1}, S_0, \dots, S_{q-1}$ be optimisers for the SDP in (S36) for $\kappa_q(\rho)$. Therefore,

$$-S_i \leq S_{i-1}^\Gamma \leq S_i \quad \forall i = 0, \dots, q-1, \quad S_{-1} = \rho, \quad S_{q-1}^\Gamma \geq 0. \quad (\text{S91})$$

For an arbitrary positive integer $n \in \mathbb{N}^+$, Lemma S19 implies that

$$-S_i^{\otimes n} \leq (S_{i-1}^\Gamma)^{\otimes n} \leq S_i^{\otimes n} \quad \forall i = 0, \dots, q-1, \quad (S_{q-1}^\Gamma)^{\otimes n} \geq 0 \quad (\text{S92})$$

Let us define

$$\tilde{S}_n := \sum_{i=0}^{q-1} \left(S_i^{\otimes n} + (S_i^\Gamma)^{\otimes n} \right). \quad (\text{S93})$$

We will now show that \tilde{S}_n is in fact a feasible point of the minimisation problem that defines $\kappa_1(\rho^{\otimes n}) = 2^{E_\kappa(\rho^{\otimes n})}$. To this end, we need to verify that

$$-\tilde{S}_n \stackrel{?}{\leq} (\rho^{\otimes n})^\Gamma \stackrel{?}{\leq} \tilde{S}_n, \quad \tilde{S}_n^\Gamma \stackrel{?}{\geq} 0. \quad (\text{S94})$$

To prove the first two inequalities, note that

$$\begin{aligned} \tilde{S}_n \pm (\rho^{\otimes n})^\Gamma &= S_0^{\otimes n} + \sum_{i=0}^{q-2} \left(S_{i+1}^{\otimes n} + (S_i^\Gamma)^{\otimes n} \right) + (S_{q-1}^\Gamma)^{\otimes n} \pm (\rho^{\otimes n})^\Gamma \\ &\geq S_0^{\otimes n} + (S_{q-1}^\Gamma)^{\otimes n} \pm (\rho^{\otimes n})^\Gamma \\ &\geq S_0^{\otimes n} \pm (\rho^{\otimes n})^\Gamma \\ &\geq 0, \end{aligned} \quad (\text{S95})$$

where all three inequalities come from (S92). Consequently, we have proved that $-\tilde{S}_n \leq (\rho^{\otimes n})^\Gamma \leq \tilde{S}_n$. Note that the latter implies that $\tilde{S}_n \geq 0$. In addition, by exploiting the definition of \tilde{S}_n in (S93), we also have that $\tilde{S}_n^\Gamma = \tilde{S}_n$, and hence $\tilde{S}_n^\Gamma \geq 0$. We have thus proved that \tilde{S}_n is a feasible point of the minimisation problem that defines (up to a logarithm) $E_\kappa(\rho^{\otimes n})$. Hence, it follows that

$$\begin{aligned} E_\kappa(\rho^{\otimes n}) &\leq \log_2 \text{Tr } \tilde{S}_n \\ &= \log_2 \text{Tr} \left[\sum_{i=0}^{q-1} \left(S_i^{\otimes n} + (S_i^\Gamma)^{\otimes n} \right) \right] \\ &\stackrel{(i)}{=} \log_2 \left(2 \sum_{i=0}^{q-1} (\text{Tr } S_i)^n \right) \\ &\stackrel{(ii)}{\leq} \log_2 \left(2 (\text{Tr } S_{q-1})^n \right) \\ &= \log_2(2q) + n \log_2(\text{Tr } S_{q-1}) \\ &\stackrel{(iii)}{=} \log_2(2q) + n E_{\kappa, q}(\rho). \end{aligned} \quad (\text{S96})$$

Here, in (i) we exploited (S7), (ii) follows from the inequalities $\text{Tr } S_0 \leq \text{Tr } S_1 \leq \dots \leq \text{Tr } S_{q-1}$ (Lemma S9), and finally (iii) holds by the optimality of $S_{-1}, S_0, \dots, S_{q-1}$. As a consequence of (S96), we have that for all fixed $q \in \mathbb{N}^+$

$$E_\kappa^\infty(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} E_\kappa(\rho^{\otimes n}) \leq E_{\kappa,q}(\rho), \quad (\text{S97})$$

which establishes (S89) and thus concludes the proof. \square

Remark S27. Theorem S26 also shows that

$$E_{\kappa,q}^\infty(\rho) := \lim_{n \rightarrow \infty} \frac{1}{n} E_{\kappa,q}(\rho^{\otimes n}) = E_\kappa^\infty(\rho) = E_{c,\text{PPT}}^{\text{exact}}(\rho) \quad \forall \rho, \quad \forall q \in \mathbb{N}^+. \quad (\text{S98})$$

B. A key technical result: proof of Proposition 5

This subsection is devoted to the detailed proof of the crucial Proposition 5, which links the χ - and the κ -hierarchies in a profound way. This connection will be the keystone on which all of our main results are build.

Proposition 5. *For all bipartite states $\rho = \rho_{AB}$ on a system of minimal local dimension $d := \min\{|A|, |B|\} \geq 2$, and all positive integers $p \in \mathbb{N}^+$, it holds that*

$$\kappa_p(\rho) \leq \chi_p(\rho) + \left(\frac{d}{2} - 1\right) (\chi_p(\rho) - \chi_{p-1}(\rho)) \quad (\text{S99})$$

In particular,

$$E_{\chi,p}(\rho) \geq \frac{2}{d} E_{\kappa,p}(\rho) + \left(1 - \frac{2}{d}\right) E_{\chi,p-1}(\rho). \quad (\text{S100})$$

To obtain the above result we first need a key technical lemma, proved below. Before stating it, let us fix some terminology. Given an arbitrary positive semi-definite bipartite operator $T = T_{AB}$, we set

$$d_{\max}(T \| \mathcal{PPT}) := \min \{ \text{Tr } L : T \leq L, \quad L \in \mathcal{PPT} \} = \min \{ \text{Tr } L : T \leq L, \quad L \geq 0, \quad L^\Gamma \geq 0 \}. \quad (\text{S101})$$

It is well known that for all positive semi-definite T it holds that [63]

$$d_{\max}(T \| \mathcal{PPT}) \leq d \text{Tr } T. \quad (\text{S102})$$

One way to see this is, similarly to Corollary S14, by first noticing that for the maximally entangled state one has $d_{\max}(\Phi_d \| \mathcal{PPT}) = d$ and then using the fact that any state $T/\text{Tr } T$ can be obtained from Φ_d by an LOCC transformation, which can never increase $d_{\max}(\cdot \| \mathcal{PPT})$. For completeness, let us give here a more self-contained proof of Eq. (S102).

Up to taking positive linear combinations, it suffices to prove (S102) when $T = |\psi\rangle\langle\psi|$ is the rank-one projector onto the pure state $|\psi\rangle = |\psi\rangle_{AB}$ with Schmidt decomposition $|\psi\rangle_{AB} = \sum_{i=1}^d \sqrt{\lambda_i} |e_i\rangle_A |f_i\rangle_B$. Setting $L := |\psi\rangle\langle\psi|_{AB} + \sum_{i \neq j} \sqrt{\lambda_i \lambda_j} |e_i\rangle\langle e_j|_A \otimes |f_j\rangle\langle f_i|_B$ one sees that on the one hand $L \geq T = |\psi\rangle\langle\psi|$, while on the other

$$L^\Gamma \simeq \bigoplus_i \lambda_i \oplus \bigoplus_{i < j} \sqrt{\lambda_i \lambda_j} \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix} \geq 0. \quad (\text{S103})$$

Then, using Cauchy-Schwarz and noting that $\sum_i \lambda_i = 1$ one finds that

$$\text{Tr } L = 1 + \sum_{i \neq j} \sqrt{\lambda_i \lambda_j} = \sum_i \lambda_i + \sum_{i \neq j} \sqrt{\lambda_i \lambda_j} = \left(\sum_i \sqrt{\lambda_i} \right)^2 \leq d, \quad (\text{S104})$$

which concludes the proof of (S102).

The key lemma that is needed to prove Proposition 5 is the following.

Lemma S28. *For $p \geq 1$, let $S_{-1}, S_0, \dots, S_{p-1}, S_p$ be optimisers for the SDP (S30) that defines $\chi_p(\rho)$. Then there exists a PPT operator $M = M_{AB} \in \mathcal{PPT}$ such that $S_{-1}, S_0, \dots, S_{p-2}, S_{p-1} + M$ are valid ansatzes for the SDP that defines $\kappa_p(\rho)$ as in (S36), and moreover $\text{Tr } M = d_{\max}((S_{p-1}^\Gamma)_- \| \mathcal{PPT})$.*

Proof. Since $-S_i \leq S_{i-1}^\Gamma \leq S_i$ holds for all $i = 0, \dots, p-1$ by assumption, and moreover $M \geq 0$, the only inequality that is left to verify is $(S_{p-1} + M)^\Gamma \geq 0$. By definition of $d_{\max}(\cdot \| \mathcal{PPT})$, there exists a PPT operator $L = L_{AB}$ such that $(S_{p-1}^\Gamma)_- \leq L$ and $\text{Tr } L = d_{\max}((S_{p-1}^\Gamma)_- \| \mathcal{PPT})$. We can then define $M := L^\Gamma$, so that

$$(S_{p-1} + M)^\Gamma = S_{p-1}^\Gamma + L = (S_{p-1}^\Gamma)_+ - (S_{p-1}^\Gamma)_- + L \geq (S_{p-1}^\Gamma)_+ \geq 0. \quad (\text{S105})$$

Since $\text{Tr } M = \text{Tr } L = d_{\max}((S_{p-1}^\Gamma)_- \| \mathcal{PPT})$, this concludes the proof of the lemma. \square

We are now ready to give the complete proof of Proposition 5.

Proof of Proposition 5. Let $S_{-1}, S_0, \dots, S_{p-1}, S_p$ be optimisers for the SDP (S30) that defines $\chi_p(\rho)$. Then clearly $S_{-1}, S_0, \dots, S_{p-1}$ are valid ansatzes for the SDP that defines $\chi_{p-1}(\rho)$. Therefore,

$$\text{Tr } S_{p-1} \geq \chi_{p-1}(\rho) = \chi_p(\rho) - \Delta_p = \text{Tr } S_p - \Delta_p, \quad (\text{S106})$$

where we defined $\Delta_p := \chi_p(\rho) - \chi_{p-1}(\rho)$. We deduce that

$$0 \leq \delta_p := \text{Tr } [S_p - S_{p-1}] \leq \Delta_p. \quad (\text{S107})$$

The first inequality is a consequence of Lemma S11 — basically, of the fact that $\delta_p = \text{Tr } [S_p - S_{p-1}^\Gamma]$, and the operator inside the trace is positive semi-definite. Now, by Lemma S28 there exists a PPT operator M such that $S_{-1}, S_0, \dots, S_{p-2}, S_{p-1} + M$ are legitimate ansatzes for the SDP that defines $\kappa_p(\rho)$ as in (S36), and moreover

$$\begin{aligned} \text{Tr } M &= d_{\max}((S_{p-1}^\Gamma)_- \| \mathcal{PPT}) \\ &\stackrel{(i)}{\leq} d \text{Tr}[(S_{p-1}^\Gamma)_-] \\ &= d \frac{\|S_{p-1}^\Gamma\|_1 - \text{Tr } S_{p-1}}{2} \\ &\stackrel{(ii)}{\leq} d \frac{\text{Tr } S_p - \text{Tr } S_{p-1}}{2} \\ &= \frac{d\delta_p}{2}, \end{aligned} \quad (\text{S108})$$

where the bound in (i) follows from (S102), and that in (ii) comes from Lemma S1 applied to the operator inequalities $-S_p \leq S_{p-1}^\Gamma \leq S_p$. (The latter is, in fact, an equality if the operators S_i are optimisers for $\chi_p(\rho)$; however, we shall not make use of this observation.)

Putting all together, we obtain that

$$\begin{aligned} \kappa_p(\rho) &\leq \text{Tr } [S_{p-1} + M] \\ &= \text{Tr } S_p - \delta_p + \text{Tr } M \\ &\stackrel{(iii)}{=} \chi_p(\rho) - \delta_p + \text{Tr } M \\ &\stackrel{(iv)}{\leq} \chi_p(\rho) + \left(\frac{d}{2} - 1\right) \delta_p \\ &\stackrel{(v)}{\leq} \chi_p(\rho) + \left(\frac{d}{2} - 1\right) \Delta_p \\ &= \chi_p(\rho) + \left(\frac{d}{2} - 1\right) (\chi_p(\rho) - \chi_{p-1}(\rho)), \end{aligned} \quad (\text{S109})$$

which reproduces (S99). Here, (iii) follows by the optimality of S_{-1}, S_0, \dots, S_p for $\chi_p(\rho)$, in (iv) we employed (S108), and finally (v) follows from (S107).

Rearranging, taking the logarithms, and using the concavity of the \log_2 function, one obtains also (S100). \square

C. What have we achieved so far?

Convergence

Proposition 5 is the core technical finding on which all of our main results rest. Now that we have proved it, most of the remaining proofs are comparatively straightforward. To show the importance of Proposition 5, it is instructive to pause for a second and explore some of its immediate consequences. In this subsection we do precisely that.

We first show that Proposition 5 already implies (8), i.e. that the limits of the χ - and κ -hierarchies coincide and yield the true zero-error PPT entanglement cost. In fact, taking the limit $p \rightarrow \infty$ on both sides of (S99) and using the fact that both $\lim_{p \rightarrow \infty} \chi_p(\rho)$ and $\lim_{p \rightarrow \infty} \kappa_p(\rho)$ exist by monotonicity (Proposition S11) shows that

$$\lim_{p \rightarrow \infty} \kappa_p(\rho) \leq \lim_{p \rightarrow \infty} \chi_p(\rho). \quad (\text{S110})$$

Combined with the reverse inequality, which is an obvious consequence of (S41), one deduces that indeed $\lim_{p \rightarrow \infty} \kappa_p(\rho) = \lim_{p \rightarrow \infty} \chi_p(\rho)$. Now, going back to Theorem S26, and in particular to (S90), we see that this implies that in fact

$$\lim_{p \rightarrow \infty} E_{\kappa,p}(\rho) = \lim_{p \rightarrow \infty} E_{\chi,p}(\rho) = E_{\kappa}^{\infty}(\rho) = E_{c,\text{PPT}}^{\text{exact}}(\rho), \quad (\text{S111})$$

i.e. the four central quantities appearing in (S90) coincide for all states. We will re-derive this equalities with some additional guarantees on the speed of convergence while proving Theorem 1 below.

The identity in (S111) is remarkable because it provides a precise connection between the zero-error PPT cost, given by the regularisation of E_{κ} and thus by a limit $n \rightarrow \infty$ over the number of copies n , and the limits of the hierarchies, intended as limits $p \rightarrow \infty$ on the hierarchy level. The fact that there would be any sort of connection between the asymptotic limit in the number of copies and that in the hierarchy level was a priori totally unclear, and it should be regarded as one of the neater results of our approach. However, Eq. (S111) is still not completely satisfactory from a computational standpoint, because it does not provide any rigorous guarantee on the speed of convergence to the limit. Having such guarantees (either on the limit in n or on that in p) would turn (S111) into an *algorithm*, because it would tell us at which n or p we would need to stop in order to achieve a certain approximation. If that could be done, then, due to the fact that $E_{\kappa}(\rho^{\otimes n})$, $E_{\chi,p}(\rho)$, and $E_{\kappa,p}(\rho)$ are all computable via SDPs, we would know how to approximate $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ up to an arbitrary accuracy. In short, whether the algorithm we have described works or not depends on whether or not we can say something about the speed of convergence to the limit in either n or p . Furthermore, even if the algorithm does exist, then whether it is efficient or not depends on the actual speed of convergence to the limit, either in n or in p .

The key observation underpinning our entire approach is that while there seems to be no easy way of achieving either of the above two goals when dealing with the limit in the number of copies n , the situation changes dramatically for the better when considering the limit in the hierarchy level p . Not only will we be able to give universal, uniform bounds on the convergence speed, but the resulting convergence will also turn out to be exponentially fast (Theorem 1), resulting in an *efficient* algorithm to calculate the zero-error PPT entanglement cost on all states (Theorem 3).

The local qubit case

Proposition 5 also implies an immediate simple solution to the problem of computing $E_{c,\text{PPT}}^{\text{exact}}(\rho_{AB})$ when either A or B is a single-qubit system. In this case, it turns out that both hierarchies collapse at the first level ($p = 1$), entailing that $E_{\chi} = E_{\kappa} = E_{c,\text{PPT}}^{\text{exact}}$ already gives the true zero-error PPT cost.

Corollary 2. *Let AB be a bipartite quantum system in which either A or B is a single qubit, i.e. $d = \min\{|A|, |B|\} = 2$. Then for all states $\rho = \rho_{AB}$ it holds that*

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\chi}(\rho) = E_{\kappa}(\rho), \quad (\text{S112})$$

where E_{χ} and E_{κ} are defined by (S27) and (S13), respectively.

Note. Remember that we identify $E_{\chi} = E_{\chi,1}$ (cf. (S27) and (S29)) and $E_{\kappa} = E_{\kappa,1}$ (cf. (S13) and (S35)).

Proof. Writing down (S99) for $d = 2$ and $p = 1$ shows that $\kappa_1(\rho) \leq \chi_1(\rho)$. Combining this with Theorem S26 then implies the claim. \square

Remark S29. The above Corollary 2 does not tell us what happens to the zeroth level of the χ -hierarchy, namely, the logarithmic negativity. An older result by Ishizaka [30] states that all *two-qubit* states have zero bi-negativity, implying, by Lemma S12, that

$$|A| = |B| = 2 \implies E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_N(\rho) = \log_2 \|\rho^\Gamma\|_1 \quad \forall \rho = \rho_{AB}. \quad (\text{S113})$$

When e.g. $|A| = 2$ but $|B| > 2$, Corollary 2 guarantees that $E_{c,\text{PPT}}^{\text{exact}} = E_\chi = E_\kappa$, but it would be even better if one could establish a closed-form expression for this quantity. One way to achieve this, for instance, would be by generalising Ishizaka's result so as to encompass all qubit-qudit systems. If that could be done, then again we would find that $E_{c,\text{PPT}}^{\text{exact}} = E_N$, which would be our sought closed-form expression. We leave a full understanding of the role of the logarithmic negativity in the qubit-qudit case as an open problem.

D. Exponential convergence: proof of Theorem 1

This subsection is devoted to the proof of our first main result, Theorem 1, which will be seen to be a relatively straightforward consequence of the key Proposition 5.

Theorem 1. *For all bipartite states $\rho = \rho_{AB}$ on a system of minimal local dimension $d := \min\{|A|, |B|\} \geq 2$, and all positive integers $p \in \mathbb{N}^+$, it holds that*

$$E_{\chi,p}(\rho) \leq E_{c,\text{PPT}}^{\text{exact}}(\rho) \leq E_{\kappa,p}(\rho) \leq E_{\chi,p}(\rho) + \log_2 \frac{1}{1 - (1 - \frac{2}{d})^p}. \quad (\text{S114})$$

In particular,

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_\kappa^\infty(\rho) = \lim_{p \rightarrow \infty} E_{\chi,p}(\rho) = \lim_{p \rightarrow \infty} E_{\kappa,p}(\rho), \quad (\text{S115})$$

with the convergence in p being exponentially fast uniformly on ρ .

Proof. We give here a slightly modified and extended version of the argument presented in the main text. The reason for doing so is to incorporate also the convergence of the κ -hierarchy in a single statement. To deal with the χ - and κ -hierarchies simultaneously, it is useful to set by convention $\kappa_0(\rho) := d$. Now, define the numbers

$$\varepsilon_p(\rho) := 1 - \frac{\chi_p(\rho)}{\kappa_p(\rho)}, \quad (\text{S116})$$

which satisfy $\varepsilon_p \in [0, 1]$ for all $p \in \mathbb{N}$ due to Proposition S11. Note that this definition differs from that employed in the main text (Eq. (24)) — in particular, the quantity in (S116) is larger than that in (24). We can now write

$$\begin{aligned} 1 &\stackrel{(i)}{\leq} \frac{d}{2} \frac{\chi_p(\rho)}{\kappa_p(\rho)} - \left(\frac{d}{2} - 1\right) \frac{\chi_{p-1}(\rho)}{\kappa_p(\rho)} \\ &\stackrel{(ii)}{\leq} \frac{d}{2} \frac{\chi_p(\rho)}{\kappa_p(\rho)} - \left(\frac{d}{2} - 1\right) \frac{\chi_{p-1}(\rho)}{\kappa_{p-1}(\rho)} \\ &\stackrel{(iii)}{=} 1 - \frac{d}{2} \varepsilon_p + \left(\frac{d}{2} - 1\right) \varepsilon_{p-1}. \end{aligned} \quad (\text{S117})$$

The justification of the above steps is as follows: the inequality in (i) is just a rephrasing of that established by Proposition 5, obtained by dividing both sides by $\kappa_p(\rho)$ (remember that $\kappa_p(\rho) \geq 1$ by Remark S10); in (ii) we observed that $\kappa_p(\rho) \leq \kappa_{p-1}(\rho)$ due to Proposition S11 (this is also true for $p = 1$ due to Lemma S12) and $d \geq 2$; finally, in (iii) we employed the definition of ε_p in (S116).

We now see that (S117) can be readily massaged into

$$\varepsilon_p \leq \left(1 - \frac{2}{d}\right) \varepsilon_{p-1}. \quad (\text{S118})$$

Iterating this p times and using the fact that $\varepsilon_0 \leq 1$ and $\varepsilon_p \geq 0$ yields immediately

$$0 \leq \varepsilon_p \leq \left(1 - \frac{2}{d}\right)^p, \quad (\text{S119})$$

which in turn can be rephrased as

$$\kappa_p(\rho) \leq \frac{\chi_p(\rho)}{1 - (1 - \frac{2}{d})^p}. \quad (\text{S120})$$

Taking the logarithms of both sides proves the last inequality in (S114). The first three inequalities are already known from Theorem S90.

Taking the limit $p \rightarrow \infty$ establishes also the convergence of both hierarchies to the true zero-error PPT entanglement cost (Eq. (S115)). Note that the speed of convergence depends on d but not on ρ . \square

The above result is central in our approach, because it provides a quantitative guarantee on what level of the hierarchies we have to resort to in order to obtain a prescribed approximation of the true value of the zero-error PPT entanglement cost. Before we proceed with the description of an algorithm that builds on this observation, we take note of two remarkable consequences of Theorem 1, namely, the full additivity of the zero-error PPT entanglement cost and its continuity.

Intuitively, the former means that the cheapest way to generate many copies of a state of the form $\rho_{AB} \otimes \omega_{A'B'}$, where A and A' belong to Alice, and B and B' to Bob, is to manufacture many copies of ρ_{AB} and $\omega_{A'B'}$ *separately*: in other words, there is no advantage to be gained in considering joint protocols. The continuity of the zero-error PPT cost, instead, allows us to estimate the cost of a state that is close enough to one for which the cost is known. The proofs of both of these facts are paradigmatic examples of how the knowledge we have gathered so far allows us to say a lot about the problem of zero-error PPT entanglement dilution even without a closed-form expression for the corresponding cost.

Corollary S30. *The zero-error entanglement cost under PPT operations is fully tensor additive, i.e. for all pairs of bipartite states $\rho_{AB}, \omega_{A'B'}$ it holds that*

$$E_{c,\text{PPT}}^{\text{exact}}(\rho_{AB} \otimes \omega_{A'B'}) = E_{c,\text{PPT}}^{\text{exact}}(\rho_{AB}) + E_{c,\text{PPT}}^{\text{exact}}(\omega_{A'B'}). \quad (\text{S121})$$

Proof. It suffices to take the limit $p \rightarrow \infty$ of (S66) using (S115). \square

Corollary S31 (Continuity of the zero-error PPT entanglement cost). *Let ρ, ρ' be two quantum states on a bipartite system AB with minimal local dimension $d := \min\{|A|, |B|\}$. Let $\varepsilon := \frac{1}{2} \|\rho - \rho'\|_1$. Then*

$$\left| 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} - 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho')} \right| \leq 2d\varepsilon, \quad (\text{S122})$$

and therefore

$$\left| E_{c,\text{PPT}}^{\text{exact}}(\rho) - E_{c,\text{PPT}}^{\text{exact}}(\rho') \right| \leq \log_2(1 + 2d\varepsilon) \leq 2(\log_2 e) d\varepsilon. \quad (\text{S123})$$

Proof. It suffices to take the limit $p \rightarrow \infty$ of (S82) and (S83) written for $F = \chi_p$ and $f = E_{\chi,p}$, using Theorem 1, and in particular (S115), to simplify the left-hand sides. \square

E. An explicit algorithm to compute the zero-error PPT cost: proof of Theorem 3

We will now exploit the above Theorem 1 to prove the last of our main result, Theorem 3. To this end, we will design an efficient algorithm that computes the zero-error PPT entanglement cost up to any desired accuracy.

Theorem 3 (Efficient algorithm to compute the cost). *Let AB be a bipartite quantum system of total dimension $D := |AB| = |A||B|$ and minimal local dimension $d := \min\{|A|, |B|\}$. Then there exists an algorithm that for an arbitrary state $\rho = \rho_{AB}$ computes $2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)}$ up to a multiplicative error ε — and hence, a fortiori, $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ up to an additive error ε — in time*

$$\mathcal{O}\left((dD)^6 \text{polylog}(p, D, d, 1/\varepsilon)\right) = \mathcal{O}\left((dD)^{6+o(1)} \text{polylog}(1/\varepsilon)\right). \quad (\text{S124})$$

Proof. To start, choose

$$p := \left\lceil \frac{\ln(2d/\varepsilon)}{\ln \frac{d}{d-2}} \right\rceil, \quad (\text{S125})$$

so that

$$\left(1 - \frac{2}{d}\right)^p \leq \frac{\varepsilon}{2d}. \quad (\text{S126})$$

Due to (S114), we then know that

$$1 - \frac{\varepsilon}{2d} \leq \frac{\chi_p(\rho)}{2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)}} \leq 1, \quad (\text{S127})$$

which implies that

$$\left| \chi_p(\rho) - 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \right| \leq \frac{\varepsilon}{2d} 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \leq \frac{\varepsilon}{2}, \quad (\text{S128})$$

where the last equality is a consequence of the fact that $E_{c,\text{PPT}}^{\text{exact}}(\rho) \leq \log_2 d$ for all ρ by Lemma S12.

Now, $\chi_p(\rho)$ is given by an SDP and hence it can be computed efficiently up to additive error $\varepsilon/2$. We will look at the time it takes to carry out this computation in a moment, but for the time being assume that it yields an estimator $\widehat{\chi}_p(\rho)$ with the property that

$$\left| \widehat{\chi}_p(\rho) - \chi_p(\rho) \right| \leq \frac{\varepsilon}{2}. \quad (\text{S129})$$

Then by combining (S128) and (S129) one obtains that

$$\left| \widehat{\chi}_p(\rho) - 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \right| \leq \varepsilon. \quad (\text{S130})$$

This yields the claimed computation of $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ up to an additive error ε . To estimate the error one incurs when computing $E_{c,\text{PPT}}^{\text{exact}}(\rho)$, it suffices to observe that

$$\left| E_{c,\text{PPT}}^{\text{exact}}(\rho) - \log_2 \widehat{\chi}_p(\rho) \right| \leq (\log_2 e) \left| \widehat{\chi}_p(\rho) - 2^{E_{c,\text{PPT}}^{\text{exact}}(\rho)} \right| \leq \varepsilon \log_2 e, \quad (\text{S131})$$

so the error on $E_{c,\text{PPT}}^{\text{exact}}(\rho)$ is also ε , up to a constant. In the above derivation, we used the estimate $|a-b| \leq (\log_2 e)|2^a - 2^b|$, valid for all $a, b \geq 0$.

We still need to estimate the running time of the SDP that computes the estimator $\widehat{\chi}_p(\rho)$. To do that, we use the best known SDP solvers, which have time complexity [51] (see also the quantum-friendly review in [52])

$$\mathcal{O}\left(m \left(m^2 + n^\omega + mns\right) \text{polylog}(m, n, R, 1/\varepsilon)\right) \quad (\text{S132})$$

once the SDP is put in the standard form

$$\begin{aligned} & \max. \quad \text{Tr } CX \\ & \text{s.t.} \quad X \geq 0, \\ & \quad \text{Tr } A_j X \leq b_j, \quad j = 1, \dots, m, \end{aligned} \quad (\text{S133})$$

where $\|A_j\|_\infty, \|C\|_\infty \leq 1$ for $j = 1, \dots, m$, all matrices are $n \times n$, R is any upper bound on the optimal value, $\omega \in [2, 2.373)$ is the matrix multiplication exponent, and s is the sparsity, i.e. maximum number of non-zero entries in any row of the input matrices A_j, C .

What we have to do now is to cast the SDP for $\chi_p(\rho)$ (Eq. (S30)) in the standard form (S133). This can be done by restricting the variable X to be of the form

$$X \longrightarrow \bigoplus_{i=0}^p \begin{pmatrix} S_i & S_{i-1}^\Gamma \\ S_{i-1}^\Gamma & S_i \end{pmatrix}, \quad (\text{S134})$$

where $S_{-1} = \rho$. To see why, note that for a matrix of the above block form positive semi-definiteness implies that

$$\begin{pmatrix} S_i & S_{i-1}^\Gamma \\ S_{i-1}^\Gamma & S_i \end{pmatrix} \geq 0 \quad \forall i = 0, \dots, p, \quad (\text{S135})$$

which by a simple unitary rotation can be rephrased as

$$\frac{1}{2} \begin{pmatrix} \mathbb{1} & \mathbb{1} \\ \mathbb{1} & -\mathbb{1} \end{pmatrix} \begin{pmatrix} S_i & S_{i-1}^\Gamma \\ S_{i-1}^\Gamma & S_i \end{pmatrix} \begin{pmatrix} \mathbb{1} & \mathbb{1} \\ \mathbb{1} & -\mathbb{1} \end{pmatrix}^\dagger = \begin{pmatrix} S_i + S_{i-1}^\Gamma & 0 \\ 0 & S_i - S_{i-1}^\Gamma \end{pmatrix} \geq 0 \quad \forall i = 0, \dots, p, \quad (\text{S136})$$

i.e. $S_i \pm S_{i-1}^\Gamma \geq 0$ for all $i = 0, \dots, p$, matching the positive semi-definite constraints in (S30).

The above reasoning tells us that it is a good idea to try to enforce the structure (S134) on X . This fixes the size of the matrices to be

$$n = 2(p+1)D = \mathcal{O}(pD), \quad (\text{S137})$$

where $D = |A||B|$ is the total dimension. The problem therefore becomes that of ensuring that X has the structure in (S134). To this end, it is useful to think of X as a block matrix of size $(p+1) \times (p+1)$, where each block has size $2D \times 2D$. We need to make sure that:

- (1) all the $p(p+1)$ off-diagonal blocks are zero;
- (2) inside each diagonal block X_{ii} ($i = 0, \dots, p$), which can in turn be thought of as a 2×2 block matrix $X_{ii} = \begin{pmatrix} H_i & K_i \\ L_i & M_i \end{pmatrix}$ with each block having size D , we have that $H_i = M_i$ and $K_i = L_i$;
- (3) for all $i = 1, \dots, p$, $K_i = H_{i-1}^\Gamma$; and
- (4) $K_0 = \rho^\Gamma$.

All of these constraints are equalities instead of inequalities, but any equality can be written as two inequalities, and we do not count factors of 2. How many linear equality constraints on X do we need?

- (1): $p(p+1)D^2$ constraints;
- (2): $2(p+1)D^2$ constraints;
- (3): pD^2 constraints;
- (4): D^2 constraints.

We are therefore dealing with a total of $(p+1)(p+3)D^2$ linear equality constraints. It is therefore clear that we can set

$$m = \mathcal{O}(p^2 D^2). \quad (\text{S138})$$

Now, constraints (1)–(3) are of the form $\text{Tr} AX = 0$, where A has only two non-zero elements on the same row or column, while the constraints (4) are of the form $\text{Tr} AX = b$, where A has a single non-zero entry (equal to 1). Note also that since we want $\text{Tr} CX = \text{Tr} S_p$, C needs to be block diagonal, with the first p diagonal blocks equal to zero and the last one equal to $1/2$ (all blocks are of size $2D$). We can thus set

$$s = \mathcal{O}(1). \quad (\text{S139})$$

As for R , by Lemma S12 we already know that

$$R = d \quad (\text{S140})$$

is an upper bound on $\chi_p(\rho)$. Plugging the values (S137)–(S140) into (S132) yields a time complexity

$$\begin{aligned} \mathcal{O}\left((pD)^6 \text{polylog}(p, D, d, 1/\varepsilon)\right) &= \mathcal{O}\left((dD)^6 \text{polylog}(p, D, d, 1/\varepsilon)\right) \\ &= \mathcal{O}\left((dD)^{6+o(1)} \text{polylog}(1/\varepsilon)\right), \end{aligned} \quad (\text{S141})$$

where we used the fact that $p = \mathcal{O}(d \log(d/\varepsilon))$ due to (S125). \square

Remark S32. It can be verified that the time complexity of solving the SDP that defines the negativity E_N is $\mathcal{O}(D^{6+o(1)} \text{polylog}(1/\varepsilon))$. While this is smaller than the complexity for computing $E_{c, \text{PPT}}^{\text{exact}}$, it is only marginally so — the difference being a mere factor $\text{poly}(d)$.

F. Open problem: hierarchy collapse

Although we showed that evaluating the limit $\lim_{p \rightarrow \infty} E_{\chi,p}(\rho)$ to any desired precision is computationally not much more demanding than evaluating $E_{\chi,p}$ for a fixed level of the hierarchy p , one may still wonder — perhaps from an analytical or aesthetic standpoint — whether computing the limit is truly necessary, or whether the hierarchy collapses at a certain point and there exists a finite p^* such that $E_{\chi,p} = E_{\chi,p^*} \forall p \geq p^*$. An analogous question can be asked for the κ -hierarchy.

One immediate consequence of Proposition 5 is that, if $E_{\chi,p}(\rho) = E_{\chi,p-1}(\rho)$ for some p , then $E_{\kappa,p}(\rho) \leq E_{\chi,p}(\rho)$, which would imply a complete collapse of both hierarchies, that is,

$$E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\chi,p}(\rho) = E_{\kappa,p}(\rho), \quad (\text{S142})$$

due to Proposition S11.

We were not able to confirm nor disprove that this happens. However, numerical evidence strongly suggests that the hierarchies collapse, and indeed they do so already at the second level of the χ -hierarchy. This leads us to posit the following conjecture.

Conjecture S33. *For all states $\rho = \rho_{AB}$, the χ -quantities defined by (S30) satisfy that $E_{\chi,3}(\rho) = E_{\chi,2}(\rho)$. As a consequence, the χ - and κ -hierarchies collapse and $E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\chi,2}(\rho) = E_{\kappa,3}(\rho)$.*

In fact, we were not even able to find a gap between $E_{\chi,2}$ and $E_{\kappa,2}$, so one could even make the stronger conjecture that $E_{c,\text{PPT}}^{\text{exact}}(\rho) = E_{\chi,2}(\rho) = E_{\kappa,2}(\rho)$ holds for all states, which would entail that both hierarchies collapse at the second level. For the sake of obtaining a single-letter, limit-free formula for $E_{c,\text{PPT}}^{\text{exact}}$, it would be very interesting to resolve this question in future work. However, as remarked in the main text, doing so would only amount to a $\text{poly}(d)$ reduction of the computational complexity of calculating the zero-error PPT entanglement cost $E_{c,\text{PPT}}^{\text{exact}}$.

V. FROM ZERO ERROR TO VERY SMALL ERROR

Zero-error entanglement manipulation tasks may seem on the surface un-physical, because nothing in nature happens with zero error, and small, undetectable errors should therefore always be included into the picture. Here we will do precisely that, and we will find that *if those errors are assumed to be sufficiently small (but non-zero), the resulting entanglement dilution rates are the same as in the zero-error case*. In this context, ‘sufficiently small’ could mean for example ‘going to 0 super-exponentially in the number of copies of the given state’.

This is no different from what happens in classical information theory, where the capacity for communicating on a noisy classical channel with error probability going to 0 super-exponentially equals its zero-error capacity, defined by Shannon in his landmark 1956 paper [64]. The argument to prove this claim seems to be part of the folklore in information theory, and it was brought to our attention by Andreas Winter. It goes as follows. A classical channel \mathcal{N} with input alphabet \mathcal{X} and output alphabet \mathcal{Y} can be thought of as a transition matrix $N(y|x) \geq 0$, with $\sum_y N(y|x) = 1$ for all $x \in \mathcal{X}$. A code to communicate on $\mathcal{N}^{\otimes n}$ is a list of words x_1^n, \dots, x_M^n , where $M = 2^{\lceil nR \rceil}$, and R is the rate of the code. Suppose that for a fixed n a word in $\{1, \dots, M\}$ is drawn at random. Whatever the decoder is, the probability of making an error can be estimated from below as follows. Pick two distinct $i, j \in \{1, \dots, M\}$. For a fixed $y^n \in \mathcal{Y}^n$, call $p_i := \mathcal{N}^{\otimes n}(y^n|x_i^n)$ and $p_j := \mathcal{N}^{\otimes n}(y^n|x_j^n)$ the probabilities that these two words get transformed into the same output word $y^n \in \mathcal{Y}^n$. When that happens, even if the decoding party knows that the message was either i or j all they can do is to guess i or j with a maximum likelihood rule, i.e. i with probability $\frac{p_i}{p_i+p_j}$ and j with probability $\frac{p_j}{p_i+p_j}$. The total probability of error is thus at least

$$\frac{1}{M} p_i \frac{p_j}{p_i+p_j} + \frac{1}{M} p_j \frac{p_i}{p_i+p_j} = \frac{1}{M} \frac{2p_i p_j}{p_i+p_j} \geq \frac{1}{M} \min\{p_i, p_j\}. \quad (\text{S143})$$

Now, if the right-hand side is not zero for some y^n , then it must be at least t^n/M , where $t := \min_{x,y: N(y|x)>0} N(y|x)$. Therefore the probability of error of the whole process (encoding, transmission, and decoding) is at least $t^n 2^{-\lceil nR \rceil}$, which cannot go to 0 super-exponentially. This completes the summary of the state of affairs in classical communication over noisy channels.

To arrive at analogous conclusions in the case of PPT entanglement dilution, we first need to fix some terminology. Our first task is to design a way to control the errors incurred in an arbitrary entanglement manipulation protocol.

This can be done by introducing a rate-error pair achievability region for entanglement manipulation under a given set of free operations \mathcal{F} , defined as follows.

Definition S34. For two bipartite states $\rho = \rho_{AB}$ and $\omega = \omega_{AB}$ and a class of free operations \mathcal{F} , we say that the rate-error pair (r, s) is achievable for the transformation $\rho \rightarrow \omega$ with operations in \mathcal{F} if there exists a sequence of protocols $(\Lambda_n)_{n \in \mathbb{N}_+}$, with $\Lambda_n \in \mathcal{F}(A^n B^n \rightarrow A'^n B'^n)$ for all $n \in \mathbb{N}_+$, such that

$$\liminf_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log_2 \frac{1}{2} \|\Lambda_n(\rho^{\otimes \lfloor rn \rfloor}) - \omega^{\otimes n}\|_1 \right\} \geq s. \quad (\text{S144})$$

In our case, we care about the case where $\mathcal{F} = \text{PPT}$ is the set of PPT operations, the initial state is the ebit Φ_2 , and the final state is an arbitrary ρ . In this case, a rate-error pair that is achievable for $\Phi_2 \rightarrow \rho$ under PPT operations is also called an achievable rate-error pair for PPT entanglement dilution to ρ . The main result of this section is as follows.

Theorem S35. Let $\rho = \rho_{AB}$ be an arbitrary finite-dimensional bipartite state. If (r, s) is achievable for PPT entanglement dilution to ρ , then

$$\max \{r, \log_2 d - s\} \geq E_{c, \text{PPT}}^{\text{exact}}(\rho). \quad (\text{S145})$$

In particular, for all $s > \log_2 d - E_{c, \text{PPT}}^{\text{exact}}(\rho)$ the rate-error pair (r, s) is achievable if and only if $r \geq E_{c, \text{PPT}}^{\text{exact}}(\rho)$.

Before proving the above result, let us discuss its implications. What this shows is that if the error is required to decay *sufficiently fast*, i.e. faster than $2^{-n(\log_2 d - E_{c, \text{PPT}}^{\text{exact}}(\rho))}$ asymptotically, then the optimal rate of entanglement dilution coincides with its zero-error value. In other words, once a certain threshold in s is passed, it does not make a difference for the rate r whether s increases further, even if it goes all the way to infinity. Note that Theorem S35 applies to two suggestive special cases, namely where (i) we require the error to decay faster than d^{-n} ; and therefore also when (ii) we require a super-exponential decay law. In both of these cases, the relevant optimal rate of PPT entanglement dilution coincides with the zero-error PPT entanglement cost $E_{c, \text{PPT}}^{\text{exact}}(\rho)$.

Proof of Theorem S35. Consider a sequence of PPT operations $(\Lambda_n)_{n \in \mathbb{N}_+}$ such that

$$\varepsilon_n := \frac{1}{2} \|\Lambda_n(\Phi_2^{\otimes \lfloor rn \rfloor}) - \rho^{\otimes n}\|_1 \quad (\text{S146})$$

exhibits the asymptotic decay rate

$$\liminf_{n \rightarrow \infty} \left\{ -\frac{1}{n} \log_2 \varepsilon_n \right\} \geq s. \quad (\text{S147})$$

Then,

$$\begin{aligned} 2^{\lfloor rn \rfloor} &= \kappa_1 \left(\Phi_2^{\otimes \lfloor rn \rfloor} \right) \\ &\stackrel{(i)}{\geq} \kappa_1 \left(\Lambda_n \left(\Phi_2^{\otimes \lfloor rn \rfloor} \right) \right) \\ &\stackrel{(ii)}{\geq} \kappa_1 \left(\rho^{\otimes n} \right) - 2d^n \varepsilon_n \end{aligned} \quad (\text{S148})$$

where (i) is a consequence of the monotonicity of $\kappa_1 = 2^{E_\kappa}$ under PPT operations (Proposition S13), while (ii) is an application of Proposition S23, and in particular of (S82) with $F = \kappa_1$. Therefore,

$$\begin{aligned} 1 + \max \{ \lfloor rn \rfloor, 1 + n \log_2 d + \log_2 \varepsilon_n \} &= \log_2 \left(2 \max \{ 2^{\lfloor rn \rfloor}, 2d^n \varepsilon_n \} \right) \\ &\geq \log_2 \left(2^{\lfloor rn \rfloor} + 2d^n \varepsilon_n \right) \\ &\geq \log_2 \kappa_1 \left(\rho^{\otimes n} \right) \\ &= E_\kappa \left(\rho^{\otimes n} \right). \end{aligned} \quad (\text{S149})$$

Dividing by n and taking the lim sup of both sides as $n \rightarrow \infty$ yields precisely

$$\begin{aligned}
 \max\{r, \log_2 d - s\} &\geq \limsup_{n \rightarrow \infty} \frac{1}{n} (1 + \max\{\lfloor rn \rfloor, 1 + n \log_2 d + \log_2 \varepsilon_n\}) \\
 &\geq \limsup_{n \rightarrow \infty} \frac{1}{n} E_\kappa(\rho^{\otimes n}) \\
 &= \lim_{n \rightarrow \infty} \frac{1}{n} E_\kappa(\rho^{\otimes n}) \\
 &= E_\kappa^\infty(\rho),
 \end{aligned} \tag{S150}$$

concluding the proof. □