Dirac operators on the half-line: stability of spectrum and non-relativistic limit

David Kramár and David Krejčiřík

Department of Mathematics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Trojanova 13, 12000 Prague 2, Czech Republic kramada1@fjfi.cvut.cz, david.krejcirik@fjfi.cvut.cz

16 May 2024

Abstract

We consider Dirac operators on the half-line, subject to generalised infinite-mass boundary conditions. We derive sufficient conditions which guarantee the stability of the spectrum against possibly non-self-adjoint potential perturbations and study the optimality of the obtained results. Finally, we establish a non-relativistic limit which makes a relationship of the present model to the Robin Laplacian on the half-line.

1 Introduction

According to classical physics, the electrons in atoms would have collapsed into the nucleus in a matter of nanoseconds [21], which is in direct contradiction to our experience. This paradox was one of the impetuses leading to the development of quantum mechanics which particularly explains the stability of matter [26]. Mathematically, this is a consequence of the stability of the spectrum of the Laplacian $-\Delta$ in \mathbb{R}^3 against additive perturbations $V: \mathbb{R}^3 \to \mathbb{C}$ satisfying the subordination condition [22, 15, 19]

$$\forall \psi \in W^{1,2}(\mathbb{R}^3), \qquad \int_{\mathbb{R}^3} |V| |\psi|^2 \le a \int_{\mathbb{R}^3} |\nabla \psi|^2, \tag{1}$$

with a < 1. Indeed, (1) together with the Hardy inequality implies that the spectrum of the Coulomb Hamiltonian is bounded from below. The strength of the condition (1) is that it ensures not only the set stability $\sigma(-\Delta + V) = \sigma(-\Delta)$, but also the absence of embedded eigenvalues. Moreover, the fact that V is allowed to be complex-valued makes the spectral analysis substantially more challenging, while the problem is still important for applications [25].

There exist generalisations of the subordination condition (1) to higher dimensions [22, 16, 15, 19]. Because of the absence of Hardy inequality, the condition is void in dimensions 1 and 2. However, there exist two-dimensional analogues for the magnetic Laplacian [14, 4]. To get a non-trivial result in dimension one, it is necessary to restrict the Laplacian to a half-line, subject to repulsive boundary conditions of Robin type [5, 23].

The motivation of the present paper is to establish one-dimensional analogues of (1) in the relativistic setting of Dirac operators on the half-line. While this model was considered previously [6, 13], the stability of the spectrum has not been addressed and our objective is to fill in this gap. For an analogous stability problem for the Dirac operator in higher dimensions, see [10, 4, 11, 27].

To be more specific, given $m \geq 0$ (mass), we consider the unperturbed Dirac operator

$$D_0 := \begin{pmatrix} m & -\frac{d}{dx} \\ \frac{d}{dx} & -m \end{pmatrix} \quad \text{in} \quad L^2((0,\infty), \mathbb{C}^2)$$
 (2)

with the domain consisting of functions $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \in H^1((0,\infty),\mathbb{C}^2)$ satisfying the one-parametric class of generalised infinite-mass boundary conditions

$$\psi_1(0)\cot(\alpha) = \psi_2(0) \quad \text{with} \quad \alpha \in \left(0, \frac{\pi}{2}\right).$$
 (3)

The latter restriction ensures that D_0 is self-adjoint and its spectrum is purely continuous:

$$\sigma(D_0) = \sigma_c(D_0) = (-\infty, -m] \cup [m, +\infty). \tag{4}$$

Given $V:(0,\infty)\to\mathbb{C}^{2,2}$, we consider the perturbed operator D_0+V . To be more specific, since we merely assume $V\in L^1\big((0,\infty),\mathbb{C}^{2,2}\big)$, we introduce a closed (not necessarily self-adjoint) extension D_V of the operator sum D_0+V via Kato's resolvent formula (see [22, 7] and below). For almost every $x\in(0,\infty)$, V(x) is a 2×2 matrix and we denote by |V(x)| its operator norm in \mathbb{C}^2 . Our main result reads as follows.

Theorem 1.1. Let $V \in L^1((0,\infty), \mathbb{C}^{2,2}; (1+x) dx)$ satisfy

$$\int_{0}^{\infty} \int_{0}^{\infty} |V(x)| \left[1 + \left(q + 2m \min(x, y) \right)^{2} \right] |V(y)| \, dx \, dy < 1 \,, \tag{5}$$

where $q := \max (\cot(\alpha), \cot(\alpha)^{-1})$. Then $\sigma(D_V) = \sigma_c(D_V) = \sigma(D_0)$.

The sufficient condition (5) follows as a consequence of a better (but less explicit) result which is of the same nature as (1) (see Theorem 3.3 and Remark 3.4). Moreover, the spectral stability follows as a consequence of the fact that D_V is similar to D_0 via bounded and boundedly invertible similarity transformation (see Theorem 3.3 and Corollary 3.5). Consequently, the possibly non-self-adjoint operator D_V is quasi-self-adjoint [25, 24].

We have not been able to prove the optimality of (5). To this purpose, we also establish an alternative sufficient condition for special potentials having V_{11} as the only non-zero component and this turns out to be optimal.

Our next goal is to relate Theorem 1.1 to the non-relativistic analogue given by the self-adjoint Laplacian (here m > 0)

$$S_0 := -\frac{1}{2m} \frac{d^2}{dx^2}$$
 in $L^2((0,\infty))$ (6)

with the domain consisting of functions $\psi \in H^2((0,\infty))$ satisfying the repulsive Robin boundary conditions

$$\psi'(0) = \beta\psi(0) \quad \text{with} \quad \beta \in (0, +\infty) . \tag{7}$$

The Dirichlet problem formally corresponds to $\beta = +\infty$. Given $V : (0, \infty) \to \mathbb{C}$ with $V \in L^1((0,\infty))$, the symbol S_V denotes the m-sectorial extension of the sum $S_0 + V$ obtained via the form sum. Spectral enclosures for S_V were established already in [17, 12]. However, the stability of the spectrum was not considered in these references and related results can be found only in more recent works [5, 23].

Theorem 1.2 ([23]). Let $V \in L^1((0, \infty), \mathbb{C}; (1+x) dx)$ satisfy

$$\int_{0}^{\infty} \int_{0}^{\infty} |V(x)| \left(\frac{2m}{\beta} + 2m \min(x, y)\right)^{2} |V(y)| \, dx \, dy < 1.$$
 (8)

Then $\sigma(S_V) = \sigma_c(S_V) = \sigma(S_0)$.

In this paper, we explain how to understand (8) as the non-relativistic limit of (5). We consistently introduce the speed of light c into the action of D_0 as well as to the boundary condition (3) and obtain S_0 as the limit (in a norm resolvent sense) of a suitably renormalised D_0 after sending c to $+\infty$. It turns out that the right correspondence between the relativistic and non-relativistic boundary conditions is given by

$$\beta = 2\cot\alpha. \tag{9}$$

Then (8) is a limit of a c-dependent version of (5) as $c \to +\infty$. From this perspective, the infinite-mass (also known as MIT) boundary condition given by $\alpha = \frac{\pi}{4}$ is a relativistic version of the Robin parameter $\beta = 2$.

We stress that our non-relativistic limit is in no mathematical contradiction with the other approaches [6, 1, 2], where the (generalised) infinite-mass boundary condition is interpreted as a relativistic version of the Dirichlet boundary condition (i.e., $\beta = +\infty$, irrespectively of the value of α). Indeed, the argument of the other authors is based on a different introduction of the speed of light into the operator. In particular, the boundary condition is left c-independent and that is why the dependence on α is lost in the limit, see [6, Prop. 3.1]. Our approach is based on physics arguments [28] recently adopted in a related problem in [20].

The paper is organised as follows. In Section 2 we write down the integral kernel of the resolvent of the unperturbed operator D_0 and establish two crucial uniform estimates (Lemmata 2.2 and 2.3). In Section 3 we introduce the perturbed operator D_V and establish various sufficient conditions which guarantee the similarity of D_V to D_0 . In particular, Theorem 1.1 (based on Lemma 2.2) is proved, but we also establish an alternative result, Theorem 3.6 (based on Lemma 2.3). In Section 4 we prove the optimality of Theorem 3.6 employing Dirac delta potentials. In Section 5, introducing the speed of light c in a refined way, we prove a convergence of D_0 to S_0 in a norm-resolvent sense as $c \to +\infty$. In particular, the compatibility of Theorems 1.1 and 1.2 is established.

2 The free Dirac operator and its resolvent

From now on, we abbreviate $\mathbb{R}_+ := (0, \infty)$ and consider the Hilbert space $L^2(\mathbb{R}_+, \mathbb{C}^2)$. The norm and inner product in this space (and other functional spaces) is denoted by $\|\cdot\|$ and (\cdot, \cdot) , respectively. The latter is assumed to be antilinear in its first argument. Recall that the absolute value notation $|\cdot|$ is maintained for the vector and operator norms of \mathbb{C}^2 .

The half-line Dirac operators (2), subject to the generalised infinite-mass boundary conditions (3), have already been considered in [6, 13]. We particularly follow the notation of the pioneering work of Cuenin [6], from where we overtake the following formula for the resolvent.

Proposition 2.1. Let $\alpha \in (0, \frac{\pi}{2})$. For $z \in \rho(D_0)$ the resolvent $(D_0 - z)^{-1}$ acts as an integral operator with the integral kernel

$$\mathcal{R}_{\alpha}(x,y;z) = \frac{1}{W} \left[\psi_{\alpha}(x;z) \,\phi_{\alpha}(y;z)^{\top} \,\Theta(x-y) + \phi_{\alpha}(x;z) \,\psi_{\alpha}(y;z)^{\top} \,\Theta(y-x) \right], \tag{10}$$

where Θ stands for the Heaviside function and

$$\psi_{\alpha}(x;z) := \exp(ik(z)x) \begin{pmatrix} i\zeta(z) \\ -1 \end{pmatrix},$$

$$\phi_{\alpha}(y;z) := \begin{pmatrix} \cos(k(z)y) + \zeta(z)\cot(\alpha)\sin(k(z)y) \\ -\zeta(z)^{-1}\sin(k(z)y) + \cot(\alpha)\cos(k(z)y) \end{pmatrix},$$

with coefficients

$$k(z):=\sqrt{z^2-m^2}, \qquad \zeta(z):=rac{z+m}{k(z)}, \qquad W:=1+i\zeta(z)\cot(\alpha).$$

The square root is chosen so that $\Im[k(z)] > 0$.

Proof. For the reader's convenience, we give a short proof of the resolvent formula. Moreover, the proof provides an insight into the structure, which will be useful when deriving the non-relativistic limit in Section 5.

Let us consider the whole-line Dirac operator D in $L^2(\mathbb{R}, \mathbb{C}^2)$, which acts as D_0 in (2), subject to continuity boundary conditions at zero, namely the domain of D is $H^1(\mathbb{R}, \mathbb{C}^2)$. Using the traditional trick $(D-z)^{-1} = (D+z)(D^2-z^2)^{-1}$ together with the fact that the resolvent kernel of the Schrödinger operator $D^2 = (-\Delta + m^2)I_{\mathbb{C}^2}$ is well known, one easily verifies that the integral kernel of $(D-z)^{-1}$ reads

$$\Re(x,y;z) = \frac{1}{2} \begin{pmatrix} i\zeta(z) & \operatorname{sgn}(x-y) \\ -\operatorname{sgn}(x-y) & i\zeta^{-1}(z) \end{pmatrix} \exp(ik(z)|x-y|). \tag{11}$$

Let $\phi \in L^2(\mathbb{R}_+, \mathbb{C}^2)$ be a function on the half-line and $\mathbb{A} := \operatorname{diag}(\mu_1, \mu_2) \in \mathbb{C}^{2,2}$ be a diagonal matrix. For every $x \in \mathbb{R}$, we define $\phi_{\mathbb{A}}(x) := \phi(|x|)\Theta(x) + \mathbb{A}\phi(|x|)\Theta(-x)$, which is a function in $L^2(\mathbb{R}, \mathbb{C}^2)$ but not necessarily in $H^1(\mathbb{R}, \mathbb{C}^2)$. Then

$$((D-z)^{-1}\phi_{\mathbb{A}})(x) = \int_{\mathbb{R}} \Re(x,y;z)\phi_{\mathbb{A}}(y)dy$$
$$= \int_{0}^{\infty} (\Re(x,y;z) + \Re(x,-y;z)\mathbb{A}) \ \phi(y)dy =: \begin{pmatrix} \xi_{1}(x) \\ \xi_{2}(x) \end{pmatrix}.$$

Demanding the boundary condition

$$\xi_1(0)\cot(\alpha) = \xi_2(0)$$

yields the equation

$$\int_0^\infty \left[\left(\Re_{11}(0,y;z) + \mu_1 \Re_{11}(0,-y;z) \right) \cot(\alpha) - \Re_{21}(0,y;z) - \mu_1 \Re_{21}(0,-y;z) \right] \phi_1(y) dy$$

$$= \int_0^\infty \left[\Re_{22}(0,y;z) + \mu_2 \Re_{22}(0,-y;z) - \left(\Re_{12}(0,y;z) + \mu_2 \Re_{12}(0,-y;z) \right) \cot(\alpha) \right] \phi_2(y) dy.$$

Since this equation has to be satisfied for every $\phi \in L^2(\mathbb{R}_+, \mathbb{C}^2)$, for the special choice $\phi_1 = 0$, respectively $\phi_2 = 0$, it breaks down into two independent conditions

$$(\mathcal{R}_{11}(0,y;z) + \mu_1 \mathcal{R}_{11}(0,-y;z)) \cot(\alpha) = \mathcal{R}_{21}(0,y;z) + \mu_1 \mathcal{R}_{21}(0,-y;z),$$

$$(\mathcal{R}_{12}(0,y;z) + \mu_2 \mathcal{R}_{12}(0,-y;z)) \cot(\alpha) = \mathcal{R}_{22}(0,y;z) + \mu_2 \mathcal{R}_{22}(0,-y;z),$$

to be satisfied for all $y \in \mathbb{R}_+$. From the expression (11) we obtain a unique solution for μ_1, μ_2 leading to

$$\mathbb{A} = \sigma_3 \eta(\alpha) \quad \text{with} \quad \eta(\alpha) := \frac{1 - i\zeta(z)\cot(\alpha)}{1 + i\zeta(z)\cot(\alpha)} \quad \text{and} \quad \sigma_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The integral kernel of the resolvent $(D_0 - z)^{-1}$ therefore reads

$$\Re_{\alpha}(x, y; z) = \Re(x, y; z) + \Re(x, -y; z)\sigma_{3}\eta(\alpha), \tag{12}$$

for all $x, y \in \mathbb{R}_+$. It is straightforward to check that

$$\mathcal{R}_{\alpha}(x,y;z) = R_{\alpha}(x,y;z)\Theta(x-y) + R_{\alpha}(y,x;z)^{\top}\Theta(y-x)$$

with

$$[R_{\alpha}(x,y;z)]_{11} = \frac{i\zeta(z)}{1 + i\zeta(z)\cot(\alpha)} \exp(ik(z)x) \left[\cos(k(z)y) + \zeta(z)\cot(\alpha)\sin(k(z)y)\right],$$

$$[R_{\alpha}(x,y;z)]_{12} = \frac{i\zeta(z)}{1 + i\zeta(z)\cot(\alpha)} \exp(ik(z)x) \left[\cot(\alpha)\cos(k(z)y) - \zeta(z)^{-1}\sin(k(z)y)\right],$$

$$[R_{\alpha}(x,y;z)]_{21} = \frac{-1}{1 + i\zeta(z)\cot(\alpha)} \exp(ik(z)x) \left[\cos(k(z)y) + \zeta(z)\cot(\alpha)\sin(k(z)y)\right],$$

$$[R_{\alpha}(x,y;z)]_{22} = \frac{-1}{1 + i\zeta(z)\cot(\alpha)} \exp(ik(z)x) \left[\cot(\alpha)\cos(k(z)y) - \zeta(z)^{-1}\sin(k(z)y)\right].$$

This coincides with (10).

The following lemma is the main technical result of this paper. Here it is fundamental that the boundary parameter α is assumed to belong to the interval $(0, \frac{\pi}{2})$, cf. (3). Indeed, D_0 possesses a discrete eigenvalue in the interval (-m, m) if $\alpha \in (\frac{\pi}{2}, \pi)$ and $\Re_{\alpha}(x, y; z)$ has a non-removable singularity at $z = \pm m$ even if $\alpha \in \{0, \frac{\pi}{2}\}$ (zigzag boundary conditions). Recall the notation q introduced in Theorem 1.1.

Lemma 2.2. Let $\alpha \in (0, \frac{\pi}{2})$. Then

$$\sup_{z \in \rho(D_0)} |\mathcal{R}_{\alpha}(x, y; z)|^2 = 1 + (q + 2m \min(x, y))^2.$$

Proof. The norm of the resolvent kernel $\mathcal{R}_{\alpha}(x,y;z)$ as an operator $\mathbb{C}^2 \to \mathbb{C}^2$ reads

$$|\mathcal{R}_{\alpha}(x,y;z)| = \frac{1}{|W(z)|} \left[|\psi_{\alpha}(x;z)| |\phi_{\alpha}(y;z)| \Theta(x-y) + |\psi_{\alpha}(y;z)| |\phi_{\alpha}(x;z)| \Theta(y-x) \right].$$

Without loss of generality, we shall further consider only the case x > y. To simplify further expressions, let us introduce the notation

$$\eta_1(z) := \frac{1}{4} \left(1 + |\zeta(z)|^{-2} \right) |1 - i\zeta(z) \cot(\alpha)|^2,
\eta_2(z) := \frac{1}{4} \left(1 + |\zeta(z)|^{-2} \right) |1 + i\zeta(z) \cot(\alpha)|^2,
\eta_3(z) := \frac{1}{2} \left(1 - \cot^2(\alpha) |\zeta(z)|^2 \right),
\eta_4(z) := \frac{1}{2} \left(2\Re[\zeta(z)] \cot(\alpha) \right).$$

A straightforward calculation yields

$$\begin{aligned} |\psi_{\alpha}(x;z)|^2 &= \exp(-2\Im[k(z)]x) \left(|\zeta(z)|^2 + 1 \right), \\ |\phi_{\alpha}(y;z)|^2 &= \eta_1(z) \exp(-2\Im[k(z)]y) + \eta_2(z) \exp(2\Im[k(z)]y) \\ &+ \left(1 - |\zeta(z)|^{-2} \right) \left(\eta_3(z) \cos(2\Re[k(z)]y) + \eta_4(z) \sin(2\Re[k(z)]y) \right). \end{aligned}$$

Since the sums, products, and compositions of holomorphic functions are holomorphic, it is obvious that the resolved kernel $\mathcal{R}_{\alpha}(x,y;z)$ is a holomorphic function of the spectral parameter z. Therefore, the supremum of its modulus cannot be achieved in the resolvent set. This fact follows from the maximum modulus principle stated in [8] for general Banach space-valued maps. The fact that \mathbb{C}^2 equipped with the standard norm satisfies the appropriate assumptions follows from [9].

On the other hand, the supremum itself has to exist. Therefore, either the supremum lies at complex infinity or it is achieved somewhere in the spectrum (4). We shall show that the latter is true. First, we show that $|\mathcal{R}_{\alpha}(x,y;z)|$ can be uniformly bounded as $z \to \infty$.

Estimating the exponential functions one has

$$\lim_{z \to \infty} |\mathcal{R}_{\alpha}(x, y; z)|^{2} \Theta(x - y) = \lim_{z \to \infty} \frac{1}{|W|^{2}} |\psi_{\alpha}(x; z)|^{2} |\phi_{\alpha}(y; z)|^{2} \Theta(x - y)$$

$$\leq \lim_{z \to \infty} \frac{(1 + |\zeta(z)|^{2})}{|W|^{2}} (\eta_{1}(z) + \eta_{2}(z))$$

$$+ \lim_{z \to \infty} \frac{(1 - |\zeta(z)|^{-2})}{|W|^{2}} (\cos(2\Re[k(z)]y)\eta_{3}(z) + \sin(2\Re[k(z)]y)\eta_{4}(z))$$

$$= 2$$

The last equality follows from the facts that $|1 \pm i\zeta(z)\cot(\alpha)|^2 \to 1 + \cot^2(\alpha)$ and $|\zeta(z)|^2 \to 1$ as $z \to \infty$, which is a consequence of the asymptotics of k(z) and $\zeta(z)$. Indeed, it can be easily checked that, under the notation $z = z_1 + iz_2$ with $z_1, z_2 \in \mathbb{R}$, it holds that

$$k(z) = \frac{z_1 z_2}{a(z)} + ia(z),$$

with

$$a(z) := \frac{1}{\sqrt{2}} \left(z_2^2 - z_1^2 + m^2 + \sqrt{(z_2^2 - z_1^2 + m^2)^2 + 4z_1^2 z_2^2} \right)^{1/2}.$$

Furthermore, for $z_2 \neq 0$, by simple calculations one finds that,

- $\lim_{|z_2| \to +\infty} a(z) = +\infty$ as $o(|z_2|)$,
- $\bullet \lim_{|z_1| \to +\infty} a(z) = |z_2|,$
- $\lim_{|z_1| \to +\infty} \Im(\zeta(z)) = \lim_{|z_2| \to +\infty} \Im(\zeta(z)) = 0.$

Now, we investigate the behaviour of the restriction $|\mathcal{R}_{\alpha}(x,y;z)|^2$ to the spectrum. That is, $u := z \in \sigma(D_0) = (-\infty, -m] \cup [m, +\infty)$. The coefficients k(u) and $\zeta(u)$ are now purely real, and the restriction of $|\mathcal{R}_{\alpha}(x,y;z)|^2$ reads

$$|\mathcal{R}_{\alpha}(x,y;u)|^{2}\Theta(x-y) = \frac{2u^{2}}{u^{2}-m^{2}} + \cos(2k(u)y)\frac{u-m-\cot^{2}(\alpha)(u+m)}{u-m+\cot^{2}(\alpha)(u+m)}\frac{2mu}{u^{2}-m^{2}} + \sin(2k(u)y)\frac{\cot(\alpha)4m}{\sqrt{u^{2}-m^{2}}}\frac{1}{u-m+\cot^{2}(\alpha)(u+m)} =: \chi(u,y).$$

We claim that the function $\chi(u,y)$ has no local extremes for $u \in \sigma(\mathcal{D}_{\alpha})$. Indeed, the derivative

$$\frac{\partial \chi}{\partial u}(u,y) = \xi_1(u) + \xi_2(u)\cos(2k(u)y) + \xi_3(u)\cos(2k(u)y)y
+ \xi_4(u)\sin(2k(u)y) + \xi_5(u)\sin(2k(u)y)y,$$
(13)

with some functions ξ_1, \ldots, ξ_5 that we do not need to make explicit now, can be viewed as the linear combination of linearly independent functions in variable y for every fixed $u \in \sigma(D_0)$. Therefore, the only possible solution of the equation when the derivative is put equal to zero is $\xi_i = 0$ for every $i \in \{1, \ldots, 5\}$. Since

$$\xi_1(u) = \frac{4mu}{u^2 - m^2},$$

we conclude that $u = 0 \notin (-\infty, -m] \cup [m, +\infty)$ if m > 0 or that $\xi_1(u) = 0$ cannot be satisfied if m = 0.

Since we have established that $u \mapsto \chi(u, y)$ has no local extremes, its supremum must the maximum of the limits at the "boundaries". These are

$$\lim_{u \to \pm \infty} \chi(u, y) = 2,$$

$$\lim_{u \to -m} \chi(u, y) = 1 + \left(\cot(\alpha) + 2my\right)^2,$$

$$\lim_{u \to m} \chi(u, y) = 1 + \left(\cot(\alpha)^{-1} + 2my\right)^2.$$

Thus, we conclude that the claim of the lemma holds true.

As an alternative to Lemma 2.2, we shall use the following uniform bound on the first component of the resolvent kernel.

Lemma 2.3. Let $\alpha \in (0, \frac{\pi}{2})$. Then

$$\sup_{z \in \rho(D_0)} |\mathcal{R}_{\alpha,11}(x,y;z)| = \max\left(1, \frac{1}{\cot(\alpha)} + 2m\min(x,y)\right).$$

Proof. It is not difficult to see that

$$|\mathcal{R}_{\alpha,11}(x,y;z)|^2 \Theta(x-y) = \frac{|\zeta(z)|^2 \exp(-2\Im(k(z))x)}{|1+i\zeta(z)\cot(\alpha)|^2} \left(\Lambda_1(y;z) + \Lambda_2(y;z)\right),$$

where for x > y > 0 we have

$$\begin{split} & \Lambda_1(x,y;z) := \frac{1}{4} \left[\exp(-2\Im(k(z))y) |1 - i\zeta(z)\cot(\alpha)|^2 + \exp(2\Im(k(z))y) |1 + i\zeta(z)\cot(\alpha)|^2 \right], \\ & \Lambda_2(x,y;z) := \frac{1}{2} \left[\cos(2\Re(k(z))y) \left(1 - |\zeta(z)|^2 \cot^2(\alpha) \right) + \sin(2\Re(k(z))y) 2\cot(\alpha)\Re(\zeta(z)) \right]. \end{split}$$

Adopting the approach in the general setting of Lemma 2.2, we see that

$$\sup_{z \in \rho(D_0)} |\mathcal{R}_{\alpha,11}(x,y;z)| = \max(l_\infty, l_{\pm \infty}, l_{\pm m}),$$

where we denote

$$l_{\infty} := \limsup_{z \to \infty} |\mathcal{R}_{\alpha,11}(x,y;z)|^2 \Theta(x-y),$$

$$l_{\pm \infty} := \limsup_{z \to \pm \infty} |\mathcal{R}_{\alpha,11}(x,y;z)|^2 \Theta(x-y),$$

$$l_{\pm m} := \lim_{z \to \pm m} |\mathcal{R}_{\alpha,11}(x,y;z)|^2 \Theta(x-y).$$

In the present case, the problem is more delicate, since the alternating part Λ_2 does not vanish as z tends to a complex infinity. However, in general for any $\nu, \mu \in \mathbb{R}$ it holds that

$$\lim_{x \to +\infty} \sup (\nu \cos(x) + \mu \sin(x)) = \sqrt{\nu^2 + \mu^2}.$$
 (14)

Calculation of l_{∞} . The fact that z tends to a complex infinity means that either $|\Im(z)| \to +\infty$ or $|\Re(z)| \to +\infty$. The only relevant case is the latter one, since $|\Re_{\alpha,11}(x,y;z)|^2\Theta(x-y) \to 0$ whenever $|\Im(z)|$ tends to $+\infty$, see the asymptotics of k(z) and $\zeta(z)$ in the proof of Lemma 2.2. Combining (14) and the fact pointed out above, one can see that for any curve $\gamma: \mathbb{R}_+ \to \rho(D_0)$ such that $\lim_{t\to +\infty} |\Re(\gamma(t))| = +\infty$ we have

$$\limsup_{t \to +\infty} |\mathcal{R}_{\alpha,11}(x,y;\gamma(t))|^2 \Theta(x-y) \le \frac{\exp(-2\gamma_{\infty}^I x)}{2} \left(\cosh(2\gamma_{\infty}^I y) + 1\right) \le 1,$$

where $\gamma_{\infty}^{I} := \liminf_{t \to \infty} |\Im(\gamma(t))|$ which is, without loss of generality, assumed to be finite. Thus we conclude with $l_{\infty} \leq 1$.

Calculation of $l_{\pm\infty}$. Restricted to the spectrum, i.e. $z = u \in (-\infty, -m] \cup [m, +\infty)$, the absolute value of $\mathcal{R}_{\alpha,11}(x,y;u)$ reads

$$|\mathcal{R}_{\alpha,11}(x,y;u)|^2 \Theta(x-y) = \frac{1}{2} \frac{u+m}{u-m} + \cos(2k(u)y) \frac{1}{2} \frac{u-m-\cot^2(\alpha)(u+m)}{u-m+\cot^2(\alpha)(u+m)} \frac{u+m}{u-m} + \sin(2k(u)y) \frac{\cot(\alpha)(u+m)}{\sqrt{u^2-m^2}} \frac{1}{u-m+\cot^2(\alpha)(u+m)}.$$

Once again, from (14) it follows that

$$l_{\pm\infty} = \limsup_{u \to \pm \infty} |\Re_{\alpha}(x, y; u)|^2 \Theta(x - y) = 1.$$

Limits at $\pm m$. By a direct calculation we obtain

$$l_{-m} = \lim_{u \to -m} |\mathcal{R}_{\alpha}(x, y; u)|^2 \Theta(x - y) = 0,$$

$$l_{m} = \lim_{u \to m} |\mathcal{R}_{\alpha}(x, y; u)|^2 \Theta(x - y) = \left(\frac{1}{\cot(\alpha)} + 2my\right)^2.$$

The proof is completed by comparing the values of $l_{\infty}, l_{\pm \infty}, l_{\pm m}$.

3 Perturbations and the stability of the spectrum

Given a potential $V: \mathbb{R}_+ \to \mathbb{C}^{2,2}$, we would like to consider a closed realisation D_V of the perturbed operator $D_0 + V$. Already the latter is a closed operator, with $\text{dom}(D_0 + V) = \text{dom}(D_0)$, provided that V is bounded or, more generally, relatively bounded with respect to D_0 with the relative bound less than 1. This results in the requirement $V \in (L^2 + L^{\infty})(\mathbb{R}_+, \mathbb{C}^{2,2})$.

To include a larger class of potentials, it is possible to introduce D_V via the pseudo-Friedrichs extension [19]. This approach leads to $V \in (L^{1+\delta} + L^{\infty})(\mathbb{R}_+, \mathbb{C}^{2,2})$ with any positive δ . The advantage is a quite explicit form-wise interpretation of D_V and the strong apparatus of the Birman–Schwinger principle developed in [19]. However, the desired setting $\delta = 0$ does not seem to be reachable.

To reach the optimal scale $V \in (L^1 + L^{\infty})(\mathbb{R}_+, \mathbb{C}^{2,2})$, it seems necessary to introduce D_V via Kato's resolvent formula [22, 7]. We adopt this approach in this paper. The abstract Kato's theorem, in a weaker form sufficient to our purposes, reads as follows.

Theorem 3.1 ([22]). Let H_0 be a self-adjoint operator in a Hilbert space \mathcal{H} and suppose that A, B are closed operators in \mathcal{H} with $dom(H_0) \subset dom(A) \cap dom(B)$ which are H_0 -smooth and such that

$$\sup_{z \in \rho(H_0)} ||A(H_0 - z)^{-1}B^*|| < 1.$$
(15)

Then there exists a closed extension H_V of $H_0 + B^*A$ which is similar to H_0 satisfying for all $z \in \rho(H_0)$

$$(H_0 - z)^{-1} - (H_V - z)^{-1} = \overline{(H_0 - z)^{-1}B^*}A(H_V - z)^{-1}.$$

Here the similarity means that there exists an operator $W \in \mathcal{B}(\mathcal{H})$ such that $W^{-1} \in \mathcal{B}(\mathcal{H})$ and $H_V = WH_0W^{-1}$ holds. Consequently, the operators H_V and H_0 are isospectral. What is more, also the nature of the spectrum is preserved, in particular $\sigma_c(H_V) = \sigma_c(H_0)$. (The continuous spectrum $\sigma_c(H)$ of any closed operator H in a Hilbert space \mathcal{H} is defined as the set of all complex points λ which are not eigenvalues of H such that the range of $H - \lambda$ is not equal to \mathcal{H} but its closure is.)

Instead of giving the very definition of H_0 -smoothness [22, Def. 1.2], we use the equivalent criterion suitable to our purposes [22, Thm. 5.1] which says that A is H_0 -smooth if, and only if,

$$\sup_{z \in \rho(H_0), \, \psi \in \text{dom}(A^*) \setminus \{0\}} \frac{|(A^*\psi, [R_0(z) - R_0(z^*)] \, A^*\psi)|}{\|\psi\|^2} < \infty \tag{16}$$

where $R_0(z) := (H_0 - z)^{-1}$.

In our case, we set $H_0 := D_0$ and $\mathcal{H} := L^2(\mathbb{R}_+, \mathbb{C}^2)$ and use the matrix polar decomposition $V(x) = U(x)(V(x)^*V(x))^{1/2}$ with U(x) unitary to set $A := (V^*V)^{1/4}$ and $B^* := U(V^*V)^{1/4}$ as maximal operators of matrix multiplication in \mathcal{H} . One has $B := (V^*V)^{1/4}U^*$ and $\mathrm{dom}(A) = \mathrm{dom}(B) = \mathrm{dom}(V^*V)^{1/4}$.

We proceed with verifying the hypotheses of Theorem 3.1. Since the domain of D_0 is a subspace of $H^1(\mathbb{R}_+, \mathbb{C}^2)$, the following proposition ensures the right inclusion properties between domains of A, B and D_0 .

Proposition 3.2. If $V \in L^1(\mathbb{R}_+, \mathbb{C}^{2,2})$, then $H^1(\mathbb{R}_+, \mathbb{C}^2) \subset \text{dom}(A)$.

Proof. Let $\psi \in C_0^{\infty}(\mathbb{R}, \mathbb{C}^2)$ and $\varepsilon > 0$. Then we have

$$|\psi(x)|^2 = \int_{\infty}^{x} |\psi|^{2'} = 2 \int_{\infty}^{x} \Re(\bar{\psi}^{\top}\psi') \le 2 \|\psi\| \|\psi'\| \le \varepsilon \|\psi'\|^2 + \varepsilon^{-1} \|\psi\|^2,$$

for all $x \in \mathbb{R}_+$ and therefore also $\|\psi\|_{\infty}^2 \le \varepsilon \|\psi'\|^2 + \varepsilon^{-1} \|\psi\|^2$, where $\|\psi\|_{\infty} := \||\psi|\|_{L^{\infty}(\mathbb{R}_+)}$. Consequently,

$$||A\psi||^2 = \left(\psi, (V^*V)^{1/2}\psi\right) \le \int_{\mathbb{R}_+} |V(x)| \, |\psi(x)|^2 \, dx \le ||V||_1 \, ||\psi||_{\infty}^2 \le ||V||_1 \, \left(\varepsilon \, ||\psi'||^2 + \varepsilon^{-1} \, ||\psi||^2\right),$$

where $||V||_1 := |||V|||_{L^1(\mathbb{R}_+)}$. In the second inequality, we have used the equality of the operator norms $|(V(x)^*V(x))^{1/2}| = |V(x)|$ in \mathbb{C}^2 . The density of restrictions of $C_0^{\infty}(\mathbb{R}, \mathbb{C}^2)$ to \mathbb{R}_+ in $H^1(\mathbb{R}_+, \mathbb{C}^2)$ concludes the proof.

To prove (15) with $R_0 := (D_0 - z)^{-1}$, we employ the fact that $K_z := AR_0(z)B^*$ is an integral operator in $\mathcal{H} = L^2(\mathbb{R}_+, \mathbb{C}^2)$ with the kernel (denoted by the same letter)

$$K_z(x,y) := |V(x)^*V(x)|^{1/4} \, \mathcal{R}_{\alpha}(x,y;z) \, U(y) \, |V(y)^*V(y)|^{1/4} \, .$$

The operator norm in \mathbb{C}^2 of the latter can be estimated as follows:

$$|K_z(x,y)| \le |V(x)|^{1/2} |\Re_{\alpha}(x,y;z)| |V(y)|^{1/2} =: L_z(x,y).$$
 (17)

The integral operator L_z in $L^2(\mathbb{R}_+,\mathbb{C})$ generated by the kernel $L_z(x,y)$ obviously satisfies

$$||K_z|| \leq ||L_z||$$

(the norms are determined by the spaces which the operators act in).

What is more, also the D_0 -smoothness of A and B is reduced to the uniform boundedness of L_z . This follows from the triangle inequality applied to (16) and estimating the individual terms as in (17).

Finally, Lemma 2.2 yields $||L_z|| \le ||L||$, where L is the integral operator in $L^2(\mathbb{R}_+, \mathbb{C})$ generated by the z-independent kernel

$$L(x,y) := |V(x)|^{1/2} \sqrt{1 + (q + 2m\min(x,y))^2} |V(y)|^{1/2}.$$
 (18)

Applying Theorem 3.1, we therefore conclude with the following theorem, which is the central result of this paper.

Theorem 3.3. Let $V \in L^1(\mathbb{R}_+, \mathbb{C}^{2,2})$ satisfy ||L|| < 1. Then D_V is similar to D_0 .

Remark 3.4. The sufficient condition of Theorem 3.3 is of the same nature as the subordination requirement (1). To see it, note (cf. [23, Rem. 21]) that (8) implies that the non-relativistic variant

$$L_{\infty}(x,y) := |V(x)|^{1/2} \left(\frac{2m}{\beta} + 2m \min(x,y) \right) |V(y)|^{1/2}$$

as an integral operator in $L^2(\mathbb{R}_+,\mathbb{C})$ has norm less than one. Since $L_{\infty} = |V|^{1/2} S_0^{-1} |V|^{1/2}$, the last requirement is equivalent to the subordination condition

$$\forall \psi \in W^{1,2}(\mathbb{R}_+), \qquad \int_0^\infty |V| |\psi|^2 \le \frac{a}{2m} \left(\int_0^\infty |\psi'|^2 + \beta |\psi(0)|^2 \right),$$

with a < 1. Note that the right-hand side is the quadratic form of S_0 multiplied by a. We refer to [15, Sec. 2] and [19, Sec. 7.1.1] for the idea of the equivalence. Now, $L(x,y) \le L_1(x,y) + L_2(x,y)$, where $L_1(x,y) := |V(x)|^{1/2} |V(y)|^{1/2}$ is a rank-one operator and

$$L_2(x,y) := |V(x)|^{1/2} (q + 2m \min(x,y)) |V(y)|^{1/2}$$

has the same structure as L_{∞} . Consequently, ||L|| < 1 provided that $||L_1|| \le 1 - a$ and $||L_2|| \le a$ with some $a \in (0,1)$. This is equivalent to the simultaneous validity of $||V||_1 \le 1 - a$ and

$$\forall \psi \in W^{1,2}(\mathbb{R}_+), \qquad \int_0^\infty |V| \, |\psi|^2 \le \frac{a}{2m} \left(\int_0^\infty |\psi'|^2 + \frac{2m}{q} \, |\psi(0)|^2 \right).$$

Estimating the operator norm of L by the Hilbert–Schmidt norm, we get the following corollary of Theorem 3.3, which particularly implies the spectral claim of Theorem 1.1 from the introduction.

Corollary 3.5. Let $V \in L^1(\mathbb{R}_+, \mathbb{C}^{2,2}; (1+x) dx)$ satisfy (5). Then D_V is similar to D_0 .

The condition $V \in L^1(\mathbb{R}_+, \mathbb{C}^{2,2}; (1+x) dx)$ ensures the finiteness of the integral in (5). Indeed, using $\min(x, y) \leq \sqrt{xy}$ for every $x, y \in \mathbb{R}_+$, condition (5) follows as a consequence of

$$\left(\int_0^\infty |V(x)| \, dx\right)^2 + \left(\int_0^\infty |V(x)| \left(q + 2mx\right)^2 dx\right)^2 < 1.$$

Finally, we state the following alternative result to Corollary 3.5 for very particularly structured potentials. Its proof follows the same chain of reasonings as above, just the usage of Lemma 2.2 is replaced by Lemma 2.3.

Theorem 3.6. Let $V \in L^1((0,\infty), \mathbb{C}^{2,2}; (1+x) dx)$ be such that $V_{12} = V_{21} = V_{22} = 0$ and satisfy

$$\int_{0}^{\infty} \int_{0}^{\infty} |V(x)| \, \max\left(1, \frac{1}{\cot(\alpha)} + 2m\min(x, y)\right)^{2} \, |V(y)| \, dx \, dy < 1. \tag{19}$$

Then D_V is similar to D_0 .

4 The Dirac delta potentials and optimality

It is natural to ask the question whether our sufficient condition (5) is optimal. More specifically, by the optimality we mean that for every $\varepsilon > 0$ there is a potential $V \in L^1(\mathbb{R}_+, \mathbb{C}^{2,2}; (1+x) dx)$

$$||V||_{m}^{2} := \int_{0}^{\infty} \int_{0}^{\infty} |V(x)| \left[1 + \left(q + 2m \min(x, y) \right)^{2} \right] |V(y)| \, dx \, dy = 1 + \varepsilon \,,$$

and D_V has an eigenvalue outside the interval $\sigma(D_0) = (-\infty, -m] \cup [m, +\infty)$.

In the case of Schrödinger operators on the half-axis [17] as well as Dirac operators on the whole line [7], an optimality is always achieved by Dirac delta potentials. We try to follow this approach in this paper too.

We restrict to m > 0. Given $t \in \mathbb{R}$ and a > 0, let V_t be formally given by

$$V_t(x) := t \, \delta(x - a) \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \tag{20}$$

Then $||V||_m^2 = t^2 [1 + (q + 2ma)^2]$, so the question is whether D_{V_t} has an eigenvalue in (-m, m) whenever $t > t_*$ or $t < -t_*$ with

$$t_* := \frac{1}{\sqrt{1 + (q + 2ma)^2}}.$$

Rigorously, we understand D_{V_t} as the self-adjoint operator which acts as D_0 in $(0, a) \cup (a, +\infty)$ and functions $\psi \in H^1(\mathbb{R}_+, \mathbb{C}^2)$ in its domain satisfy, in addition to the boundary condition (3), also the interface condition (see, e.g., [20])

$$\begin{pmatrix} it/2 & -i \\ 1 & 0 \end{pmatrix} \psi(a^+) + \begin{pmatrix} it/2 & i \\ -1 & 0 \end{pmatrix} \psi(a^-) = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Then we rely on the property that D_{V_t} can be approximated by a family $D_0 + \epsilon^{-1}V(\epsilon^{-1}\cdot)$ with regular potentials V in a norm-resolvent sense, cf. [29] (the adaptation of the proof of [29] to the present model is straightforward, because the resolvent kernel of our operator D_0 on the half-line is derived from its whole-line realisation D).

Solving the eigenvalue problem $D_{V_t}\psi = \lambda \psi$ with $\lambda \in (-m, m)$, we arrive at the implicit equation

$$t = -\frac{\cot(\alpha)\sqrt{m^2 - \lambda^2} + m - \lambda}{\cot(\alpha)(m + \lambda)\sinh(a\sqrt{m^2 - \lambda^2}) + \sqrt{m^2 - \lambda^2}\cosh(a\sqrt{m^2 - \lambda^2})} e^{a\sqrt{m^2 - \lambda^2}}.$$
 (21)

Since the right-hand side is negative for any $\lambda \in (-m, m)$, we get the expected result that there no solutions for all $t \geq t_0$ with some $t_0 < 0$. On the other hand, $\lambda = 0$ is clearly a solution of (21) for a certain negative value of t. Hence, the critical value t_0 can be chosen such that D_{V_t} possesses a discrete eigenvalue for all t less than but close to t_0 . This eigenvalue emerges for $t = t_0$ either from m or -m. Since the right-hand side of (21) diverges as $\lambda \to -m$, the eigenvalue must emerge from m. Taking the limit $\lambda \to m$ in (21), we find

$$t_0 = -\frac{\cot(\alpha)}{1 + 2ma\cot(\alpha)}. (22)$$

The dependence of the eigenvalue on the parameter t is depicted in Figure 1.




Figure 1: The eigenvalue of D_{V_t} as a function of t for m=1, a=2 and $\alpha=\frac{\pi}{4}$.

Since $t_0 < -t_*$, we conclude that Theorem 1.1 is not optimal for the choice (20). We also tried different matrices with different entries or combinations being the Dirac delta interactions,

but never achieved the optimality. We leave as an open problem whether the optimality of Theorem 1.1 can be achieved by a different choice of the potential.

Despite failing to establish the optimality of Theorem 1.1, the present analysis proves the optimality of the alternative result stated in Theorem 3.6.

Proposition 4.1. Theorem 3.6 is optimal for the choice (20). More specifically, let $\alpha \in (0, \frac{\pi}{2})$, m > 0 and a > 0 be such that

 $\frac{1}{\cot(\alpha)} + 2ma > 1.$

Then there exists $\varepsilon_0 > 0$ such that, given any $\varepsilon \in (0, \varepsilon_0)$, the operator D_{V_t} with $t = t_0 \sqrt{1 + \varepsilon}$ possesses a discrete eigenvalue in the interval (-m, m) and the left-hand side of (19) equals $1 + \varepsilon$.

5 The non-relativistic limit

In order to calculate the non-relativistic limit, the speed of light c, as a positive parameter to be sent to infinity, needs to be recovered in D_0 . As for the action of D_0 , we follow the classical literature [28] and conventionally set

$$D_{0,c} := \begin{pmatrix} mc^2 & -c\frac{d}{dx} \\ c\frac{d}{dx} & -mc^2 \end{pmatrix}.$$

Contrary to the recent approaches in [6, 1, 2], however, we also make the boundary condition c-dependent (and in fact also m-dependent):

$$dom(D_{0,c}) := \left\{ \psi \in H^1(\mathbb{R}_+, \mathbb{C}^2) : \ \psi_1(0) \frac{\cot(\alpha)}{mc} = \psi_2(0) \right\}.$$

This particular form is motivated by the fact that the boundary condition can be considered as a point interaction, which is known to be approximated by regular potentials [29], and the latter require a c-dependence, see [28, Chap. 6], [3] and [20].

Throughout this section we assume m > 0. Physically, it makes sense because there is no non-relativistic quantum theory for massless particles. Mathematically, the non-relativistic Hamiltonian (6), which we expect to be obtained after sending c to infinity, requires positive m.

The sufficient condition (5) restated for $D_{0,c}$ perturbed by c-independent potentials V reads

$$\frac{1}{c^2} \int_0^\infty \int_0^\infty |V(x)| \left[1 + \left(q_c + 2mc \min(x, y) \right)^2 \right] |V(y)| \, dx \, dy < 1 \,,$$

where $q_c := \max\left(\frac{\cot(\alpha)}{mc}, \frac{mc}{\cot(\alpha)}\right)$. Sending the c to infinity, we formally (here V is a matrix-valued function) arrive at the sufficient condition (8), which guarantees the stability of the spectrum of the non-relativistic operator S_V under the identification (9). Let us now prove that not only the sufficient conditions (5) and (8) are compatible in the non-relativistic limit, but the operators do so as well.

Theorem 5.1. Let $\alpha \in (0, \frac{\pi}{2})$ and m > 0. If $z \in \rho(S_0)$, then, for all sufficiently large c, $z \in \rho(D_{0,c} - mc^2)$ and

$$\lim_{c \to +\infty} \left\| (D_{0,c} - mc^2 - z)^{-1} - \begin{pmatrix} (S_0 - z)^{-1} & 0 \\ 0 & 0 \end{pmatrix} \right\| = 0.$$

Proof. In the present c-dependent setting, the integral kernel of $(D_{0,c}-z)^{-1}$ reads (cf. (12))

$$\mathcal{R}_{\alpha,c}(x,y;z) = \mathcal{R}_c(x,y;z) + \mathcal{R}_c(x,-y;z)\sigma_3\eta_c(\alpha) \quad \text{with} \quad \eta_c(\alpha) := \frac{1 - i\zeta_c(z)\frac{\cot(\alpha)}{mc}}{1 + i\zeta_c(z)\frac{\cot(\alpha)}{mc}},$$

where $z \in \rho(D_{0,c}) = \mathbb{C} \setminus (-\infty, -mc^2] \cup [mc^2, +\infty)$ and

$$\mathcal{R}_c(x,y;z) = \frac{1}{2c} \begin{pmatrix} i\zeta_c(z) & \operatorname{sgn}(x-y) \\ -\operatorname{sgn}(x-y) & i\zeta_c^{-1}(z) \end{pmatrix} \exp(ik_c(z)|x-y|),$$

is the resolvent kernel of D_c , the free c-dependent variant of the Dirac operator D acting on the whole line, with

$$\zeta_c(z) := \frac{z + mc^2}{ck_c(z)}$$
 and $k_c(z) := c^{-1}\sqrt{z^2 - (mc^2)^2}$.

On the other hand, for $z \in \rho(S_0) = \mathbb{C} \setminus [0, +\infty)$, the resolvent kernel of S_0 is given by (cf. [23])

$$\mathcal{G}_{\alpha}(x,y;z) := (S_0 - z)^{-1}(x,y;z) = \mathcal{G}(x,y;z) + \mathcal{G}(x,-y;z)\,\xi(\alpha),$$

where

$$\xi(\alpha) := \frac{\sqrt{2mz} - i2\cot(\alpha)}{\sqrt{2mz} + i2\cot(\alpha)}$$
(23)

represents the boundary condition and

$$\mathfrak{G}(x,y;z) := \frac{im}{\sqrt{2mz}} \exp(i\sqrt{2mz}|x-y|),$$

is the resolvent kernel of the whole-line Schrödinger operator S in $L^2(\mathbb{R}, \mathbb{C})$, which acts as the Schrödinger operator S_0 in (6), subject to continuity boundary conditions at zero up to the first derivative, namely the domain of S is $H^2(\mathbb{R}, \mathbb{C})$.

The proof of the theorem follows by comparing the integral operators generated by the kernels $\mathcal{R}_{\alpha,c}$ and \mathcal{G}_{α} . We essentially employ the well-known relativistic limit of D_c to S on the whole line. To this purpose, we denote by $\|\cdot\|_{\mathbb{R}_+}$ and $\|\cdot\|_{\mathbb{R}}$ (operator) norms on $L^2(\mathbb{R}_+,\mathbb{C}^2)$ and $L^2(\mathbb{R},\mathbb{C}^2)$, respectively.

Given any $\phi \in L^2(\mathbb{R}_+, \mathbb{C}^2)$, we introduce

$$\phi_{\mathbb{A}_c}(x) := \phi(|x|)\Theta(x) + \mathbb{A}_c\phi(|x|)\Theta(-x),$$

with $\mathbb{A}_c := \sigma_3 \eta_c(\alpha)$. Note that

$$\|\phi_{\mathbb{A}_c}\|_{\mathbb{R}}^2 = \int_{\mathbb{R}} |\phi_{\mathbb{A}_c}(x)|^2 dx = \int_{\mathbb{R}_+} |\phi(x)|^2 dx + \int_{\mathbb{R}_+} |\mathbb{A}_c \phi(x)|^2 dx$$
$$\leq (1 + |\eta_c(\alpha)|^2) \int_{\mathbb{R}_+} |\phi(x)|^2 dx = (1 + |\eta_c(\alpha)|^2) \|\phi\|_{\mathbb{R}_+}^2.$$

Denoting $\mathbb{E}_1 := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, we have

$$\begin{split} \|(D_{c} - mc^{2} - z)^{-1} - (S - z)^{-1} \mathbb{E}_{1}\|_{\mathbb{R}}^{2} &= \sup_{\phi \in L^{2}(\mathbb{R}, \mathbb{C}^{2})} \frac{\|((D_{c} - mc^{2} - z)^{-1} - (S - z)^{-1} \mathbb{E}_{1})\phi\|_{\mathbb{R}}^{2}}{\|\phi\|_{\mathbb{R}}^{2}} \\ &\geq \sup_{\phi_{\mathbb{A}_{c}} \in L^{2}(\mathbb{R}, \mathbb{C}^{2})} \frac{\|((D_{c} - mc^{2} - z)^{-1} - (S - z)^{-1} \mathbb{E}_{1})\phi_{\mathbb{A}_{c}}\|_{\mathbb{R}}^{2}}{\|\phi_{\mathbb{A}_{c}}\|_{\mathbb{R}}^{2}} \\ &\geq \sup_{\phi \in L^{2}(\mathbb{R}_{+}, \mathbb{C}^{2})} \frac{\|((D_{0,c} - mc^{2} - z)^{-1} - (S_{\mathbb{A}_{c}} - z)^{-1} \mathbb{E}_{1})\phi\|_{\mathbb{R}_{+}}^{2}}{(1 + |\eta_{c}(\alpha)|^{2})\|\phi\|_{\mathbb{R}_{+}}^{2}} \\ &= \frac{1}{(1 + |\eta_{c}(\alpha)|^{2})} \|(D_{0,c} - mc^{2} - z)^{-1} - (S_{\mathbb{A}_{c}} - z)^{-1} \mathbb{E}_{1}\|_{\mathbb{R}_{+}}^{2}, \end{split}$$

where

$$(S_{\mathbb{A}_c} - z)^{-1}(x, y; z) := \mathfrak{G}(x, y; z) + \mathfrak{G}(x, -y; z)\eta_c(\alpha).$$

Note that although $(S_{\mathbb{A}_c}-z)^{-1}$ is a well defined integral operator, it is not necessarily a resolvent of a closed, densely defined operator, symbolically denoted as $S_{\mathbb{A}_c}$. It is well known [18] that

$$\|(D_c - mc^2 - z)^{-1} - (S - z)^{-1} \mathbb{E}_1\|_{\mathbb{R}} \xrightarrow[c \to +\infty]{} 0.$$
 (24)

At the same time, $\eta_c(\alpha) \to \xi(\alpha)$ as $c \to +\infty$. So it remains to verify that $(S_{\mathbb{A}_c} - z)^{-1}$ is close to $(S_0 - z)^{-1}$ in the limit $c \to +\infty$. This follows by the Hilbert–Schmidt bound:

$$||(S_{\mathbb{A}_c} - z)^{-1} - (S_0 - z)^{-1}||_{HS}^2 = \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} |\Im(x, -y; z)|^2 |\eta_c(\alpha) - \xi(\alpha)|^2 dx dy$$

$$= \frac{m}{2z} |\eta_c(\alpha) - \xi(\alpha)|^2 \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} \exp(-2\Im(\sqrt{2mz})|x + y|) dx dy$$

together with the aforementioned convergence $\eta_c(\alpha) \to \xi(\alpha)$ as $c \to +\infty$.

Acknowledgments

We are grateful to Lukáš Heriban for valuable discussions on the non-relativistic limit. The authors were supported by the EXPRO grant No. 20-17749X of the Czech Science Foundation.

References

- [1] N. Arrizibalaga, L. Le Treust, and N. Raymond, On the MIT bag model in the non-relativistic limit, Comm. Math. Phys. **354** (2017), 641–669.
- [2] J. Behrndt, D. Frymark, M. Holzmann, and Ch. Stelzer-Landauer, *Nonrelativistic limit of generalized MIT bag models and spectral inequalities*, (2023), arXiv:2312.14550.
- [3] S. Benvegnu and L. Dabrowski, *Relativistic point interaction in one dimension*, Lett. Math. Phys. **30** (1994), 159–167.
- [4] L. Cossetti, L. Fanelli, and D. Krejčiřík, Absence of eigenvalues of Dirac and Pauli Hamiltonians via the method of multipliers, Comm. Math. Phys. **379** (2020), 633–691.

- [5] L. Cossetti and D. Krejčiřík, Absence of eigenvalues of non-self-adjoint Robin Laplacians on the half-space, Proc. London. Math. Soc. 121 (2020), 584–616.
- [6] J.-C. Cuenin, Estimates on complex eigenvalues for Dirac operators on the half-line, Integral Equ. Oper. Theory **79** (2014), 377–388.
- [7] J.-C. Cuenin, A. Laptev, and Ch. Tretter, Eigenvalue estimates for non-selfadjoint Dirac operators on the real line, Ann. Henri Poincaré 15 (2014), 707–736.
- [8] S. Dineen, Complex Analysis in Locally Convex Spaces, North Holland, 2012.
- [9] P. N. Dowling and B. Turret, Complex strict convexity of absolute norms on \mathbb{C}^n and direct sums of Banach spaces, J. Math. Anal. **323** (2006), 930–937.
- [10] P. D'Ancona, L. Fanelli, D. Krejčiřík, and N.M. Schiavone, Localization of eigenvalues for non-self-adjoint Dirac and Klein-Gordon operators, Nonlinear Anal. 214 (2022), 112565.
- [11] P. D'Ancona, L. Fanelli, and N.M. Schiavone, Eigenvalue bounds for non-selfadjoint Dirac operators, Math. Ann. 383 (2022), 621–644.
- [12] A. Enblom, Estimates of eigenvalues of Schrödinger operators on the half-line with complex-valued potentials, Oper. Matrices 11 (2017), 369–380.
- [13] _____, Resolvent estimates and bounds on eigenvalues for Dirac operators on the half-line, J. Phys. A: Math. Theor. **51** (2018), 165203.
- [14] L. Fanelli, D. Krejčiřík, and L. Vega, Absence of eigenvalues of two-dimensional magnetic Schrödinger operators, J. Funct. Anal. 275 (2018), 2453–2472.
- [15] L. Fanelli, D. Krejčiřík, and L. Vega, Spectral stability of Schrödinger operators with subordinated complex potentials, J. Spectr. Theory 8 (2018), 575–604.
- [16] R. L. Frank, Eigenvalue bounds for Schrödinger operators with complex potentials, Bull. London Math. Soc. 43 (2011), 745–750.
- [17] R. L. Frank, A. Laptev, and R. Seiringer, A sharp bound on eigenvalues of Schrödinger operators on the half-line with complex-valued potentials, Spectral Theory and Analysis (J. Janas, P. Kurasov, A. Laptev, S. Naboko, and G. Stolz, eds.), Oper. Theory Adv. Appl., vol. 214, Springer, Basel, 2011, pp. 39–44.
- [18] F. Gesztesy, H. Grosse, and B. Thaller, A rigorous approach to relativistic corrections of bound state energies for spin-1/2 particles, Annales de l'I.H.P., section A 40 (1984), 159— 174.
- [19] M. Hansmann and D. Krejčiřík, The abstract Birman-Schwinger principle and spectral stability, J. Anal. Math. 148 (2022), 361–398.
- [20] L. Heriban and M. Tušek, Non-self-adjoint relativistic point interaction in one dimension,
 J. Math. Anal. Appl. 516 (2022), no. 2, 126536.
- [21] J. H. Jeans, The mathematical theory of electricity and magnetism, Cambridge University Press, 1915, third edition.
- [22] T. Kato, Wave operators and similarity for some non-selfadjoint operators, Math. Ann. 162 (1966), 258–279.

- [23] D. Krejčiřík, A. Laptev, and F. Štampach, Spectral enclosures and stability for non-self-adjoint discrete Schrödinger operators on the half-line, Bull. London. Math. Soc. 54 (2022), 2379–2403.
- [24] D. Krejčiřík and P. Siegl, Elements of spectral theory without the spectral theorem, In Non-selfadjoint operators in quantum physics: Mathematical aspects (432 pages), F. Bagarello, J.-P. Gazeau, F. H. Szafraniec, and M. Znojil, Eds., Wiley-Interscience, 2015.
- [25] D. Krejčiřík, P. Siegl, M. Tater, and J. Viola, *Pseudospectra in non-Hermitian quantum mechanics*, J. Math. Phys. **56** (2015), 103513.
- [26] E. H. Lieb and R. Seiringer, *The stability of matter in quantum mechanics*, Cambridge University Press, 2009.
- [27] H. Mizutani and N. M. Schiavone, Spectral enclosures for Dirac operators perturbed by rigid potentials, Rev. Math. Phys. **34** (2022), 2250023.
- [28] B. Thaller, The Dirac equation, Springer-Verlag, Berlin Heidelberg, 1992.
- [29] M. Tušek, Approximation of one-dimensional relativistic point interactions by regular potenials revised, Lett. Math. Phys. 110 (2020), 2585–2601.