

Pentagon equations, Voronoï tilings and pure braid groups invariant

Illia E. Rohozhkin

May 17, 2024

I express my gratitude to my scientific supervisor Manturov Vasily Olegovich for valuable advice and assistance in writing this article and to Igor G. Korepanov for the original ideas that formed its basis. I also express my gratitude to Kim Seongjeong, Igor M. Nikonov, Alexey V. Sleptsov, Wan Zheyuan and Louis Kauffman for their review and comments.

Abstract

We construct $(2n-4) \times (2n-4)$ matrices corresponding to the motion of points on standard round sphere from the point of view of Delaunay triangulations. We define homomorphism from spherical pure braids on n strands to the product of these matrices for $n > 5$.

Keywords: spherical braid group, Delaunay triangulation, matrices, photography method, pentagon equation.

MSC: 57M25, 57M27.

1 Introduction

In [3, 5, 2] a family of groups Γ_n^4 was defined. These groups depend on the parameter $n > 4$ and naturally describe the motion of points on the plane. The map $f : PB_n \rightarrow \Gamma_n^4$ is constructed by using Voronoï diagrams and Delaunay triangulations defined from these points. Isotopic braids correspond to equivalent word in Γ_n^k due to relations of this group.

In [6] the 2×2 matrix is assigned to each Γ_n^4 generator.

In this paper we will consider a moving of n points on the sphere. The $(2n-4) \times (2n-4)$ matrices will be defined. These matrices will satisfy specified relations. We will build a map from spherical braid group to the $GL_{2n-4}(F)$ group and prove that this map is well defined homomorphism.

2 Basic definitions

In [1] the n -strand braid group is defined to be isomorphic to the fundamental group $\pi_1(\text{UConf}_n(X))$, where X is two-dimensional manifold, $\text{UConf}_n(X)$ is n -unordered configuration space for X .

The n -strands pure braid group is defined as $\pi_1(\text{Conf}_n(X))$ group, where $\text{Conf}_n(X)$ is n -ordered configuration space for two-dimensional manifold X .

In [8] the spherical braid group on n strands, denoted SB_n or $B_n(S^2)$, is defined as the fundamental group of the configuration space of the sphere:

$$B_n(S^2) = \pi_1(\text{Conf}_n(S^2)).$$

In [3] pure braids are considered as dynamical systems whose initial and final states coincide, see Fig. 1.




Figure 1: Dynamical system corresponding to $b_{ij} \in PB_n$ generator.

Consider now n points moving on the sphere. Spherical pure n -strand braids can be considered as n -point dynamical systems. By SPB_n denote the spherical pure n -strand braid group.

Convention. The field of n ($n > 5$) pairwise distinct elements is denoted by F . The general linear group of degree $2n - 4$ over F is denoted by $\text{GL}_{2n-4}(F)$. The standard round sphere of radius 1 is denoted by S .

3 Delaunay triangulation on a sphere

Definition 1. We say that n points are in general position¹ on the sphere if there are no four points placed on same circle such that remaining $n - 4$ points are located at a larger distance from one of the centers² of this circle.

Consider P be a set of points in general position on a sphere S . Voronoï tiling associated with P is the set of all points on the sphere S defined as:

$$U_i = \{\forall j \neq i : u, |u - u_i| \leq |u - u_j|\}.$$

The Delaunay triangulation of P corresponds to the dual graph of the Voronoï diagram for P . Let's count the number of triangles in such triangulation

¹The “general position of points” has a broader meaning. See book [2] for more details.

²A circle on a sphere has two centers.

on the sphere S . The Euler characteristic of sphere is equal to 2:

$$V - E + F = 2,$$

where V , E , and F are respectively the numbers of vertices, edges and faces of the triangulated sphere.

Let us express the number of edges in terms of the number of faces for triangulation on the sphere. Since each face contains exactly 3 edges and each edge connects 2 faces consequently, it follows that: $V - \frac{3}{2}E + F = 2$. Finally, we obtain:

$$F = 2V - 4. \quad (1)$$

As the points move, they may not be in general position at some moment. In this case the Delaunay triangulation undergoes a flip, see Fig. 2.




Figure 2: A flip.

However the number of triangles in a triangulation on a sphere always remains the same.

4 The pure braid group and matrices

In [3, 4] both Γ_n^4 group and map $f_n : PB_n \rightarrow \Gamma_n^4$ were described. The generators of Γ_n^4 correspond to the flips of Delaunay triangulation.

It was shown in [6] that each Γ_n^4 generator can be associated with 2×2 matrix of the form:

$$\begin{pmatrix} \frac{\zeta_k - \zeta_j}{\zeta_l - \zeta_j} & \frac{\zeta_i - \zeta_j}{\zeta_l - \zeta_j} \\ \frac{\zeta_k - \zeta_l}{\zeta_j - \zeta_l} & \frac{\zeta_i - \zeta_l}{\zeta_j - \zeta_l} \end{pmatrix}. \quad (2)$$

The sum of the elements in each column of this matrix is 1.

The pentagon relation in the Γ_n^4 group can be written in 3×3 matrices. These matrices have the following form:

$$\begin{pmatrix} \Delta_{ijk} & \Delta_{ikl} & \Delta \\ \Delta_{jkl} & \Delta_{ijl} & \Delta \\ \Delta & \Delta & \Delta \end{pmatrix} = \begin{pmatrix} \frac{\zeta_k - \zeta_j}{\zeta_l - \zeta_j} & \frac{\zeta_i - \zeta_j}{\zeta_l - \zeta_j} & 0 \\ \frac{\zeta_k - \zeta_l}{\zeta_j - \zeta_l} & \frac{\zeta_i - \zeta_l}{\zeta_j - \zeta_l} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (3)$$

This matrix corresponds to one of the flips inside the pentagon.

5 $(2n - 4) \times (2n - 4)$ matrices

Consider n moving points on the sphere S . Assume these points are in general position over a period of time t . In this case the Voronoï diagram is graph of valency 3. Let $\mathcal{T}_t$ be a Delaunay triangulation corresponding to the Voronoï diagram during t . To $\mathcal{T}_t$ associate the $(2n - 4)$ -dimensional vector space $V_{\mathcal{T}_t}$ over F . A basis of the $V_{\mathcal{T}_t}$ are triangles of the Delaunay triangulation.

As above, the Delaunay triangulation undergoes a flip at some critical moment when moving points are not in general position. Denote by $\mathcal{T}_p$ the new Delaunay triangulation and denote by $V_{\mathcal{T}_p}$ the vector space related to this triangulation.

To each flip assign the linear operator γ :

$$\gamma : V_{\mathcal{T}_t} \rightarrow V_{\mathcal{T}_p}.$$

As in (4), let A_{ijkl} be a matrix that corresponds to γ operator. This matrix has the form:

$$A_{ijkl} = \begin{matrix} & \Delta_{\dots}^1 & \Delta_{\dots}^2 & \dots & \Delta_{jkl}^x & \dots & \Delta_{ijl}^y & \dots & \Delta_{\dots}^{s-1} & \Delta_{\dots}^s \\ \Delta_{\dots}^1 & 1 & 0 & \dots & 0 & \dots & 0 & \dots & 0 & 0 \\ \Delta_{\dots}^2 & 0 & 1 & \dots & 0 & \dots & 0 & \dots & 0 & 0 \\ \vdots & & & & \vdots & & & & & \\ \Delta_{ijk}^x & 0 & 0 & \dots & \frac{\zeta_k - \zeta_j}{\zeta_l - \zeta_j} & \dots & \frac{\zeta_i - \zeta_j}{\zeta_l - \zeta_j} & \dots & 0 & 0 \\ \vdots & & & & \vdots & & & & & \\ \Delta_{ikl}^y & 0 & 0 & \dots & \frac{\zeta_k - \zeta_l}{\zeta_j - \zeta_l} & \dots & \frac{\zeta_i - \zeta_l}{\zeta_j - \zeta_l} & \dots & 0 & 0 \\ \vdots & & & & \vdots & & & & & \\ \Delta_{\dots}^{s-1} & 0 & 0 & \dots & 0 & \dots & 0 & \dots & 1 & 0 \\ \Delta_{\dots}^s & 0 & 0 & \dots & 0 & \dots & 0 & \dots & 0 & 1 \end{matrix}, \quad (4)$$

where $s = 2n - 4$, $i, j, k, l \in \{1, 2, \dots, n\}$, $x, y \in \{1, 2, \dots, s\}$, $x < y$, i, j, k, l are four points are on the same circle such that other points are at a futher distance from one of the centers of this circle.

We claim that the set of all these matrices is a subgroup of $\text{GL}_{2n-4}(F)$. Indeed, it can easily be checked that these matrices are non-singular.

Statement 1. A_{ijkl} matrices satisfy the following relations:

- (a) $A_{ijkl}A_{stuv} = A_{stuv}A_{ijkl}$, where $|\{i, j, k, l\} \cap \{s, t, u, v\}| < 3$,
- (b) $A_{iklm}A_{ijkm}A_{jklm}A_{ijlm}A_{ijkl} = I_s$, where i, j, k, l, m are five points are located on the same circle such that other points are at a futher distance from one of the centers of this circle.

The first relation is called a *far commutativity relation*. The second relation is called a *pentagon relation*.

The proof of the commutativity relation is trivial. The matrices commute because they are block matrices with non-unit blocks at different places.

The pentagon relation for 3×3 matrices is verified by direct computation (see [6]). Larger matrices are obtained from 3×3 matrices by adding diagonal entries 1 corresponding to the fixed triangles. The matrix blocks either do not change or the size of blocks changes the same for all matrices. The pentagon relation is preserved in this case.

Both (a) and (b) relations are not unique. All other relations follow from (a), (b) trivially by conjugation, i.e. by rearranging rows and columns of the matrices.

6 The main theorem

Assume we have n moving points on the sphere S .

Let us consider a generator of n -strand spherical pure braid group, denoted b , as the dynamical system on the sphere, described in section (2). Let us enumerate $0 < t_1 < t_2 < \dots < t_l < 1$, such that at the moment t_z ($0 < z < 1$) four points are positioned on the one circle see Fig. 3. This moment corresponds




Figure 3: Codimension 1 event; it corresponds to the A_{ijkl} matrix.

to *codimension 1 event*. Each t_z corresponds to A_{ijkl} ($i, j, k, l \in n$). Denote A_{ijkl} by A_t . With the generator $b \in SPB_n$ we associate the following product:

$$f_n(b) = \prod_{t=1}^l A_t.$$

Proposition 1. *The map $f_n : SPB_n \rightarrow GL_{2n-4}$, which is defined above, is a well defined homomorphism.*

Proof. We consider isotopies between two spherical pure n -strand braids as isotopies of general position state of n points on the sphere. A finite number of events of codimension at most two correspond to the general position state isotopies. These events were classified in [5]. Now we list them explicitly.

- (i) One point moving on the sphere is tangent to the circle, which passes through three points, see Fig. 4. This means that the inverse matrix A_{ijkl}^{-1} exists for each A_{ijkl} generator.




Figure 4: Codimension 2 event; point l moves, being tangent to the circle, which passes through i , j and k .

- (ii) There are two sets A and B of four points, which are on the same circles such that $|A \cap B| \leq 2$, see Fig. 5. All other points are at a further distance from one of the centers of each circles. This means that corresponding matrices commute because corresponding to triangles transformations rows and columns are different in those matrices. This event corresponds to the far commutativity relation (a).




Figure 5: Codimension 2 event; two sets A and B of four points on the circles such that $|A \cap B| = 2$.

- (iii) There are five points $\{i, j, k, l, m\}$ on the same circle; all other points are at a further distance from one of the centers of this circle. We obtain the sequence of five subsets of $\{i, j, k, l, m\}$ with four points on the same circle, which corresponds to the flips on the pentagon, see Fig. 6. This corresponds to the pentagon relation (b).

□

Geometric interpretation of the (iii) event from the point of view of the braids group was described in [4]. In the same way as described in [6] we may verify the pentagon relation by direct computation. In this paper the “photography method” was also described. This method helps to understand pentagon relation geometrically.

7 Further directions

Among further directions which may appear in further papers is knots invariants constructing.




Figure 6: Codimension 2 event; a sequence of five subsets of $\{i, j, k, l, m\}$, corresponding to flips on the pentagon.

A braid closure allows to construct knot from the braid. Also it is possible to transform one braid to another by a sequence of Markov movements. The first move is conjugation. The second move is adding a strand to the braid and twisting it once with the last strand.

Having a braids invariant in matrices, we need to find some characteristics of the matrix that are invariant under the first and second Markov movements. It can be a trace or determinant of the matrix. This will allow us to construct an invariant for knots.

References

- [1] Vassily Manturov Knot Theory, second edition, 2018, Taylor & Francis Group.
- [2] V. Manturov, D. Fedoseev, S. Kim, I. Nikonov, Invariants and Pictures, May 11, 2020, World Scientific Publishing Company, ISBN-10: 9811220115, ISBN-13: 978-9811220111, 388 pages.
- [3] S. Kim and V. O. Manturov Artin's braids, braids for three space, and groups Γ_n^k and G_n^k , [arXiv:1902.11238v3](#) [math.GT] 8 Mar 2019.
- [4] V.O.Manturov, I.M.Nikonov. The groups Γ_n^4 , braids, and 3-manifolds, [arXiv:2305.06316v1](#) [math.GT] 10 May 2023.
- [5] Denis A. Fedoseev, Vassily O. Manturov and Igor M. Nikonov. Manifolds of triangulations, braid groups of manifolds, and the groups Γ_n^k , [arXiv:1912.02695v2](#) [math.GT] 21 Dec 2020.
- [6] Vassily Olegovich Manturov and Zheyang Wan, [arXiv:2305.11945v3](#) [math.GT] 7 Nov 2023, MSC 2020: 20F36, 13F60, 57K20, 57K31.
- [7] E.-K. Lee, A positive presentation for the pure braid group, Journal of the Chungcheong mathematical society, Vol. 23, No. 3, Sep., 2010.
- [8] Lei Chen and Nick Salter Section problems for configurations of points on the Riemann sphere, [arXiv:1807.10171v1](#) [math.GT] 26 Jul 2018.