# THE FREE BOUNDARY FOR SEMILINEAR PROBLEMS WITH HIGHLY OSCILLATING SINGULAR TERMS

MARK ALLEN, DENNIS KRIVENTSOV, AND HENRIK SHAHGHOLIAN

ABSTRACT. We investigate general semilinear (obstacle-like) problems of the form  $\Delta u = f(u)$ , where f(u) has a singularity/jump at  $\{u = 0\}$  giving rise to a free boundary. Unlike many works on such equations where f is approximately homogeneous near u = 0, we work under assumptions allowing for highly oscillatory behavior.

We establish the  $C^{\infty}$  regularity of the free boundary  $\partial \{u > 0\}$  at flat points. Our approach is to first establish that flat free boundaries are Lipschitz, using a comparison argument with the Kelvin transform. For higher regularity, we study the highly degenerate PDE satisfied by ratios of derivatives of u, using changes of variable and then the hodograph transform. Along the way, we prove and make use of new Caffarelli-Peral type  $W^{1,p}$  estimates for such degenerate equations. Much of our approach appears new even in the case of Alt-Phillips and classical obstacle problems.

# Contents

1. Introduction	1
1.1. Assumptions on nonlinearity	2
1.2. Main results	3
2. Preliminaries	4
3. Flat implies Lipschitz	10
4. Lipschitz implies $C^{1,\beta}$	12
5. Hodograph Transform	19
6. $W^{1,p}$ estimates	23
Acknowledgements	27
Declarations	27
References	28

#### 1. INTRODUCTION

In this paper, we consider one-phase semilinear (obstacle-like) problems of the form

(1.1) 
$$\Delta u = f(u), \quad u \ge 0,$$

where f is subject to certain conditions, listed below in (A1–4) One variation of our problem is a simple perturbation of the obstacle problem, such as the equation

(1.2) 
$$\Delta u = (2 - \log(u))\chi_{\{u>0\}}$$

studied in [9] and [19]. A second example is variations of the Alt-Phillips problem [3] given by equations like

$$\Delta u = u^{\gamma} (2 - \log(u)) \chi_{\{u > 0\}}, \qquad (-1/3 < \gamma < 1).$$

Our analysis encompasses many modifications of these problems, even allowing f to fluctuate between the sub- and supercritical ranges for the Alt-Phillips problem such as in

$$\Delta u = u^{\alpha \sin(\log(2 - \log u))}$$

For this kind of one-phase problem, obtaining optimal estimates on the growth of u away from the free boundary  $\{u > 0\}$  is not difficult, and we do so in Section 2. Our main interest, however, is in the regularity of the free boundary itself. Here it is helpful to consider (1.2), and observe that at any free boundary point, the blow-up limits of solutions in the natural scaling for the equation solve the *classical obstacle problem* (see [19]). As such, there are two kinds of free boundary points: ones where the blow-up limit is a regular free boundary point for the obstacle problem (i.e. flat points) and ones where it is a singular point. In this paper, our goal is to study the free boundary near flat points, and show that it is locally a smooth hypersurface. For an equation like (1.2), this turns out to be quite challenging, in part due to how different the PDE is from the one satisfied by the blow-up limit (they do not even have the same natural scaling).

Since the seminal paper [6], many techniques have been developed to study both the initial free boundary regularity as well as higher regularity for obstacle-type problems. Semiconvexity estimates [13] are useful for obtaining Lipschitz regularity and then  $C^{1,\beta}$  regularity of the free boundary. In our situation when f is in the subcritical range of Alt-Phillips (or even a perturbation of the obstacle problem such as (1.2)), the solution u will no longer be semiconvex. Methods to obtain directional monotonicity from the equation that  $\partial_e u$  satisfies (see [20]) are also unavailable due to the possible singular behavior of f. Since [4] the boundary Harnack principle has been a powerful tool to move from Lipschitz to  $C^{1,\beta}$  regularity of the free boundary. This technique has seen renewed interest in recent years by utilizing the boundary Harnack inequality for inhomogeneous equations [1,2,21,24]. Unfortunately, the boundary Harnack inequality corresponding to our situation may not hold. Typically, as time progresses, robust techniques are developed which apply to more general problems. Here, with a variety of modern robust techniques appearing to not apply, to prove that flat free boundaries are Lipschitz graphs we instead return to a more rigid comparison technique using the Kelvin transform as introduced in [22].

Higher regularity of the free boundary for the obstacle problem was originally obtained with the Legendre transform [18]. More recently, the higher order boundary Harnack principle [12] has proven quite adaptable for obstacle-type problems such as the thin obstacle problem [11, 17, 21]. An alternative approach to obtaining the higher-order boundary Harnack principle arises from acquiring Schauder estimates for a highly degenerate elliptic equation. This approach is discussed in both [23] and [25], where  $C^1$  regularity of the boundary is assumed in both references. We utilize this new technique to prove Cordes-Nirenberg estimates for the highly degenerate elliptic equation to move from Lipschitz to  $C^{1,\beta}$  regularity of the free boundary. In both [23,25] the Schauder theory gives  $C^{\infty}$  and real-analyticitly respectively. But again in our problem, the method as given in [23] and [25] (in particular how the free boundary is flattened) fails to move beyond  $C^{1,\beta}$  regularity, see the discussion in Section 5. Instead, we more carefully flatten the free boundary by utilizing a modified partial hodograph transform. We could then prove Schauder estimates as in [23,25] on the nonlinear equation obtained in the hodograph variables, but opt instead to prove Caffarelli-Peral estimates. These estimates are more easily obtained, and when applied in the hodograph variables they show that a  $C^{1,\beta}$  free boundary is  $C^{\infty}$ .

1.1. Assumptions on nonlinearity. Below are the assumptions we make regarding the nonlinearity f(t). Note that since our focus lies on the local behavior of solutions near u = 0, assumptions pertaining to  $t \to \infty$  are not relevant for questions regarding existence and boundedness. Define

$$F(t) = \int_0^t f(s) ds,$$

which is guaranteed to exist under Assumptions (A1,2) below for all f we consider.

For  $-1 < \gamma_1 < \gamma_2 < 1$  and M > 0, we say that  $f \in \mathcal{F} = \mathcal{F}(M, \gamma_1, \gamma_2)$  if the following assumptions are satisfied:

(A1) f is continuous and nonnegative on (0, 1].

(A2) f satisfies the bounds

(1.3) 
$$\frac{1}{M}t^{\gamma_2} \le f(t) \le Mt^{\gamma_1} \quad \text{for } 0 < t \le 1.$$

(A3) the primitive F satisfies

(1.4) 
$$1 + \gamma_1 \le t \frac{F'(t)}{F(t)} \le 1 + \gamma_2 \quad \text{for } 0 < t \le 1.$$

(A4) f satisfies the additional bound

(1.5) 
$$f(at) \ge a^{(n+2)/(n-2)} f(t)$$
 whenever  $0 \le a < t < 1$ .

For the higher regularity results (Lipschitz free boundaries smooth), our approach also requires the additional assumption

$$\gamma_1 > -\frac{1}{3}.$$

The case of  $\gamma_1 \leq -\frac{1}{3}$  results in a linearized problem with a differently structured boundary condition, and we leave this case as an intersting open problem.

The following easier-to-verify assumption on the derivative of f implies (A2–4) and already accounts for most of the examples we have in mind: f is continuously differentiable on (0, 1] and

$$\gamma_1 \le t \frac{f'(t)}{f(t)} \le \gamma_2 \quad \text{for } 0 < t \le 1.$$

In particular, (4) follows from this if  $\gamma_2 \leq \frac{n+2}{n-2}$ , which is weaker than the standing assumption that  $\gamma_2 < 1$ .

To explain the nature of these assumptions, first recall that for homogeneous  $f(t) = t^{\gamma}$  (the Alt-Phillips problem, studied in [3]), the range  $\gamma \in (-1, 1)$  exhibits free boundary behavior. If  $\gamma \geq 1$ , the resulting equation instead satisfies a strong maximum principle and has no free boundary. The range  $\gamma \leq -1$  has very different monotonicity properties and free boundary behavior, and we do not consider it here (it has been studied recently in [10]). Our assumption (A2) asks that f remains in the free boundary Alt-Phillips range without demanding any homogeneity of f at all.

This turns out to be not scale-invariant under the natural (implicit) scaling of the problem. The purpose of assumption (A3) is to guarantee that solutions to the ODE h'' = f(h), if rescaled via  $h_r(t) = h(rt)/h(r)$ , continue to satisfy (A2) This allows for f to oscillate between different homogeneities, but only in a controlled fashion (on a scale comparable to  $\log t$ ). The assumption (A4) will be used to ensure that solutions interact favorably with their Kelvin transforms, and in particular they satisfy a comparison-type property.

Some examples of f(t) satisfying our assumptions would be  $f(t) \equiv 1$  or  $f(t) = c + g(t) \ge 0$ with  $g(t) \in C^1$  such as in the classical obstacle problem. We allow for singular and degenerate perturbations of the classical obstacle problem such as  $f(t) = 2 - \log(t)$  or  $f(t) = 1/(2 - \log(t))$ . We allow for Alt-Phillips equations  $f(t) = t^{\gamma}$  for  $\gamma \in (-1, 1)$  as well as perturbations such as  $f(t) = t^{\gamma}(2 - \log t)^{\pm 1}$ . We allow f(t) to oscillate between the sub and supercritical regions such as  $f(t) = t^{\alpha \sin(\log(2 - \log t))}$  for numbers |a| small enough, as well as other functions oscillating at this rate. To allowing oscillations, our methods do not utilize a particular blow-up limit of u, and hence that of f, since in the rescaling we only obtain a function in the same class  $\mathcal{F}(M, \gamma_1, \gamma_2)$ .

1.2. Main results. The solutions of (1.1) we consider are minimizers of the functional

(1.6) 
$$J(w) := \int_{\Omega} \frac{1}{2} |\nabla w|^2 + F(w) \, dx,$$

over the class of functions

$$\{w \in W^{1,2}(\Omega) : 0 \le w \le 1, w - v \in W^{1,2}(\Omega)\}$$

for some fixed  $v \in W^{1,2}(\Omega)$ ; we call these  $H_{\leq 1}$ -minimizers. Any minimizer of J over competitors in  $W^{1,2}(\Omega)$  will be a  $H_{\leq 1}$ -minimizer on a small enough neighborhood of a free boundary point. See Definition 2.4 for further discussion.

Throughout the paper, we will refer to the unique positive solution h (see Proposition 2.2) of the ordinary differential equation

(1.7) 
$$\begin{cases} h''(t) = f(h(t)) & \text{in } t > 0, \\ h(0) = h'(0) = 0. \end{cases}$$

In Section 2 we show that a minimizer u of (1.6) will exhibit the same asymptotic growth as h from a free boundary point, namely

$$ch(r) \le \sup_{B_r(x)} u \le Ch(r), \quad \forall x \in \partial \{u > 0\}.$$

We then examine *flat points* of the free boundary: ones where, after a possible rotation and translation,

(1.8) 
$$h(x \cdot e_n - \epsilon) \le u(x) \le h(x \cdot e_n + \epsilon).$$

Our main results regarding the free boundary near these flat points are outlined below. In Section 3, we utilize the Kelvin transform to demonstrate that flatness implies directional monotonicity and Lipschitz regularity of the free boundary.

**Theorem 1.1.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_1$ , and fix  $\delta > 0$ . There exists  $\epsilon > 0$  depending on  $\delta, n, M, \gamma_1, \gamma_2$ , such that if u satisfies (1.8) then  $\partial_{\nu} u \geq 0$  in  $B_{1/2}$  whenever  $\nu \cdot e_n \geq \delta$ . In particular the free boundary  $\partial \{u > 0\} \cap B_{1/2}$  is a Lipschitz graph.

In Section 4 to move from Lispchitz to  $C^{1,\beta}$  regularity of the free boundary, we use the technique in [23] and [25] to obtain Cordes-Nirenberg estimates on the equation satisfied by ratios of derivatives  $u_e/u_{e_n}$ , where e is any unit vector and  $e_n$  is a direction u is monotone from Theorem 1.1. This, together with appropriate change-of-variables arguments

**Theorem 1.2.** Assume  $\gamma_1 > -\frac{1}{3}$ . Let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_1$ , and fix  $0 < \beta < 1$ . There exists  $\epsilon > 0$  depending on  $\beta, n, M, \gamma_1, \gamma_2$  such that if u satisfies (1.8) then  $\partial \{u > 0\} \cap B_{1/2}$  is a  $C^{1,\beta}$  graph.

In Section 5, we employ a variant of the partial hodograph transform to flatten the free boundary. We then elaborate on how to use  $W^{1,p}$  Caffarelli-Peral estimates to yield  $C^{\infty}$  regularity of the free boundary. In Section 6, we establish these Caffarelli-Peral estimates to achieve

**Theorem 1.3.** Assume  $\gamma_1 > -\frac{1}{3}$ . Let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_1$ . There exists  $\epsilon > 0$  depending on  $n, M, \gamma_1, \gamma_2$  such that if u satisfies (1.8) then  $\partial \{u > 0\} \cap B_{1/2}$  is  $C^{\infty}$ .

## 2. Preliminaries

We list some additional consequences for f and F from our assumptions (A1–4).

**Proposition 2.1.** Let  $F \in \mathcal{F}$ . For 0 < t, at < 1 we have

$$a^{1+\gamma_2}F(t) \le F(at) \le a^{1+\gamma_1}F(t) \quad for \ a \le 1.$$
  
 $a^{1+\gamma_1}F(t) \le F(at) \le a^{1+\gamma_2}F(t) \quad for \ a > 1.$ 

*Proof.* First let  $a \leq 1$ . By scaling and considering  $F_r := F(tr)/F(r)$  we may assume t = 1 and F(1) = 1. Consider  $G(t) := F(t) - t^{1+\gamma_2}$ . Then G(1) = 0 and  $G(0) \geq 0$ . If G obtains an interior minimum at  $t_0$  with  $G(t_0) \leq 0$ , then

$$0 = G'(t_0) = F'(t_0) - (1 + \gamma_2)t_0^{\gamma_2},$$

or

$$F'(t_0) = (1 + \gamma_2) t_0^{\gamma_2}.$$

Since  $G(t_0) \leq 0$ , then  $F(t_0) \leq t_0^{1+\gamma_2}$ , so that

$$(1 + \gamma_2) \le t_0 F'(t_0) / F(t_0).$$

From assumption (1.4) we must have equality above so that  $G(t) \ge 0$ , and this gives the first inequality. A similar argument shows

$$F(at) \le a^{1+\gamma_1} F(t).$$

By reversing the roles of t, at we obtain the inequalities for a > 1.

Here, we compile properties of the unique positive solution h to the ordinary differential equation (1.7). While equation (1.7) doesn't admit a single solution, as  $h(t) \equiv 0$  is one solution and  $h(t-t_0)$  is another whenever h is a solution and  $t_0$  is positive, we prove here the existence of a unique positive solution.

**Proposition 2.2.** Assume that  $F \in \mathcal{F}$ . Then there exists a unique positive solution h to (1.7). Furthermore h satisfies the following properties. There exists a constant  $C_1$  such that

$$C_{1}^{-1}t^{2/(1-\gamma_{2})} \leq h(t) \leq C_{1}t^{2/(1-\gamma_{1})} \quad \text{for } 0 < t < 1$$

$$\frac{2}{1-\gamma_{1}} \leq t\frac{h'(t)}{h(t)} \leq \frac{2}{1-\gamma_{2}} \quad \text{for } 0 < t < 1$$

$$\frac{1+\gamma_{1}}{1-\gamma_{1}} \leq t\frac{h''(t)}{h'(t)} \leq \frac{1+\gamma_{2}}{1-\gamma_{2}} \quad \text{for } 0 < t < 1$$

$$K^{C_{1}}t \leq h^{-1}(Kh(t)) \leq K^{C_{1}^{-1}}t \quad \text{for all } 0 < K \leq 1.$$

$$K^{C_{1}^{-1}}t \leq h^{-1}(Kh(t)) \leq K^{C_{1}}t \quad \text{for all } K > 1.$$

*Proof.* We note that a solution  $h_{\gamma}$  to

$$\begin{cases} y'' = cy^{\gamma} \\ y(0) = y'(0) = 0 \end{cases}$$

is given by

$$h_{\gamma}(t) = \left[\frac{C}{2} \frac{(1-\gamma)^2}{(1+\gamma)}\right]^{1/(1-\gamma)} t^{2/(1-\gamma)}.$$

Thus,  $h_{\gamma_2} \leq h_{\gamma_1}$ , and  $h_{\gamma_2}$  is a subsolution to (1.7) whereas  $h_{\gamma_1}$  is a supersolution. Thus, by the method of sub and supersolutions, there exists a positive solution h to (1.7) over the interval  $0 < t \leq 1$ . We now show that there is a maximal such positive solution  $h_{\text{max}}$ . Indeed, consider the solution to

$$\begin{cases} y'' = Ch(y) \\ y(0) = \epsilon, \quad y'(0) = 0. \end{cases}$$

which is bounded by below by h. Then as  $\epsilon \to 0$  we obtain  $h_{\text{max}}$ . Furthermore, any positive solution is bounded from below by  $h_{\text{max}}(t-\epsilon)$ . Then  $h = h_{\text{max}}$  is a unique positive solution. We note that (1.7) can be rewritten as solving

$$\begin{cases} h'(t) = \sqrt{2F(h(t))} \\ h(0) = 0. \end{cases}$$

The existence and uniqueness for t > 1 is immediate since h is increasing and  $\sqrt{2F}$  is  $C^1$  when F is positive.

If we rescale with  $h_r(t) := h(rt)/h(r)$ , then

$$h_r'(t) = \sqrt{2F_r(h_r(t))},$$

with  $F_r(t) := (r^2/h^2(r))F(th(r))$ . We remark that

$$F_r(1) = \frac{r^2}{2} \frac{[h'(r)]^2}{h^2(r)},$$

and that  $F_r$  also satisfies (1.4). Also,  $h_r(1) = 1$ . From Proposition 2.1 we have  $F_r(t) \ge t^{1+\gamma_2}F_r(1)$ . Consider now the function  $g(t) = t^{2/(1-\gamma_2)}$ , and suppose that

$$r\frac{h'(r)}{h(r)} = \sqrt{2F_r(1)} \ge (1+\epsilon)\frac{2}{1-\gamma_2}$$

Then

$$g'(t) = \frac{2}{1 - \gamma_2} t^{(1 + \gamma_2)/(1 - \gamma_2)} < \sqrt{2F_r(1)} t^{(1 + \gamma_2)/(1 - \gamma_2)} = \sqrt{2F_r(1)} g(t)^{(1 + \gamma_2)/2} \le \sqrt{2F_r(g(t))}.$$

Then g(t) is a subsolution and lies below the solution  $h_r$  in a neighborhood of t = 1. However,  $h_r(1) = 1 = g(1)$ , and this gives a contradiction. Then

(2.2) 
$$r\frac{h'(r)}{h(r)} = \sqrt{2F_r(1)} \le \frac{2}{1-\gamma_2}.$$

A similar argument with  $g(t) = t^{2/(1-\gamma_1)}$  as a supersolution gives

(2.3) 
$$F_r(1) = r \frac{h'(r)}{h(r)} \ge \frac{2}{1 - \gamma_1}.$$

Next, we consider th''/h'. We have

$$t\frac{h''}{h'} = t\frac{f(h)}{h'} = t\frac{f(h)}{h'}\frac{h'}{h'}\frac{h}{h} = \frac{hf(h)}{2F(h)}t\frac{h'}{h}.$$

Using (2.2), (2.3), and assumption (1.4) we conclude the estimate for th''/h.

Finally, just as in the proof for F in Proposition 2.1, we obtain for h that

$$h(K^{C_1}t) \le Kh(t) \le h(K^{C_1^{-1}}t) \quad \text{for } K \le 1$$
  
 $h(K^{C_1^{-1}}t) \le Kh(t) \le h(K^{C_1}t) \quad \text{for } K > 1.$ 

By applying  $h^{-1}$  to all expressions in both inequalities above, we obtain the final two inequalities in (2.1).

We state here a scaling result for F.

**Proposition 2.3.** Assume  $F \in \mathcal{F}(M, \gamma_1, \gamma_2)$ . Then the rescaling

$$F_r(t) := \frac{r^2}{h^2(r)} F(th(r))$$

satisfies  $F \in \mathcal{F}(M, \gamma_1, \gamma_2)$  for all 0 < r < 1.

*Proof.* It is easy to check that  $F_r(t)$  satisfies (1.4). Then from Proposition 2.1 we have

$$F_r(1)t^{1+\gamma_2} \le F_r(t) \le F_r(1)t^{1+\gamma_1}$$

Now  $F_r(1) = rh'(r)/h(r)$ , so from Proposition 2.2 we have

$$\frac{2}{1-\gamma_1}t^{1+\gamma_2} \le F_r(t) \le \frac{2}{1-\gamma_2}t^{1+\gamma_1}.$$

Finally, since f satisfies (1.5)

$$\frac{f_r(at)}{f_r(t)} = \frac{f(ath(r))}{f(th(r))} \ge a^{(n+2)/(n-2)},$$

so that  $f_r$  also satisfies (1.5).

To avoid making global assumptions on f, we will work with the following weaker notion of minimizer:

**Definition 2.4.** Let  $H_{\leq 1} = \{u \in W^{1,2}(\Omega) : 0 \leq u \leq 1\}$ . Given an  $F \in \mathcal{F}$ , we say that u is a  $H_{\leq 1}$ -minimizer on  $\Omega$  if  $u \in H_{\leq 1}$ , and for any  $v \in H_{\leq 1}$  with  $u - v \in W_0^{1,2}(\Omega)$  we have  $J(u) \leq J(v)$  (with J as in (1.6)).

Occasionally, we will utilize a rescaling such as  $u_r(x) := u(rx)/h(r)$ . In those instances, the rescaled function is a now a minimizer among functions  $w(x)h(r) \leq 1$ .

Any minimizer of J is a  $H_{\leq 1}$ -minimizer on a neighborhood of a point in  $\partial \{u > 0\}$  at which u is continuous, and so our results will apply to it. We refer to the classic works [14] and [15] on the existence and basic regularity theory for minimizers in general, which is not our primary focus here. However, we do note the following proposition, which will be useful for constructing comparison functions:

**Proposition 2.5.** Assume  $F \in \mathcal{F}$ , and fix a  $\phi \in H_{\leq 1}$ . Then there exists a  $H_{\leq 1}$ -minimizer u with  $u - \phi \in W_0^{1,2}$ , and there exists a constant  $\beta > 0$  such that any  $H_{\leq 1}$ -minimizer will be in the class  $C^{1,\beta}(\Omega)$ .

We provide only a sketch of the proof to show how to fit our assumptions into the standard framework of the calculus of variations. It is not difficult to see that minimizers of J over *arbitrary*  $W^{1,2}$  functions need not be regular, or even exist, as we made no assumptions on F when  $t \notin (0, 1)$ .

*Proof.* Let  $\tilde{F}$  be defined via

$$\tilde{F}(t) = \begin{cases} F(0) & t < 0\\ F(t) & 0 \le t \le 1\\ F(1) & t > 1, \end{cases}$$

and  $\tilde{J}$  the corresponding functional as in (1.6). Take any function  $u \in H_{\leq 1}$  with  $u - \phi \in W_0^{1,2}(\Omega)$ . Then we claim that u is a  $H_{\leq 1}$ -minimizer of J if and only if it is a minimizer of  $\tilde{J}$  (over arbitrary functions  $v \geq 0$  in  $W^{1,2}(\Omega)$  with  $v - \phi \in W_0^{1,2}(\Omega)$ ). Indeed, as  $J = \tilde{J}$  on  $H_{\leq 1}$ , it is clear for all such competitors in  $H_{\leq 1}$ . On the other hand, if u is a  $H_{\leq 1}$ -minimizer of J and v is a competitor in  $W_0^{1,2}(\Omega)$ , note that  $w = \max\{\min\{1, v\}, 0\} \in H_{\leq 1}$  is also a valid competitor and has  $\tilde{F}(w) = \tilde{F}(v)$ , so

$$J(v) = J(w) = J(w) \ge J(u).$$

It is clear from Assumption (A2) that  $\tilde{F} \in C^{0,\alpha}(\mathbb{R})$ , and it follows from the direct method that  $\tilde{J}$  admits a minimizer u with  $u - \phi \in W_0^{1,2}(\Omega)$  (we omit the details here). Arguing as above,  $u \in H_{\leq 1}$ : if not,  $w = \max\{\min\{1, u\}, 0\}$  has  $\tilde{J}(w) \geq \tilde{J}(u)$ , which leads to  $|\nabla u| = 0$  a.e. over  $\{u < 0\} \cup \{u > 1\}$ , a contradiction. This proves the existence claim. The  $C^{1,\beta}$  regularity follows directly from [14], Theorem 3.1.

We now bound the growth away from the free boundary  $\partial \{u > 0\}$ .

**Theorem 2.6.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_1$  with  $F \in \mathcal{F}$ , and assume  $x_0 \in B_{3/4} \cap \partial \{u > 0\}$ . 0. There exists C > 0 depending on  $M, \gamma_1, \gamma_2$  and  $\|u\|_{H^1(B_1)}$  such that

$$\sup_{B_r(x_0)} u \le Ch(r).$$

*Proof.* Suppose by way of contradiction the result is not true. Notice that by the  $C^{1,\beta}$  regularity, we need only prove the statement for r small. Then there exist

$$u_k \quad H_{\leq 1}\text{-minimizers of (1.6)}$$
$$x_k \in B_{3/4} \cap \partial \{u_k > 0\}$$
$$r_k \to 0$$

such that

(2.4) 
$$S_{r_k} := \sup_{B_{r_k}(x_k)} u_k \ge kh_k(r_k).$$

Furthermore, since the conclusion is true for  $r > r_0$  we may assume additionally<sup>1</sup>

$$\frac{S_{2r_k}}{h_k(2r_k)} \le \frac{S_{r_k}}{h_k(r_k)}$$

so that

$$(2.5) S_{2r_k} \le \frac{h_k(2r_k)}{h_k(r_k)} S_{r_k} \le CS_{r_k}$$

We now rescale with

$$\tilde{u}_k := \frac{u_k(x_k + r_k x)}{S_{r_k}},$$

noting that  $\sup_{\partial B_1} \tilde{u}_k = 1$ , it is bounded on  $B_2$  and that  $\tilde{u}_k$  is a  $H_{\leq 1}$ -minimizer on  $B_{1/r_k}$  of (1.6) with

$$\tilde{F}_k(w) = \frac{r^2}{S_{r_k}^2} F_k(wS_{r_k}).$$

If  $t_k$  is the unique point satisfying  $h_k(t_k) = S_{r_k} \ge kh(r_k)$ , then  $r_k/t_k \to 0$ . From Proposition 2.3 we have that

$$\tilde{F}_k(w) = \frac{r_k^2}{t_k^2} \frac{t_k^2}{h_k^2(t_k)} F_k(wh_k(t_k)) \to 0 \text{ as } k \to \infty.$$

In the limit we then have the following true in  $B_2$ 

$$\begin{cases} u_k \rightarrow u_0 \text{ in } H^1 \\ u_k \rightarrow u_0 \text{ uniformly} \\ u_0 \ge 0 \\ \Delta u_0 = 0. \end{cases}$$

Moreover, from uniform convergence  $u_0(0) = 0$ . However, we also have in the limit that

$$\sup_{\partial B_1} u_0 = 1,$$

but this will contradict the minimum principle.

We now aim to prove a nondegeneracy growth condition from a free boundary point.

**Lemma 2.7.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6) on  $B_1$  and assume  $F \in \mathcal{F}$ . There exists  $\epsilon = \epsilon(M, \gamma)$  such that if  $u \leq \epsilon$  on  $\partial B_1$ , then

$$u(x) \equiv 0 \quad for \ x \in B_{1/2}.$$

<sup>&</sup>lt;sup>1</sup>This is achieved by selecting  $r_k$  as the maximum radius for which (2.4) is valid, ensuring that it also holds with equality.

Proof. Let  $\overline{u}$  be a  $H_{\leq 1}$ -minimizer for (1.6) with  $\overline{F}(t) = \frac{M^{-1}}{\gamma+1}t^{\gamma+1}$  and with  $\overline{u} = \epsilon$  on  $\partial B_1$ . Since  $\gamma + 1 \geq 1$ , the corresponding functional  $\overline{J}$  is convex, so the  $H_{\leq 1}$ -minimizer  $\overline{u}$  is unique. From symmetrization techniques, we have that  $\overline{u}$  is radially symmetric and increasing as a function of the radius. We may apply the nondegeneracy estimate for supercritical Al-Phillips; however one can show directly that  $\overline{u} \equiv 0$  on  $B_{1/2}$ , or we may apply the nondegeneracy estimate from [3].

We will now show that if u is a minimizer of J, then  $u \leq \overline{u}$ . We note that since  $\overline{F}' \leq F'$ , then

$$F(u) - F(\overline{u}) \ge \overline{F}(u) - \overline{F}(\overline{u})$$

whenever  $u \geq \overline{u}$ , so that

(2.6) 
$$F(u) + \overline{F}(\overline{u}) \ge \overline{F}(u) + F(\overline{u})$$
 whenever  $u \ge \overline{u}$ .

Let us define  $w_1 = \min\{u, \overline{u}\}$  and  $w_2 = \max\{u, \overline{u}\}$ . Then

$$\begin{split} J(u) + \overline{J}(\overline{u}) &= \int_{B_1} \frac{1}{2} |\nabla u|^2 + \frac{1}{2} |\nabla \overline{u}|^2 + F(u) + \overline{F}(\overline{u}) \\ &= \int_{\{u \geq \overline{u}\}} \frac{1}{2} |\nabla u|^2 + \frac{1}{2} |\nabla \overline{u}|^2 + F(u) + \overline{F}(\overline{u}) \\ &+ \int_{\{u < \overline{u}\}} \frac{1}{2} |\nabla w_2|^2 + \frac{1}{2} |\nabla w_1|^2 + F(u) + \overline{F}(\overline{u}) \\ &= \int_{\{u \geq \overline{u}\}} \frac{1}{2} |\nabla w_2|^2 + \frac{1}{2} |\nabla w_1|^2 + F(u) + \overline{F}(\overline{u}) \\ &+ \int_{\{u < \overline{u}\}} \frac{1}{2} |\nabla w_2|^2 + \frac{1}{2} |\nabla w_1|^2 + F(u) + \overline{F}(\overline{u}) \\ &= \int_{\{u \geq \overline{u}\}} \frac{1}{2} |\nabla w_1|^2 + \frac{1}{2} |\nabla w_2|^2 + F(u) + \overline{F}(\overline{u}) \\ &+ \int_{\{u < \overline{u}\}} \frac{1}{2} |\nabla w_1|^2 + \frac{1}{2} |\nabla w_2|^2 + F(w_1) + \overline{F}(w_2) \\ &\geq \int_{\{u \geq \overline{u}\}} \frac{1}{2} |\nabla w_2|^2 + \frac{1}{2} |\nabla w_1|^2 + F(\overline{u}) + \overline{F}(u) \\ &+ \int_{\{u < \overline{u}\}} \frac{1}{2} |\nabla w_2|^2 + \frac{1}{2} |\nabla w_2|^2 + F(w_1) + \overline{F}(w_2) \\ &\geq \int_{\{u \geq \overline{u}\}} \frac{1}{2} |\nabla w_1|^2 + \frac{1}{2} |\nabla w_2|^2 + F(w_1) + \overline{F}(w_2) \\ &= J(w_1) + \overline{J}(w_2). \end{split}$$

Thus,  $w_1$  is a  $H_{\leq 1}$ -minimizer of J and  $w_2$  is a  $H_{\leq 1}$ -minimizer of  $\overline{J}$ . Since the minimizer of  $\overline{J}$  is unique, we conclude that  $w_2 \equiv \overline{u}$ .

We will now show that  $\overline{u} \equiv 0$  on  $B_{1/2}$ . Now  $\overline{u}/\epsilon$  is a  $H_{\leq 1}$ -minimizer with  $\epsilon^{\gamma-1}F$ . We will use  $v(x) = 2^{2/(1-\gamma)} (|x| - 1/2)_{+}^{2/(1-\gamma)}$  as a barrier. Note that

$$\Delta v(x) = 2^2 [\beta(\beta + n - 2) + |x|/(|x| - 1/2)]v^{\gamma}.$$

As long as  $2^2[\beta(\beta + n - 2) + |x|/(|x| - 1/2)] \leq \epsilon^{\gamma - 1}M^{-1}$ , then the argument above (showing that  $u \leq \overline{u}$ ) will also apply so that  $\overline{u} \leq v\epsilon$ . This shows that  $\overline{u} \equiv 0$  on  $B_{1/2}$ .

**Theorem 2.8.** (Optimal non-degeneracy) Let u be a  $H_{\leq 1}$ -minimizer of (1.6) on  $B_1$  and assume  $F \in \mathcal{F}$ . There exists c depending on  $M, \gamma$  such that if  $0 \in \partial \{u > 0\}$ , then

$$\sup_{B_r} u \ge ch(r) \quad \text{for all } r \le 1/2.$$

*Proof.* We choose  $\epsilon$  from Lemma 2.7 for  $F \in \mathcal{F}$  with C from Proposition 2.3. We then apply Lemma 2.7 to the rescaling

$$u_r(x) := \frac{u(rx)}{h(r)}$$

to conclude that  $\sup_{\partial B_{r/2}} \geq \epsilon$ .

## 3. FLAT IMPLIES LIPSCHITZ

We begin this section by giving the heuristic ideas behind combining the Kelvin transform and the lattice principle for functionals to conclude monotonicity in a cone of directions. This method was used in [22] and will work for functionals of the form

$$J(w) := \int_{B_R} \frac{1}{2} |\nabla w|^2 + F(w),$$

as long as F satisfies (A4) and is nondecreasing.

We use the Kelvin transformation on the ball  $B_1(te_n)$  with  $0 < t \le 3/2$  and denote

(3.1) 
$$x^* = \frac{x - te_n}{|x - te_n|^2} + te_n$$

We will let 0 < h < t, and let R = 2 + 1/(t - h). Suppose that we have a  $H_{\leq 1}$ -minimizer u in  $B_R(0)$  which satisfies  $u(x) \ge |x - te_n|^{2-n}u(x^*)$  whenever  $x = (x', x_n) \in B_1(te_n) \cap \{x_n = h\}$ . From Assumption (A4) we have  $f(a^{n-2}t) \ge a^{n+2}f(t)$ , so that

(3.2) 
$$a^{-2n}F(a^{n-2}t) - F(t) \ge a^{-2n}F(a^{n-2}s) - F(s),$$

whenever  $0 < a \le 1$  and  $0 < s \le t \le 1$ . If  $\Omega = B_1(te_n) \cap \{x_n \le h\}$ , then  $|x - te_n|^{2-n}u(x^*) \le u(x)$ on  $\partial\Omega$ . We now show how the lattice principle will give the same inequality on all of  $\Omega$ . If  $x \in \Omega$ , we define  $v(x) = |x - te_n|^{2-n}u(x^*)$  and note that v is a minimizer to

$$\tilde{J}(w) := \int_{\Omega} \frac{1}{2} |\nabla w|^2 + |x - te_n|^{-2n} F(|x - te_n|^{n-2}w),$$

among the class  $w(x)|x - te_n|^{n-2} \leq 1$ . We let  $w_1 = \min\{v, u\}$  and  $w_2 = \max\{v, u\}$ . Then

$$\begin{split} J(u) + \tilde{J}(v) &= J(u) + J(v) + \int_{\Omega} (|x - te_n|^{-2n} F(|x - te_n|^{n-2}v) - F(v)) \\ &= \int_{\{u \ge v\}} |\nabla u|^2 + F(u) + \int_{\{u < v\}} |\nabla u|^2 + F(u) \\ &+ \int_{\{u < v\}} |\nabla v|^2 + F(v) + \int_{\{u \ge v\}} |\nabla v|^2 + F(v) \\ &+ \int_{\Omega} (|x - te_n|^{-2n} F(|x - te_n|^{n-2}v) - F(v)) \\ &= J(w_2) + J(w_1) + \int_{\Omega} (|x - te_n|^{-2n} F(|x - te_n|^{n-2}v) - F(v)) \\ &\ge J(w_2) + J(w_1) + \int_{\Omega} (|x - te_n|^{-2n} F(|x - te_n|^{n-2}w_1) - F(w_1)) \\ &= J(w_2) + \tilde{J}(w_1). \end{split}$$

where the penultimate inequality is coming from (3.2). Since u and v are respective minimizers of J and  $\tilde{J}$  we conclude that  $w_1 = v$ , so that  $v \leq u$  in  $\Omega$ . We then have that  $u(x^*) \leq |x - te_n|^{2-n}u(x^*) \leq u(x)$  for  $x \in \Omega$ .

We now explain how this inequality can give directional monotonicity. Given a point  $x \in B_{1/2}$ , a direction  $\nu \in S^{n-1}$  with  $\nu \cdot e_n > 0$ , and  $\tau$  chosen small, we consider  $x + \tau \nu$ . By a translation, we perform the Kelvin transform on the ball  $B_1(te_n + y')$  for a y chosen such that  $(x + \tau \nu)^* = x$ . By choosing the appropriate value of h, then necessarily  $x \in \Omega$ , so that from the argument above we conclude  $u(x + \tau \nu) \ge u(x)$ . This will give monotonicity of u in the  $\nu$  direction. To apply this method, one must consider a specific problem to determine which values of  $t, h, \nu$  are admissible. In our specific free boundary problem, we will consider the free boundary trapped in the strip  $\{|x_n| < \epsilon\}$  for small  $\epsilon$  over a large ball  $B_R$ . By choosing  $\epsilon$  small enough, we will be able to choose t small which will allow us to reflect not only near the south pole (when t is close to 1) which will give a small cone of monotonicity, but we will also be able to reflect near the equator (when t is close to zero) and obtain a very large cone of monotonicity. As  $\epsilon$  becomes smaller, the cone of monotonicity increases, and we are able to conclude  $C^1$  regularity of the free boundary. The next two lemmas carry out this argument. We begin by showing that in our situation  $|x - te_n|^{2-n}u(x^*) \le u(x)$  on  $\partial B_1(te_n) \cap \{x_n = h\}$ .

**Lemma 3.1.** Fix  $0 < t_0 < 3/2$ , and let  $x^*$  be the reflected point via the Kelvin transform on the ball  $B_1(te_n)$  as defined in (3.1). There exists  $l_0 > 0$  depending on  $n, t_0, M, \gamma$  such that if  $0 \le x_n \le l_0$ , and  $t_0 < t < 3/2$ , and if  $x = (x', x_n) \in B_1(te_n)$ , then

$$(3.3) |x - te_n|^{2-n}h(x^* \cdot e_n) \le h(x \cdot e_n).$$

Proof. We denote  $\rho = |x'|$ . When  $\rho^2 = \rho_1^2 := 1 - (t - x_n)^2$ , so that  $x = (\rho_1, h) \in \partial B_1(te_n)$ , the inequality (3.3) is trivially satisfied since  $x = x^*$ . Also, if  $\rho^2 \leq \rho_0^2 = (t - x_n)/t - (t - x_n)^2$ , then  $u(x^*) = 0$  and (3.3) is trivially satisfied. Thus, we only need to check when  $\rho$  is in the interval  $(\rho_0, \rho_1)$ . Now  $x_n$  is fixed, and so we need

$$G(\rho) := h(x_n)|x - te_n|^{n-2} - h(x^* \cdot e_n) \ge 0.$$

Since  $x_n$  is fixed, the term  $|x - te_n|$  only depends on  $|x'| = \rho$ , so for notational convenience we define  $g(\rho) = |x - te_n|$ . Suppose now that  $G'(\rho) = 0$  so that

$$0 = G'(\rho) = h(x_n)(n-2)g^{n-3}(n-2)g' - h'(x^* \cdot e_n)2(t-x_n)g^{-3}g'.$$

Using that q' > 0 we have

$$h(x_n)g^n(n-2) = 2(t-x_n)h'(x^* \cdot e_n)$$

If we now assume that at the same  $\rho$ , that  $G(\rho) < 0$ , then

$$(n-2)g^{2}(\rho)h(x^{*} \cdot e_{n}) > h(x_{n})g^{n}(\rho)(n-2) = 2(t-x_{n})h'(x^{*} \cdot e_{n}).$$

Using (2.1) we have

$$(n-2)g^2(\rho)h(x^* \cdot e_n) \ge \frac{C(t_0 - x_n)}{x^* \cdot e_n}h(x^* \cdot e_n).$$

Taking  $x_n$  small enough (so that  $x^* \cdot e_n$  is arbitrarily small), we obtain a contradiction.

**Lemma 3.2.** Fix  $\eta > 0$ , and let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_R(0)$  with  $R = 2 + 2/\eta$ . There exists  $\epsilon_0 > 0$  depending on  $\eta$ , n such that if  $h(x_n - \epsilon) \leq u \leq h(x_n + \epsilon)$  in  $B_R(0)$ , then  $\partial_{\nu} u \geq 0$  for any  $x \in B_1$  and  $\nu \in \partial B_1$  with  $e_n \cdot \nu \geq 2\eta$ .

Proof. We first let  $\epsilon < \eta/16$ . By letting  $\epsilon \to 0$ , and by compactness and  $C^{1,\beta}$  convergence, we have for small enough  $\epsilon$  that  $\partial_{\nu}u(x) \ge 0$  for  $e_n \cdot \nu \ge 2\eta$  and  $x \in B_{R/2} \cap \{x_n \ge \eta/8\}$ . Since  $u(x', x_n) = 0$  for  $x_n \le -\epsilon$ , we also have  $\partial_{\nu}u(x', x_n) \ge 0$  if  $x_n \le -\epsilon$ . We now consider the more difficult strip where  $\{|x_n| < \epsilon\}$ , and we will employ the ideas from Lemma 3.1. For our ball  $B_1(te_n)$  we will assume that  $t \ge \eta$ . This now gives a fixed  $l_0$  coming from Lemma 3.1, and if necessary we will choose  $\epsilon$  even smaller so that  $\epsilon < l_0/4$ . Since we used a first derivative argument in Lemma 3.1 to prove (3.3), we will utilize  $C^{1,\beta}$  convergence (as  $\epsilon \to 0$ ) for  $x_n \ge l_0/2$ . Then  $u(x', x_n) \ge |(x', x_n) - te_n|^{n-2}u(x^*)$  as

long as  $t \ge \eta/4$  and  $x_n = l_0$ . Then from the argument explained at the beginning of this section, we have that  $\partial_{\nu} u(x) \ge 0$  for  $x \in B_1$  and  $\nu \cdot e_n \ge 2\eta$ .

From Lemma 3.2 and rescaling we obtain Theorem 1.1.

# 4. Lipschitz implies $C^{1,\beta}$

In this section we obtain  $C^{1,\beta}$  regularity of the free boundary near flat points. It will be convenient for the remainder of the paper to label  $\mathbb{R}^n = \mathbb{R}^{n-1} \times \mathbb{R}$  with  $(x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$ . We refer to  $\mathbb{R}^{n-1}$ as the thin space, and label  $B'_r := B_r \cap \mathbb{R}^{n-1}$ . From now on, we also assume that  $\gamma_1 > -\frac{1}{3}$ .

Based on the results of the previous section, throughout this section we make the following assumption for the free boundary of the minimizer:  $\partial \{u > 0\}$  is parametrized by  $x_n = q(x')$  with  $(0,0) \in \partial \{u > 0\}$ . From the Lipschitz regularity in the previous section we may assume that  $||q||_{C^{0,1}}(B_1) \leq \epsilon$  and that  $\partial_e u(x', x_n) \geq 0$  for any  $(x', x_n) \in B_1$  and any  $e \in S^{n-1}$  such that  $e \cdot e_n \geq \eta$  for some small  $\eta > 0$ .

By writing  $u(x', g(x', x_n)) = x_n$  and differentiating in the  $x_i$  direction, we have that

$$u_{x_i}(x', g(x', x_n)) + u_{x_n}g_{x_i}(x', x_n) = 0$$

so that

$$g_{x_i}(x', g(x', x_n)) = -\frac{u_{x_i}(x', g(x', x_n))}{u_{x_n}(x', g(x', x_n))}$$

If the quotient  $u_{x_i}/u_{x_n}$  is Hölder continuous, then we may conclude that  $g_{x_i}(x,0)$  is Hölder continuous, so that the free boundary  $\partial \{u > 0\}$  is  $C^{1,\beta}$ . Notice it is sufficient to show  $u_e/u_{x_n}$  is Hölder continuous for any e with  $e \cdot e_n \ge 1 - \delta$  for some small  $\delta$ .

Utilizing the Lipschitz nature of the free boundary, we state here improved growth rates from the free boundary.

**Lemma 4.1.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6). Then there exists a universal constant C such that for any  $(x', x_n) \in B_{1/2}$ ,

(1) 
$$u(x', x_n + q(x')) \approx h(x_n)$$
  
(2)  $u_{x_n}(x', x_n + q(x')) \approx h(x_n)/x_n$   
(3)  $|\nabla u(x', x_n + q(x'))| \leq Ch(x_n)/x_n$   
(4)  $|D^2 u(x', x_n + q(x'))| \leq Ch(x_n)/x_n^2$   
(5)  $|\nabla \frac{u_e}{u_y}(x', x_n + q(x'))| \leq \frac{C}{x_n}.$ 

*Proof.* The upper bound in (1) follows directly from optimal regularity. Using the directional monotonicity of u we have that  $u(x', x_n + q(x')) \ge u(x)$  for all  $x \in B_{x_n/2}(x')$ . Then by the nondegeneracy estimate we have

$$u(x', x_n + q(x')) \ge \sup_{B_{x_n/2}(x')} \ge ch(x_n/2) \ge c_1h(x_n).$$

This gives the lower bound in (1).

Estimates (2)-(4) then follow from rescaling and interior estimates. To obtain (5) we differentiate and utilize (2) - (4) to get

(4.1) 
$$\left|\nabla \frac{u_e}{u_{x_n}}(x', x_n + q(x'))\right| = \left|\frac{u_{x_n}\nabla u_e - u_e\nabla u_{x_n}}{u_{x_n}^2}\right| \le \frac{C}{x_n}$$

In the introduction, we noted the unavailability of the boundary Harnack principle in our problem, due to its singular nature. Instead, we opt to flatten the boundary and examine the resulting degenerate elliptic equation, following the approach outlined in [23,25]. While we could potentially employ the Hodograph transform to flatten the free boundary, as discussed in Section 5, we choose this particular method partly to elucidate why it does not extend beyond  $C^{1,\beta}$ .

We will show that  $u_e/u_{x_n}$  satisfies a degenerate elliptic equation. Subsequently, leveraging the regularity theory for degenerate elliptic equations, we can establish the Hölder continuity of  $u_e/u_{x_n}$ .

**Lemma 4.2.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6). Then

$$div\left(u_{x_n}^2 \nabla \frac{u_e}{u_{x_n}}\right) = 0$$

in the following weak sense

(4.2) 
$$\int_{B_1} u_{x_n}^2 \langle \nabla \frac{u_e}{u_{x_n}}, \nabla \psi \rangle = 0$$

for every  $\psi \in C_0^1(B_1)$ .

*Proof.* Since  $u \in C^{\infty}$  in  $\{u > 0\}$ , if u(x) > 0 we may directly compute

$$\operatorname{div}\left(u_{x_{n}}^{2}\nabla\frac{u_{e}}{u_{x_{n}}}\right) = \sum_{i=1}^{n} \frac{\partial}{\partial x_{i}} \left(u_{x_{n}}^{2}\frac{u_{x_{n}}u_{ex_{i}} - u_{e}u_{x_{n}x_{i}}}{u_{x_{n}}^{2}}\right) = \sum_{i=1}^{n} \frac{\partial}{\partial x_{i}} \left(u_{x_{n}}u_{ex_{i}} - u_{e}u_{x_{n}x_{i}}\right)$$
$$= \sum_{i=1}^{n} \left(u_{x_{n}x_{i}}u_{ex_{i}} + u_{x_{n}}u_{ex_{i}x_{i}} - u_{ex_{i}}u_{x_{n}x_{i}} - u_{e}u_{x_{n}x_{i}x_{i}}\right)$$
$$= u_{x_{n}}\Delta u_{e} - u_{e}\Delta u_{x_{n}} = u_{x_{n}}f'(u)u_{e} - u_{e}f'(u)u_{x_{n}} = 0.$$

If  $\Omega \subset \{u > 0\}$ , then using (4.1) we have

$$\int_{\Omega} u_{x_n}^2 \langle \nabla \frac{u_e}{u_{x_n}}, \nabla \psi \rangle = \int_{\partial \Omega} \psi \langle u_{x_n} \nabla u_e - u_e \nabla u_{x_n}, \zeta(x) \rangle,$$

where  $\zeta$  is the outward unit normal vector for  $\partial\Omega$ . Since we assume  $\partial\{u>0\}$  to be Lipschitz and parametrized by  $q \in C^{0,1}(B'_1)$  with  $|\nabla u(x', x_n + q(x'))| \approx h(x_n)/x_n$  and  $|D^2 u(x', x_n + q(x'))| \approx h(x_n)/x_n^2$ , it follows that  $|\nabla u||D^2 u|(x', x_n + q(x')) \leq Ch^2(x_n)/x_n^3$ . Then with (2.1)

$$\left| \int_{\Omega} u_{x_n}^2 \langle \nabla \frac{u_e}{u_{x_n}}, \nabla \psi \rangle \right| \le \frac{h^2(x_n)}{x_n^3} \le C x_n^{-3+4/(1-\gamma_1)}$$

If  $-3 + 4/(1 - \gamma_1) > 0$ , so that  $\gamma_1 > -1/3$ , then (4.2) holds.

If  $x \in \{u > 0\}$ , we define a(x) by

(4.3) 
$$u_{x_n}(x', x_n + q(x')) = a(x)h'(x_n)$$

The primary focus of this section is to smoothen the boundary, which enables the treatment of the quotient  $u_e/u_{x_n}$  as it pertains to a degenerate variable coefficient problem. Given our assumption that the free boundary is Lipschitz rather than  $C^1$ , continuity in the coefficients is not known or assumed. However, in proximity to the free boundary, these coefficients may display slight oscillations. Consequently, we will utilize Cordes-Nirenberg type estimates. From Lemma 4.1, we deduce that a(x) is bounded. Next, we demonstrate that osc  $a \to 0$  as  $|q|_{C^{0,1}} \to 0$ .

**Lemma 4.3.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6) in  $B_1$  and fix  $0 < \epsilon < 1/4$ . There exists  $\delta > 0$  such that if  $\|q\|_{C^{0,1}}(B'_{1/2}) \leq \delta$ , then

$$\operatorname{osc} a(x) \leq \epsilon$$
 whenever  $x \in B_{\delta} \cap \{x_n \geq \delta\epsilon\}$ .

*Proof.* Fix  $\epsilon > 0$ . We will utilize the directional monotonicity. Suppose by way of contradiction that there exist  $H_{\leq 1}$ -minimizers  $u_k$  in  $B_1$  with  $||q_k||_{C^{0,1}(B'_1)} \leq 1/k$  and points  $z^k, y^k \in B_{1/k} \cap \{x_n \geq \epsilon/k\}$  with  $|a_k(z^k) - a_k(y^k)| \geq \epsilon$ . We rescale with

$$\tilde{u}_k(x) := \frac{u_k(x/k)}{h_k(1/k)}, \qquad \tilde{a}_k(x) := a_k(x/k), \qquad \tilde{z}^k := z^k/k \qquad \tilde{y}^k := y^k/k$$

so that

(4.4) 
$$|\tilde{a}_k(\tilde{z}^k) - \tilde{a}_k(\tilde{y}^k)| \ge \epsilon.$$

Then we obtain a subsequence of  $\tilde{u}_k$  converging in  $C^{1,\beta}$  on compact sets to a  $H_{\leq 1}$ -minimizer  $\tilde{u}_0$ which is monotone in every direction  $\nu$  with  $\nu \cdot e_n \geq 0$ . Thus  $\tilde{u}_0$  is one-dimensional and consequently a constant multiple of the limiting function  $\tilde{h}_0$ . Now  $\tilde{a}_k$  converges uniformly on  $B_1 \cap \{x_n \geq \epsilon\}$  to  $\tilde{a}_0$  which must also be constant. Now if  $\tilde{z}^k \to z^0 \in \partial B_1$  and  $\tilde{y}^k \to y^0 \in \partial B_1$ , then we obtain a contradiction to (4.4).

In both this and the following section, we will need Sobolev estimates for the weighted measure  $\alpha(x_n) dx$  with  $\alpha(x_n) = (h'(x_n))^2$ . Throughout the remainder of the paper, whenever we state that a constant depends on  $\alpha$  we mean that the constant depends on  $M, \gamma_1, \gamma_2$  where  $f \in \mathcal{F}(M, \gamma_1, \gamma_2)$ . After flattening the boundary, we will evenly reflect our functions, so that  $u(x', x_n) = u(x', -x_n)$ . For smooth functions we consider the weighted norm

(4.5) 
$$||u||_{W^{1,p}(\alpha,\Omega)} := \left(\int_{\Omega} \alpha(x_n)(|u|^P + |Du|^P)\right)^{1/p}, \quad (\alpha(x_n) = (h'(x_n))^2).$$

Following the standard notation in the literature, we denote  $H^{1,p}(\alpha,\Omega)$  as the closure of  $C^{\infty}$  functions with the above norm. The space  $W^{1,p}(\alpha,\Omega)$  typically refers to the set of  $L^1_{\text{loc}}$  functions such that the function and weak derivative are bounded in the above norm. If  $\gamma_2 > 2$ , then  $W^{1,p}(\alpha,\Omega) \subsetneq H^{1,p}(\alpha,\Omega)$ , so we will work in the space  $H^{1,p}(\alpha,\Omega)$  which is complete. The following inequality illustrates that if  $\gamma_2 \leq 2$ , the estimates in this section would be simpler. If  $\gamma_2 \leq 2$ , then  $H^{1,2}(\alpha,\Omega) \subset L^1_{\text{loc}}(\Omega)$ . We show later in Proposition 4.5 that  $H^{1,p}(\alpha,\Omega) \subset L^1_{\text{loc}}(\Omega)$  for p large enough.

**Proposition 4.4.** Let  $v \in H_0^{1,2}(\alpha, B_1)$ , then for any  $0 < \gamma_3 \le \min\{1, 2(1 - \gamma_1)/(1 + \gamma_1)\}$ , there exists a constant C depending on  $\gamma_1, \gamma_2, \gamma_3, n$  such that<sup>2</sup>

(4.6) 
$$\int_{B_1} \frac{\alpha(x_n)}{|x_n|^{\gamma_3+1}} v^2 \le C \int_{B_1} \alpha(x_n) |\nabla v|^2$$

*Proof.* Let  $\phi \in C_0^1(B_1)$ , and extend it by zero outside  $B_1$ . For fix  $x' \in \mathbb{R}^{n-1}$  we have

$$0 = \int_{-1}^{1} \frac{d}{dt} \left[ \frac{\alpha(t)}{|t|^{\gamma_3}} \phi^2(x', t) \right] dt = \int_{-1}^{1} \left[ \frac{\alpha'(t)}{|t|^{\gamma_3}} - \gamma_3 \frac{\alpha(t)}{|t|^{\gamma_3+1}} \right] \phi^2(x', t) dt + \int_{-1}^{1} \frac{\alpha(t)}{|t|^{\gamma_3}} 2\phi(x', t) \phi_{x_n}(x', t) dt$$

Using that  $\alpha(t) = [h'(t)]^2$  and (2.1) we have

$$\frac{\alpha'(t)}{|t|^{\gamma_3}} - \gamma_3 \frac{\alpha(t)}{|t|^{\gamma_3+1}} \ge c \frac{\alpha(t)}{|t|^{\gamma_3+1}}.$$

Then using Young's inequality we obtain

$$\int_{-1}^{1} \frac{\alpha(t)}{|t|^{\gamma_3+1}} \phi^2(x',t) \le C \int_{-1}^{1} \alpha(t) \phi_{x_n}^2(x',t) \ dt + \epsilon \int_{-1}^{1} \frac{\alpha(t)}{|t|^{2\gamma_3}} \phi^2(x',t) \ dt$$

Using that  $\gamma_3 \leq 1$  we can absorb the second term on the right hand side to conclude (4.6).

We also have the following inequality for  $p \ge 2$ .

<sup>&</sup>lt;sup>2</sup>Observe that  $\gamma_1, \gamma_2$  dependent come in indirectly by the definition of  $\alpha(x_n)$  in (4.5) and Proposition 2.2.

**Proposition 4.5.** Let  $v \in H_0^{1,p}(\alpha, B_R)$ . If  $p > 2(1 + \gamma + 2)/(1 - \gamma_2)$ , then there exists C depending on  $\gamma, R$  such that

(4.7) 
$$\int_{B_R} v^p \le C \int_{B_R} |x_n|^{2(1+\gamma_2)/(1-\gamma_2)} |\nabla v|^p \le \int_{B_R} \alpha(x_n) |\nabla v|^p.$$

*Proof.* Fix  $x' \in \mathbb{R}^{n-1}$ . By density we need only prove the result for  $\phi \in C_0^1(B_R)$ . We will integate over  $B_R^+$ , and the result will be the same for  $B_R^-$ . We have

$$\begin{split} \phi^p(x',r) &= \left(\int_r^R \phi_{x_n}(x',x) \ ds\right)^p \\ &= \left(\int_r^R s^{-\tau} \phi_{x_n}(x',s) s^{\tau} \ ds\right)^p \\ &\leq \left(\int_r^R s^{-\tau p/(p-1)}\right)^{p-1} \left(\int_r^R s^{\tau p} \phi_{x_n}^p(x',s)\right) \\ &\leq Cr^{p(1-\tau)} \left(\int_r^R s^{\tau p} \phi_{x_n}^p(x',s) \ ds\right) \\ &\leq Cr^{p(1-\tau)} \left(\int_0^R s^{\tau p} \phi_{x_n}^p(x',s) \ ds\right) \end{split}$$

If  $p > 2(1 + \gamma)/(1 - \gamma)$ , then we choose  $\tau$  so that  $p\tau = 2(1 + \gamma_2)/(1 - \gamma_2)$ . Then  $\tau < 1$ , so that we may integrate in r and then x' to obtain (4.7).

We also have the following weighted Sobolev inequality.

**Proposition 4.6.** Let  $v \in H_0^{1,2}(\alpha, B_1)$ . There exists C and p > 2 both depending on  $n, \gamma_1, \gamma_2$  such that<sup>3</sup>

$$\|v\|_{L^p(\alpha, B_1)} \le C \|\nabla v\|_{L^2(\alpha, B_1)}.$$

*Proof.* By density we need only prove the result for  $\phi \in C_0^1(B_1)$ . We will adapt the usual proof of the Sobolev inequality in the subcritical case. For  $y \in \Gamma := \{y \in \partial B_2 : e_n \cdot y > 1/4\}$ , we have  $\phi(x+2y) = 0$ , so that

$$(4.8) -c(n)\phi(x) = -\int_{\Gamma} \phi(x) \, d\sigma(y) = \int_{\Gamma} \phi(x+2y) - \phi(x) \, d\sigma(y) = \int_{\Gamma} \int_{0}^{2} \langle \nabla \phi(x+ty), y \rangle \, dt \, d\sigma(y).$$

Using Hölder's inequality on the right side on the product space  $\Gamma \times (0,2)$ , we have

$$-c(n)\phi(x) \le \left(\int_{\Gamma} \int_{0}^{2} \frac{1}{\alpha(x_{n} + ty_{n})} \ dt \ d\sigma(y)\right)^{1/2} \left(\int_{\Gamma} \int_{0}^{2} |\nabla\phi(x + ty)|^{2} \alpha(x_{n} + ty_{n}) \ dt \ d\sigma(y)\right)^{1/2}.$$

We now square both sides of the above inequality and multiply both sides by  $\alpha(x_n)^{1/q}$  to obtain

 $^{3}$ Recall footnote 2

Now since  $\alpha$  is an increasing function, we have that  $\alpha(x_n) \leq \alpha(x_n + ty_n)$  for  $y_n \geq 0$ , so that

$$\begin{aligned} \alpha(x_n)^{1/q} \int_{\Gamma} \int_{0}^{2} \frac{1}{\alpha(x_n + ty_n)} \, dt \, d\sigma(y) &\leq \int_{\Gamma} \int_{0}^{2} \frac{\alpha(x_n + ty_n)^{1/q}}{\alpha(x_n + ty_n)} \, dt \, d\sigma(y) \\ &= \int_{\Gamma} \int_{0}^{2} \alpha(x_n + ty_n)^{1/q - 1} \, dt \, d\sigma(y) \\ &\leq \int_{\Gamma} \int_{0}^{2} \alpha(ty_n)^{1/q - 1} \, dt \, d\sigma(y) \\ &\leq C \int_{\Gamma} \int_{0}^{2} t^{2(1 - \gamma_2)/(1 + \gamma_2)(1 - 1/q)} \, dt \, d\sigma(y) \quad \text{by (2.1)} \\ &\leq C(n, \gamma_2, q) \end{aligned}$$

as long as q > 1 is chosen close enough to 1. Then continuing with the usual proof of the Sobolev inequality in the subcritical case, we have

$$\alpha(x_n)^{1/q} |u(x)|^2 \le C(\alpha |\nabla u|^2 * \chi_{B_2} |x|^{1-n})$$

Applying Young's convolution theorem, we conclude

$$\|\alpha(x_n)^{1/q}\|u(x)\|^2\|_{L^q} \le C\|\alpha(x_n)|\nabla u|^2\|_{L^1}\|\chi_{B_2}|x|^{1-n}\|_{L^q}.$$

By taking the square root of both sides we obtain the result with p = 2q.

In the Hilbert space setting, the following result is immediate.

**Lemma 4.7.** Assume that  $\lambda I \leq A \leq \Lambda I$  and let  $g^i \in L^2(\alpha, B_r)$  for  $0 \leq i \leq n$  and label  $G = (g^1, \ldots, g^n)$ . Let  $\psi \in H^{1,2}(\alpha, B_r)$ , then there exists a unique w with  $w - \psi \in H^{1,2}_0(\alpha, B_r)$  such that

(4.9) 
$$\int_{B_r} \alpha(x_n) \langle A \nabla w, \nabla v \rangle = \int_{B_r} \alpha(x_n) g^0 v - \langle \boldsymbol{G}, \nabla w \rangle.$$

for all  $v \in W_0^{1,2}(\alpha, B_r)$ .

Finally, we state the Caccioppoli inequality which is obtained with the usual proof.

**Proposition 4.8.** If w is a subsolution in the sense that

$$\int_{B_1} \alpha(x_n) \langle A \nabla w, \nabla v \rangle \le 0 \quad \text{for all } v \in H_0^{1,2}(B_1) \text{ with } v \ge 0.$$

then there exists a constant C depending on dimension  $n, \lambda, \Lambda$  such that for  $0 < r < R \leq 1$ 

$$\int_{B_r} \alpha(x_n) |\nabla w|^2 \le \frac{C}{(R-r)^2} \int_{B_R} \alpha(x_n) w^2.$$

We now flatten the boundary. If  $\partial \{u > 0\}$  is parametrized by  $x_n = q(x')$ , then we flatten  $\partial \{u > 0\}$  by  $\Phi(x', x_n) = (x', x_n - q(x))$  and let  $\Phi^{-1} = \Psi(x', x_n) = (x', x_n + q(x'))$ . We define

(4.10) 
$$w(x,y) = \frac{u_e}{u_{x_n}}(\Psi(x',x_n)) = \frac{u_e}{u_{x_n}}(x',q(x')+x_n).$$

We have that  $|\det \Psi| = 1$ . We then have that w satisfies

(4.11) 
$$\operatorname{div}\left((h'(x_n)a(x))^2a^{ij}\nabla w\right) = 0,$$

where

$$a^{ij} = \delta_{ij} \quad \text{for } 1 \le i, j < n,$$
  

$$a^{in} = a^{ni} = -q_{x_i} \quad \text{for } 1 \le 1 < n,$$
  

$$a^{nn} = 1 + |\nabla q|^2.$$

The matrix  $a^{ij}$  is uniformly elliptic and continuous. From Lemma 4.3 we have that a(x) is almost constant and uniformly bounded from above and away from zero.

We will need estimates on the limiting solution.

**Lemma 4.9.** If w is a solution to (4.9) in  $B_1$  with  $g^i \equiv 0$  for all i and  $A \equiv I$ , then there exists a  $C, \beta$  depending on  $\alpha, n$  such that

(4.12) 
$$\|\partial_{\nu}w\|_{C^{2,\beta}(B_{1/2})} \le C\|w\|_{L^{2}(\alpha,B_{1})},$$

whenever  $\nu \cdot e_n = 0$ . Furthermore,  $w_{x_n}(x', 0) = 0$  and

$$\|w_{x_n}\|_{C^{0,1}(B_{1/2})} \le C \|w\|_{L^2(\alpha, B_1)}$$

*Proof.* We first utilize the fact that our weight  $\alpha$  is independent of the first n-1 variables, to show derivatives in the first n-1 variables are also solutions. If e is a unit vector orthogonal to  $x_n$ , and we consider the difference quotient

$$\phi_{e,\tau}(x',x_n) := \frac{\phi(x'+e,x_n) - \phi(x',x_n)}{\tau},$$

for a smooth function  $\phi$ , then

$$\phi_{e,\tau}(x',x_n) = \frac{1}{\tau} \int_0^\tau \langle \nabla \phi(x'+\tau,x_n), e \rangle,$$

so that if r < 1 - h, then

$$\begin{split} \int_{B_r} \alpha(x_n) |\phi_{e,\tau}(x',x_n)|^2 &\leq \int_{B_r} \alpha(x_n) \left(\frac{1}{\tau} \int_0^\tau |\nabla \phi(x'+t,x_n)| \ dt\right)^2 \\ &\leq \frac{1}{\tau} \int_0^\tau \int_{B_1} \alpha(x_n) |\nabla \phi(x',x_n)|^2 \ dt \\ &= \int_{B_1} \alpha(x_n) |\nabla \phi(x',x_n)|^2. \end{split}$$

Thus, by density of  $C^{\infty}$  in  $H^{1,2}(\alpha, B_1)$  we have that for small enough  $\tau$ ,

(4.14) 
$$\int_{B_r} \alpha(x_n) w_{e,\tau}^2 \le \int_{B_1} \alpha(x_n) |\nabla w|^2.$$

By linearity,  $w_{e,\tau}$  is a solution to (4.9). Applying the Caccioppoli inequality we have that

$$\int_{B_{r_1}} \alpha(x_n) |\nabla w_{e,\tau}|^2 \le C \int_{B_{r_2}} \alpha(x_n) |w_{e,\tau}|^2 \le C \int_{B_{r_3}} \alpha(x_n) |\nabla w|^2 \le C \int_{B_{r_4}} \alpha(x_n) w^2 + C \int_{B_{r_4}} \alpha(x_n) |\nabla w|^2 \le C \int_{B_{r_4}} \alpha(x_n) |w_{e,\tau}|^2 \le C \int_{B_{r_4}} \alpha(x_n) |\nabla w|^2 \le C \int_{B_{r_4}} \alpha(x_n)$$

Thus,  $w_e \in H^{1,2}(\alpha, B_{r_1})$ . By iterating this procedure we obtain  $D^{\beta}u \in H^{1,2}(\alpha, B_{1/2})$  as long as the multi-index  $\beta$  is only in the first n-1 coordinates. By utilizing the Sobolev inequality in Proposition 4.6 and the Caccioppoli inequality in Proposition 4.8, the standard Moser iteration technique for subsolutions gives

(4.15) 
$$\lim_{p \to \infty} \left( \int_{B_{3/4}} \alpha(x_n) (D^{\beta} w)^p \right)^{1/p} \le C \int_{B_{7/8}} \alpha(x_n) (D^{\beta} w)^2.$$

The standard proof shows that this implies

$$||D^{\beta}w||_{L^{\infty}(B_{3/4})} \le C \int_{B_{7/8}} \alpha(x_n) (D^{\beta}w)^2.$$

Since  $D^{\beta}w$  is bounded, we may repeatedly apply the *unweighted* Sobolev embedding theorem finitely many times on each n-1 dimensional, so that for  $|x_n| \leq 1/4$ 

$$||w(\cdot, x_n)||_{C^{2,\beta}(B'_{1/4})} \le ||w||_{L^{\infty}(B_{1/2})}.$$

To obtain regularity of  $D^{\beta}w$  as  $x_n$  changes as well as  $w_{x_n}$ , we now consider regularity in the  $x_n$  direction. From (4.12) we have that  $(\alpha(x_n)u_{x_n})_{x_n} = \alpha(x_n)\Delta_{x'} \in C^{0,\beta}(B'(\cdot, x_n))$ . Since w is even and bounded (from (4.15)), the weak formulation (4.9) implies that  $\lim_{x_n\to 0} \alpha(x_n)w(x', x_n) = 0$  for  $H^{n-1}$  almost every x'. Then for  $x' \in B_{1/4}$ 

$$\begin{aligned} \alpha(x_n)w_{x_n}(x',x_n) &= \alpha(x_n)w_{x_n}(x',x_n) - 0 = \int_0^{x_n} \partial_{x_n}(\alpha(y)w_{x_n}(x',y)) \, dy \\ &= \int_0^{x_n} \alpha(y)\Delta_{x'}w(x',y) \, dy \le C \int_0^{x_n} \alpha(y) \|w\|_{L^{\infty}(B_{1/4})}. \end{aligned}$$

Now

$$\frac{d}{dt}\alpha(t) = \frac{d}{dt}(h'(t))^2 = 2h'(t)h''(t) = 2\sqrt{2F(h(t))}f(h(t)) \ge 0$$

Then we conclude

$$\begin{aligned} |\alpha(x_n)w_{x_n}(x',x_n)| &\leq \int_0^{x_n} \alpha(y) |\Delta_{x'}w(x',y)| \ dy \leq C \|w\|_{L^{\infty}(B_{1/2})} \int_0^{x_n} \alpha(y) \ dy \\ &\leq C \|w\|_{L^{\infty}(B_{1/2})} \int_0^{x_n} \alpha(x_n) \ dy = C \|w\|_{L^2(\alpha,B_1)} x_n \alpha(x_n), \end{aligned}$$

so that

$$|w_{x_n}(x', x_n)| \le C x_n ||w|| L^2(\alpha, B_1).$$

This proves the Lipschitz estimate of  $w_{x_n}$  on the thin space; the estimate off the thin space follows from interior regularity and scaling, and we have (4.13). Finally, having shown  $w_{x_n}$  is Lipschitz, (so in particular  $D^{\beta}w_{x_n}$  is Lipschitz), we conclude (4.12).

We now proceed to obtain Cordes-Nirenberg type estimates.

**Lemma 4.10.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6), and let w be given as in (4.10), so that  $w(x, y) = \frac{u_e}{u_{x_n}}(x', q(x') + x_n)$ . For fixed  $\beta < 1$  and r < 1/2, there exists a constant  $C(\beta)$  and  $\epsilon > 0$  such that if  $\|q\|_{C^{0,1}(B_1)} \leq \epsilon$  then

(4.16) 
$$S_r(x,w) := \frac{1}{\mu(\alpha, B_r)} \int_{B_r(x)} \alpha(x_n) (w - \overline{w}_r(x))^2 \le C(\alpha) r^{2\beta},$$

for any  $x = (x', 0) \in B'_{1/2}$  and where

$$\overline{w}_r(x) = \frac{1}{\mu(\alpha, B_r)} \int_{B_r(x)} \alpha(x_n) w \quad and \quad \mu(\alpha, B_r) := \int_{B_R} \alpha(x_n) \, dx.$$

*Proof.* Fix  $0 < \beta < 1$ . Suppose by way of contradiction that (4.16) is not true. Then there exists a sequence  $\epsilon_j \to 0$ , solutions  $w_j$ , radii  $r_j \to 0$  and points  $x_j \in B'_{1/2}$  (which we may assume by taking a subsequence that  $x_j \to x_0$ ) such that

(1) 
$$\frac{S_{r_j}(x_j, w_j)}{r_j^{2\beta}} = C_j \to \infty$$
  
(2) 
$$S_{2^{-k}r_j}(x_j, w_j) \ge 2^{-2\beta k} S_{r_j}(x_j, w_j) \quad \text{whenever } 1 \le k \le j.$$

We now rescale with

$$\tilde{w}_j(x) := \frac{w(r_j/2x + x_j) - \overline{w}_{r_j/2}(x_j)}{\sqrt{S_{r_j/2}(x_j, w_j)}}.$$

We will use  $\tilde{\alpha}_j(x_n) := \alpha_j(r_j x_n/2)/\alpha_j(r_j/2)$ . From the Caccioppoli inequality we have that

$$\int_{B_{3/2}} \tilde{\alpha}_j(x_n) |\nabla \tilde{w}_j|^2 \le C \int_{B_2} \tilde{\alpha}_j(x_n) |\tilde{w}_j|^2.$$

19

By the Arzela-Ascoli theorem, taking a subsequence we obtain  $\tilde{\alpha}_j \to \tilde{\alpha}_0$  in  $C^1([0,2])$ . We also have a limiting function  $\tilde{w}_0$  with the convergence in  $C^2$  away from the thin space. From the Caccioppoli inequality above  $\sqrt{\tilde{\alpha}_j} \nabla \tilde{w}_{r_j} \to \sqrt{\tilde{\alpha}_0} \nabla \tilde{w}_0$  in  $L^2(B_1)$ . If  $v \in C_0^1$ , then  $\sqrt{\tilde{\alpha}_j} \nabla v A_j \to \sqrt{\tilde{\alpha}_0} \nabla v$  in  $L^2$ . Thus,  $w_0$  is a solution to

$$\int_{B_1} \tilde{\alpha}_0(x_n) \langle A_0 \nabla \tilde{w}_0, \nabla v \rangle = 0,$$

for all smooth  $v \in C_0^1(B_1)$ , and where  $A_0$  is the limiting coefficient matrix. Also, from the definition of the rescaling we have that  $S_1(0, \tilde{w}_j) = 1$ . We now show that  $\tilde{w}_j \to \tilde{w}_0$  in  $L^2(\alpha, B_1)$ , so that the limiting function satisfies  $S_1(0, \tilde{w}_0) = 1$ . From the weighted Poincare inequality in Proposition 4.5 we have

$$\int_{B_{1}\cap|x_{n}|\leq t} \tilde{\alpha}_{j}(x_{n})|\tilde{w}_{j}|^{2} \leq t^{1+\gamma_{3}} \int_{B_{1}\cap|x_{n}|\leq t} \frac{\tilde{\alpha}_{j}}{x_{n}^{1+\gamma_{3}}} |\tilde{w}_{j}|^{2} \leq \int_{B_{3/2}} \tilde{\alpha}_{j}(x_{n})|\nabla \tilde{w}_{j}|^{2} \\ \leq \int_{B_{2}} \tilde{\alpha}_{j}(x_{n})|\tilde{w}_{j}|^{2} \leq \int_{B_{1}} \tilde{\alpha}_{j}(x_{n})|\tilde{w}_{j}|^{2} \leq C,$$

with the penultimate inequality coming from how j is chosen and the final inequality is due to how  $\tilde{w}_j$  is defined. Thus,

$$\int_{B_1 \cap \{|x_n| \le t\}} |\tilde{w}_j|^2 \le C t^{1+\gamma_3},$$

so  $\tilde{w}_j$  cannot concentrate on the thin space, and so  $\tilde{\alpha}_j \tilde{w}_j \to \tilde{\alpha}_0 \tilde{w}_0$  in  $L^2(\alpha, B_1)$ , and we conclude  $S_1(0, \tilde{w}_0) = 1$ .

Now from 4.3 we have that  $A_0$  is a constant multiple of the identity for  $|x_n| \ge \epsilon$ , but  $\epsilon$  can be taken to be arbitrary small. Thus the limiting function  $\tilde{w}_0$  is an even weak solution to  $\operatorname{div}(\tilde{\alpha}_0 \nabla \tilde{w}_0) = 0$ . From the Lipschitz regularity of  $\tilde{w}_0$  from Lemma 4.9, we have

(4.17) 
$$S_{2^{-k}}(0, \tilde{w}_0) \le C_1 2^{-2k} S_1(0, \tilde{w}) \quad \text{for } k \ge 1.$$

From our assumption (2) we have

(4.18) 
$$S_{2^{-k}}(0,\tilde{w}_j) \ge C_2 2^{-2k\beta} S_1(0,\tilde{w}_j) \quad \text{for } 1 \le k \le j.$$

From the  $L^2$  convergence we have for a fixed k that  $S_{2^{-k}}(0, \tilde{w}_j) \to S_{2^{-k}}(0, \tilde{w}_0)$ . Then for a fixed k large enough we obtain a contradiction between (4.17) and (4.18).

We conclude this section by stating a theorem whose proof is standard and hence left to readers.

**Theorem 4.11.** Let u be a  $H_{\leq 1}$ -minimizer of (1.6), and let w be given as in (4.10), so that  $w(x,y) = \frac{u_e}{u_{x_n}}(x',q(x')+x_n)$ . For fixed  $\beta < 1$  and r < 1/2, there exists a constant  $C(\beta)$  and  $\epsilon > 0$  such that if  $\|q\|_{C^{0,1}(B_1)} \leq \epsilon$  then  $w \in C^{0,\beta}(B_{1/2})$ .

From Theorem 4.11 and the discussion at the beginning of this section, we obtain Theorem 1.2.

# 5. Hodograph Transform

From the results of the previous section, we consider a  $H_{\leq 1}$ -minimizer u of (1.6) on  $B_r$  that is monotone in the  $e_n$  direction and has  $\partial \{u > 0\}$  a  $C^{1,\alpha}$  hypersurface containing 0 and with inward unit normal  $e_n$  at 0. To obtain higher regularity we perform a modified hodograph transform with respect to h by defining, for  $y = (y', y_n) \in B_s^+ = \{(y', y_n) \in B_s : y_n > 0\}$ , the quantity v(y) to be the unique number such that

(5.1) 
$$u(y', v(y', y_n)) = h(y_n).$$

The mapping  $y \mapsto (y', v(y', y_n))$  is a homeomorphism from  $B_s^+$  to  $U \cap \{u > 0\}$ , where U a neighborhood of 0, and extends continuously up to the boundary, mapping  $\{y_n = 0\}$  to the free boundary  $\partial\{u > 0\}$ .

As u and h are  $C^2$  functions away from  $\partial \{u > 0\}$ , v is  $C^2$  away from  $\{y_n = 0\}$ . We can therefore repeatedly differentiate the equation to get the relations

$$u_{n}v_{n} = h'(y_{n})$$
  

$$u_{i} + u_{n}v_{i} = 0$$
  

$$u_{nn}v_{n}^{2} + u_{n}v_{nn} = h''(y_{n})$$
  

$$u_{in}v_{n} + u_{nn}v_{i}v_{n} + u_{n}v_{in} = 0$$
  

$$u_{ii} + 2u_{ni}v_{i} + u_{nn}v_{i}^{2} + u_{n}v_{ii} = 0.$$

Here and below, u and its derivatives are always evaluated at (y', v(y)), v and its derivatives are evaluated at  $y \in B_s$ , and i is any number  $1, \ldots, n-1$  corresponding to a tangential direction. Summing the fifth equation over i, adding a multiple of the third equation, and making substitutions for the mixed and lower-order derivatives of u, we obtain

$$\Delta u = [1 + \sum_{i} v_i^2] u_{nn} + \sum_{i} [2v_{in} \frac{h'v_i}{v_n^2} - v_{ii} \frac{h'}{v_n}] = f(h)$$

It is helpful to multiply this whole expression by h', and then eliminate the  $u_{nn}$  factor. After combining terms, this leads to the following second-order PDE for v:

(5.2) 
$$\left[1 + \sum_{i} v_{i}^{2}\right] \left[\frac{h''h'}{v_{n}^{2}} - v_{nn}\frac{(h')^{2}}{v_{n}^{3}}\right] + \sum_{i} \left[2v_{in}\frac{(h')^{2}v_{i}}{v_{n}^{2}} - v_{ii}\frac{(h')^{2}}{v_{n}}\right] = h'f(h)$$

We claim this can be rewritten in divergence form. To see this, first observe that

$$\frac{h''h'}{v_n^2} - v_{nn}\frac{(h')^2}{v_n^3} = [\frac{(h')^2}{2v_n^2}]_n$$

and so

$$[1 + \sum_{i} v_{i}^{2}][\frac{h''h'}{v_{n}^{2}} - v_{nn}\frac{(h')^{2}}{v_{n}^{3}}] = [(1 + \sum_{i} v_{i}^{2})\frac{(h')^{2}}{2v_{n}^{2}}]_{n} - \sum_{i} v_{in}\frac{(h')^{2}v_{i}}{v_{n}^{2}}.$$

Substituting into (5.2), we get

$$[(1+\sum_{i}v_{i}^{2})\frac{(h')^{2}}{2v_{n}^{2}}]_{n} + \sum_{i}[v_{in}\frac{(h')^{2}v_{i}}{v_{n}^{2}} - v_{ii}\frac{(h')^{2}}{v_{n}}] = h'f(h).$$

The remaining term can be rewritten as a tangential derivative:

$$v_{in}\frac{(h')^2v_i}{v_n^2} - v_{ii}\frac{(h')^2}{v_n} = -[\frac{(h')^2v_i}{v_n}]_i,$$

as  $h' = h'(y_n)$  has  $h'_i = 0$ . This leads to

(5.3) 
$$\left[ (1 + \sum_{i} v_i^2) \frac{(h')^2}{2v_n^2} \right]_n - \sum_{i} \left[ \frac{(h')^2 v_i}{v_n} \right]_i = h' f(h).$$

We rewrite this as

$$\sum_{k} \partial_k \boldsymbol{H}^k(y_n, \nabla v) = q(y_n),$$

where a key point is that  $q(y_n) = h'f(h)$  is independent of  $y_i$ , for  $i \neq n$ .

Set  $w = v_i$ : this satisfies (in the weak sense, using that  $v \in C^2$ ) the PDE

$$\sum_{k,j} \partial_k [\boldsymbol{H}_j^k(y_n, \nabla v) w_j] = 0,$$

as  $q_i = 0$  for  $i \neq n$ . The matrix  $\boldsymbol{H}_j^k$  can be computed directly,

$$\boldsymbol{H}_{j}^{k}(t,p) = (h')^{2} \begin{bmatrix} \frac{1}{p_{n}} & 0 & \cdots & 0 & -\frac{p_{1}}{p_{n}^{2}} \\ 0 & \frac{1}{p_{n}} & \cdots & 0 & -\frac{p_{2}}{p_{n}^{2}} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & \frac{1}{p_{n}} & -\frac{p_{n-1}}{p_{n}^{2}} \\ -\frac{p_{1}}{p_{n}^{2}} & -\frac{p_{2}}{p_{n}^{2}} & \cdots & -\frac{p_{n-1}}{p_{n}^{2}} & \frac{1}{p_{n}^{3}}(1+\sum_{i}p_{i}^{2}), \end{bmatrix},$$

and observe that this is symmetric and of the form  $(h')^2 A(\nabla v)$ . We claim that A is uniformly elliptic on  $B_s^+$ , as long as s is chosen sufficiently small.

Indeed, recall that

$$v_n = \frac{h'}{u_n}$$
  $v_i = \frac{u_i}{u_n}$ 

From Lemma 4.1(1), we have that

$$C^{-1}h(v(y) - v(y', 0)) \le u(y', v(y)) = h(y_n) \le Ch(v(y) - v(y', 0))$$

and combining with Proposition 2.2 leads to

$$C_*^{-1}[v(y) - v(y', 0)] \le h^{-1}(C^{-1}h(v(y) - v(y', 0))) \le y_n \le h^{-1}(Ch(v(y) - v(y', 0))) \le C_*[v(y) - v(y', 0)].$$

But then from Lemma 4.1(2),

$$u_n(y', v(y)) \le C \frac{h(v(y) - v(y', 0))}{v(y) - v(y', 0)} \le C \frac{h(y_n)}{y_n},$$

 $\mathbf{SO}$ 

$$v_n(y) \ge C^{-1} \frac{y_n h'(y_n)}{h(y_n)} \ge C^{-1}$$

from Proposition 2.2 again. Similarly,

 $v_n(y) \leq C.$ 

On the other hand, from Theorem 4.11, we have that  $x \mapsto \frac{u_i(x)}{u_n(x)}$  is continuous on  $\overline{\{u > 0\} \cap B_r}$ , and coincides with the *i* derivative of a level set of *u* viewed as a graph over  $\mathbb{R}^{n-1}$ . In particular,  $v_i(0) = \frac{u_i(0)}{u_n(0)} = 0$ , and so  $|v_i| \leq \delta \ll 1$  on  $B_s^+$  for *s* chosen small enough.

It follows that all the diagonal entries of A are in  $[C^{-1}, C]$  with C depending only on the M in the hypotheses on f, while the off-diagonal entries are bounded by  $\delta C$  for any  $\delta > 0$ . Choosing s (and so  $\delta$ ) small enough, A is uniformly elliptic on  $B_s^+$ . We have established:

**Proposition 5.1.** Let  $v : \bar{B}_s^+ \to \mathbb{R}$  be as defined above, and  $w = v_i$ . Then there exists an s' > 0 small such that for any  $\phi \in C_c^{\infty}(B_s^+)$ ,

$$\int (h'(y_n))^2 A(\nabla v) \nabla w \cdot \nabla \phi = 0.$$

and moreover  $\lambda I \leq A(\nabla v) \leq \Lambda I$  for some  $0 < \lambda \leq \Lambda < \infty$  depending only on n and M.

To utilize this, we also need a boundary condition on w, that surprisingly is of a simple Neumann type. Extend the function v (and w, A) to all of  $B_s$  through even reflection, such that  $v(y', -y_n) = v(y', y_n)$ .

**Proposition 5.2.** Proposition 5.1 is valid for any  $\phi \in C_c^{\infty}(B_s)$  as well, under the same assumptions.

*Proof.* From above, we have that  $C^{-1} \leq v_n \leq C$  and  $|v_i| \leq C\delta$  on  $B_s^+$ , so v is a bilipchitz map from  $\bar{B}_s^+$  onto its image, and w is a  $C^1$  function on  $B_s^+$ . Take any  $\phi \in C_c^{\infty}(B_s)$ , and consider

$$(5.4)$$
$$\int_{B_s^+} (h'(y_n))^2 A \nabla w \cdot \nabla \phi = \lim_{t \searrow 0} \int_{B_s^+ \cap y_n > t} (h'(y_n))^2 A \nabla w \cdot \nabla \phi = -\lim_{t \searrow 0} \int_{B_s^+ \cap \{y_n = t\}} (h'(t))^2 \phi A \nabla w \cdot e_n,$$

where the last expression is obtained by using integration by parts and Proposition 5.1. Now using the expressions for the derivatives  $v_{in}$  and  $v_{ii}$  and 4.1(2,4) we have

$$|\nabla w(y)| \le C \frac{|D^2 u(y', v(y))|}{u_n(y', v(y))} \le \frac{C}{v(y) - v(y', 0)} \le \frac{C}{y_n},$$

which gives the following estimate for the integrand in (5.4)

$$|(h'(t))^2 \phi A \nabla w \cdot e_n| \le C(h'(t))^2 |\nabla w(y',t)| \le \frac{C}{t^3} h^2(t) \le C \frac{t^{2/(1+\gamma)}}{t^3} = C t^{-\frac{1+3\gamma}{1+\gamma}}.$$

For  $\gamma < -\frac{1}{3}$  this expression tends to 0 uniformly, implying

$$\int_{B_s^+} (h'(y_n))^2 A \nabla w \cdot \nabla \phi = 0$$

The integral on the other half-ball similarly vanishes.

We are now in a position to prove the following regularity theorem:

**Theorem 5.3.** Let v be as in Proposition 5.1. Then for each  $k \in \mathbb{N}$ , there is an s = s(k) > 0 such that  $v(\cdot, x_n) \in C^k(B'_s)$  for each  $x_n$ , with

$$\sup_{B_s} \sum_{|\alpha| \le k} |\partial^{\alpha} v| \le C(k).$$

where the sum is over all multiindeces  $\alpha$  in the first n-1 variables only, i.e.  $\alpha \in \mathbb{Z}_{\geq 0}^{n-1} \times \{0\}$ . In particular, the level set  $\partial \{u > 0\}$  is  $C^{\infty}$  on a neighborhood of 0.

*Proof.* First, we observe that  $\nabla v$  is continuous at 0. Indeed, we have already seen that  $v_i$  is a continuous function from Theorem 4.11. As for  $v_n$ , this follows directly from Lemma 4.3 and the fact that  $\partial \{u > 0\}$  is  $C^1$ . In particular, this implies that  $y \mapsto A(\nabla v(y))$  is continuous at 0.

We will prove, by induction on k, the following estimate: for all  $2 , there is a constant <math>C_{p,k}$  and a number s = s(p,k) such that

$$\sum_{\alpha|=k} \|\partial^{\alpha} \nabla v\|_{L^p(B_s)} \le C_{p,k}.$$

The conclusion then follows from Sobolev embeddings and the observation that  $v|_{y_n=0}$  is precisely  $\partial \{u > 0\}$  parametrized as a graph over  $\mathbb{R}^{n-1}$ .

The case of k = 0 has already been established, so consider some  $k \ge 1$ . Fix  $i \le n - 1$ , and set  $w = v_i$ , which we know satisfies

div 
$$\left[(h'(y_n))^2 A(\nabla v) \nabla w\right] = 0$$

in the weak sense. Now take a multiindex  $|\alpha| = k - 1$  containing only tangential derivatives, and differentiate the equation:

div 
$$[(h'(y_n))^2 A(\nabla v) \nabla \partial^{\alpha} w] = -\text{div} \left[ (h'(y_n))^2 \sum_{\beta_1 + \beta_2 = \alpha, |\beta_1| > 0} \partial^{\beta_1} [A(\nabla v)] \partial^{\beta_2} w \right].$$

The quantity  $\partial^{\beta_1}[A(\nabla v)]$  can be expressed (if  $\partial^{\beta_1} = \partial^i \partial^{\beta_3}$ ) as

$$\partial^{\beta_1}[A(\nabla v)] = \partial^{\beta_3} \sum_{j \le n} [A_j(\nabla v)v_i j].$$

The function  $p \mapsto A(p)$  is analytic, and so all the derivatives  $(D^m A)(\nabla v)$  are bounded over  $B_s$ . By applying this computation repeatedly, we see that  $\partial^{\beta_1}[A(\nabla v)]$  is a sum of products of derivatives of v, multiplied by one derivative in the tensor  $(D^m A)(\nabla v)$ , with the following properties: at most  $|\beta_1|$  factors are present, and each factor has at most one derivative of v is in a direction other than in the multiindex  $\beta_1$ . In particular, we can bound

$$|\partial^{\beta_1}[A(\nabla v)]| \le C(|\beta_1|) \sum_{\zeta_1 + \ldots + \zeta_m = \beta_1, |\zeta_j| \ge 1} \prod_{j=1}^m |\partial^{\zeta_j} \nabla v|.$$

Fix p > 2, and estimate

$$\begin{aligned} \|\partial^{\beta_1}[A(\nabla v)]\partial^{\beta_2}w\|_{L^p(B_s)} &\leq \|\partial^{\beta_1}[A(\nabla v)]\|_{L^{2p}(B_s)} \|\partial^{\beta_2}w\|_{L^{2p}(B_s)} \\ &\leq C_{2p,k-2}\|\partial^{\beta_1}[A(\nabla v)]\|_{L^{2p}(B_s)} \\ &\leq C\sum_{\zeta_1+\ldots+\zeta_m=\beta_1,|\zeta_j|\geq 1} \prod \|\partial^{\zeta_j}\nabla v\|_{L^{2mp}} \\ &\leq C\prod_{j=1}^m C_{2mp,|\zeta_j|} \\ &\leq C\end{aligned}$$

from our inductive hypothesis used repeatedly, so long as  $s \leq s(2kp, k-2)$ . We have shown that

div 
$$[(h'(y_n))^2 A(\nabla v) \nabla \partial^{\alpha} w] = \text{div } \boldsymbol{G}$$

with  $\|\boldsymbol{G}\|_{L^p(B_s)} \leq C$ . Applying Theorem 6.1 and using that A is continuous at 0, this gives (with possibly smaller s) that

$$\|\nabla \partial^{\alpha} w\|_{L^{p}(B_{s/2})} \le C \|\boldsymbol{G}\|_{L^{p}(B_{s})} \le C.$$

This is true for any tangential multiindex  $\alpha$  with  $|\alpha| \leq k-1$  and any  $i = 1, \ldots, n-1$ , so for any  $\alpha' = \alpha + e_i$ 

$$\|\nabla \partial^{\alpha'} v\|_{L^p(B_{s/2})} \le C.$$

Summing over  $\alpha'$  completes the inductive step.

# 6. $W^{1,p}$ estimates

In this section we show how the technique of Caffarelli-Peral [8] can be adapted to give  $C^{\infty}$  regularity of the free boundary. We will adapt the presentation given in [5] which accommodates a nonhomogeneous equation. The estimates needed are

- Suitable  $W^{1,p}$  estimates for an approximating equation
- Hardy-Littlewood maximal function bounds in a measure space with a doubling measure.

We consider even solutions  $w \in H^{1,2}(\alpha, \Omega)$  over a bounded domain U to

(6.1) 
$$\int_{\Omega} \langle \alpha(x_n) A(x', x_n) \nabla w, \nabla v \rangle = -\int_{\Omega} \alpha(x_n) \langle \boldsymbol{G}, \nabla v \rangle,$$

for all  $v \in H_0^{1,2}(\alpha, \Omega)$ . We will assume that A is uniformly elliptic. In this section we will follow the presentation in [5] to prove

**Theorem 6.1.** Let  $G \in L^p$  for some 1 , and let <math>w be a solution to (6.1). There exists  $\delta > 0$  depending on  $p, \alpha$  such that if  $||A - Id|| \leq \delta$ , then for any compact  $V \subseteq \Omega$ , there exists a constant C depending on  $V, \Omega, n, \alpha, p$  such that

$$\|\nabla w\|_{L^{p}(\alpha,V)} \leq C(\|w\|_{H^{1,2}(\alpha,\Omega)} + \|G\|_{L^{p}(\alpha,\Omega)}).$$

For notational convenience, let us define  $d\mu = \alpha(x_n)dx$ .

**Lemma 6.2.** For any  $\epsilon > 0$  small, there exists a constant C(n) only depending on dimension n, and not on  $\alpha$ , such that for any weak solution w of (6.1) in  $B_4$  satisfying

(6.2) 
$$\frac{1}{\mu(B_4)} \int_{B_4} |\nabla w|^2 d\mu \le 1, \qquad \frac{1}{\mu(B_4)} \int_{B_4} |\mathbf{G}|^2 d\mu \le \epsilon^2, \qquad \|A - I\|_{L^{\infty}} \le \epsilon^2$$

then

(6.3) 
$$\frac{1}{\mu(B_4)} \int_{B_4} |w - v|^2 + |\nabla(w - v)|^2 \ d\mu \le C\epsilon^2,$$

where v is the weak solution to

$$\int_{B_4} \langle \nabla v, \nabla \phi \rangle d\mu = 0 \quad \text{for all } \phi \in H^1_0(\alpha, B_4).$$

*Proof.* Let  $\tilde{w}$  solve

$$\int_{B_4} \langle A \nabla \tilde{w}, \nabla \phi \rangle d\mu = 0,$$

with  $\tilde{w} - w \in H_0^1(\alpha, B_4)$ . Then

$$\frac{1}{\mu(B_4)} \int_{B_4} |\nabla(\tilde{w} - w)^2| d\mu \le 2\frac{1}{\mu(B_4)} \int_{B_4} \langle A\nabla(\tilde{w} - w), \nabla(\tilde{w} - w) \rangle d\mu = -2\frac{1}{\mu(B_4)} \int_{B_4} \mathbf{G} \cdot \nabla(\tilde{w} - w) d\mu \le 2\epsilon \frac{1}{\mu(B_4)} \int_{B_4} |\nabla(\tilde{w} - w)|^2 d\mu$$

Now let v solve

$$\int_{B_4} \langle \nabla v, \nabla \phi \rangle d\mu = 0,$$

with  $v - \tilde{w} \in H_0^1(\alpha, B_4)$ . Then

$$\frac{1}{\mu(B_4)} \int_{B_4} |\nabla(\tilde{w} - v)|^2 \, d\mu = \frac{1}{\mu(B_4)} \int_{B_4} \langle \nabla(\tilde{w} - v), \nabla(\tilde{w} - v) \rangle \, d\mu = \frac{1}{\mu(B_4)} \int_{B_4} \langle \nabla\tilde{w}, \nabla(\tilde{w} - v) \rangle \, d\mu \\ = \frac{1}{\mu(B_4)} \int_{B_4} \langle (I - A) \nabla\tilde{w}, \nabla(\tilde{w} - v) \rangle \, d\mu \le C(n) \epsilon^2 \frac{1}{\mu(B_4)} \int_{B_4} |\nabla(\tilde{w} - v)|^2 \, d\mu.$$

From the triangle inequality we obtain (6.3) for the gradient term. The bound for the function term follows from (4.7) with p = 2.

We now state some essential tools for the Caffarelli-Peral method. The maximal function is defined as

$$(\mathcal{M}g)(x) = \sup_{r>0} \frac{1}{\mu(B_r)} \int_{B_r(x)} |g| \ d\mu$$

Since  $\mu$  is a doubling measure, we have the Vitali covering lemma, from which we obtain maximal function theorems. (see Theorem 2.2 in [16]).

**Theorem 6.3.** For all t > 0 we have the weak 1-1 estimate

$$\mu(\{(\mathcal{M}g) > t\}) \le \frac{C_1}{t} \int_{\mathbb{R}^n} |g| \ d\mu$$

and the  $L^p$  estimate

$$\int_{\mathbb{R}^n} |(\mathcal{M}g)|^p d\mu \le C_p \int_{\mathbb{R}^n} |g|^p \ d\mu$$

where the constants  $C_1$  and  $C_p$  only depend on the doubling constant for  $\mu$ .

**Lemma 6.4.** There is a constant  $N_1$  so that for any  $\eta > 0$  there exists a small epsilon > 0, and if w is a weak solution of (6.1) in  $\Omega \supset B_6$  with

(6.4) 
$$B_1 \cap \{x \in \Omega : \mathcal{M}(|\nabla w|^2(x)) \le 1\} \cap \{x \in \Omega : \mathcal{M}(|\mathbf{G}|^2)(x) \le \epsilon^2\} \ne \emptyset$$
  
and  $||A - I|| \le \epsilon^2$ , then

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^2\} \cap B_1) < \eta \mu(B_1).$$

*Proof.* From (6.4), there exists a point  $x^0 \in B_1$  satisfying

(6.5) 
$$\frac{1}{\mu(B_r)} \int_{B_r(x^0) \cap \Omega} |\nabla w|^2 \, d\mu \le 1, \quad \frac{1}{\mu(B_r)} \int_{B_r(x^0) \cap \Omega} |\mathbf{G}|^2 \, d\mu \le \epsilon^2 \quad \text{for all } r > 0.$$

Since  $B_4(0) \subset B_5(x^0)$ , we have

$$\frac{1}{\mu(B_4)} \int_{B_4} |\mathbf{G}|^2 \ d\mu \le \frac{\mu(B_5(x^0))}{\mu(B_4)} \frac{1}{\mu(B_5(x^0))} \int_{B_5(x_0)} |\mathbf{G}|^2 \ d\mu \le C(n,\alpha)\epsilon^2.$$

Similarly,

$$\frac{1}{\mu(B_4)} \int_{B_4} |\nabla w|^2 \ d\mu \le C(n, \alpha).$$

Dividing u and G by  $C(n, \alpha)$ , we can apply Lemma 6.2 to the  $\alpha$ -harmonic replacement v. From Lemma 4.9 we have

$$\|\nabla v\|_{L^{\infty}(B_3)}^2 \le N_0^2$$

with  $N_0$  depending on n and  $\alpha$ . We now claim

(6.6) 
$$\{x: \mathcal{M}(|\nabla w|^2) > N_1^2\} \cap B_1 \subset \{x: \mathcal{M}(|\nabla w|^2 \chi_{B_4}) > N_0^2\} \cap B_1,$$

where  $N_1^2 = \max 4N_0^2$ , C with C only depending on  $n, \alpha$ . To show this, suppose that

$$x^1 \in \{x : \mathcal{M}(|\nabla w|^2 \chi_{B_4}) > N_0^2\} \cap B_1.$$

For  $r \leq 2$ ,  $B_r(x^1) \subset B_3$ , so that

$$\frac{1}{\mu(B_r(x^1))} \int_{B_r(x^1)} |\nabla w|^2 \ d\mu \le \frac{2}{\mu(B_r(x^1))} \int_{B_r(x^1)} |\nabla (w - v)|^2 + |\nabla v|^2 \ d\mu \le 4N_0^2.$$

Now for r > 2,  $x^0 \in B_r(x^1) \subset B_{2r}(x^0)$  so by (6.5)

$$\frac{1}{\mu(B_r(x^1))} \int_{B_r \cap \Omega} |\nabla w|^2 \ d\mu \le \frac{\mu(B_{2r}(x^0))}{\mu(B_r(x^1))} \frac{1}{\mu(B_{2r}(x^0))} \int_{B_{2r}(x^0) \cap \Omega} |\nabla w|^2 \ d\mu \le C.$$

then the above two inequalities show that

$$x^1 \in \{x : \mathcal{M}(|\nabla w|^2) \le N_1^2\} \cap B_1,$$

and our claim in (6.6) is shown. By (6.6), the weak 1-1 estimate in Theorem 6.3, and inequality (6.3) we have

$$\mu(\{x: \mathcal{M}(|\nabla w|^2)(x) > N_1^2\} \cap B_1) \le \mu(\{x: \mathcal{M}(|\nabla (w-v|^2 \chi_{B_4})(x)(x) > N_0^2\} \cap B_1) \\ \le \frac{C}{N_0^2} \int_{B_4} |\nabla (w-v)|^2 \ d\mu \ \le \frac{C}{N_0^2} \epsilon^2 = \eta \mu(B_1).$$

We now wish to apply Lemma 6.4 to balls of various radii and translations off the thin space  $\{x_n = 0\}$ .

**Lemma 6.5.** Assume that w is a weak solution of (6.1) in  $\Omega \supset B_{6r}(x^0)$ . If

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^2\} \cap B_r(x^0)) \ge \eta \mu(B_r(x^0)),$$

then

$$B_r(x^0) \subset \{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > 1\} \cup \{x \in \Omega : \mathcal{M}(|\boldsymbol{G}|^2)(x) > \epsilon^2\}.$$

*Proof.* If  $x^0$  is centered on the thin space, then the Lemma follows directly from rescaling. Suppose now that  $B_r(x^0) \subset B_{2r}(x^1)$  with  $x^1$  centered on the thin space. Suppose now

$$x^2 \in B_{B_r(x^0)} \cap \{x \in \Omega : \mathcal{M}(|\nabla w|^2(x)) \le 1\} \cap \{x \in \Omega : \mathcal{M}(|\boldsymbol{G}|^2)(x) \le \epsilon^2\}.$$

Since the Lemma holds for  $B_{2r}(x^1)$ , we have

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^2\} \cap B_r(x^0))$$
  
 
$$\subset \mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^2\} \cap B_{2r}(x^1)) \le \eta\mu(B_2) \le C\eta\mu(B_1(x^0))$$

If  $B_{2r}(x^0) \cap \{x_n = 0\} = \emptyset$ , then the proof of Lemma 6.4 goes through exactly as before for the rescaled function since the approximating  $\alpha$ -harmonic function has interior  $C^{1,1}$  estimates from interior regularity. Then rescaling backwards we obtain the Lemma for all balls.

Using induction we obtain the following corollary.

**Corollary 6.6.** Suppose that w is a weak solution to (6.1) in a domain  $\Omega \supset B_6$ . Assume that

$$|\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^2\}| < \eta \mu(B_1).$$

Let 
$$k \in \mathbb{N}$$
 and set  $\eta_1 = C_1 \eta$ . Then  

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^{2k}) \\ \leq \sum_{i=1}^k \eta_1^i \mu(\{x \in B_1 : \mathcal{M}(|\mathbf{G}|^2) > \epsilon^2 N_1^{2(k-i)}\}) + \eta_1^k \mu(\{x \in B_1 : \mathcal{M}(|\nabla w|^2) > 1\})$$

*Proof.* We proceed by induction. For the base case k = 1 we consider

$$S_1 := \{ x \in B_1 : \mathcal{M}(|\nabla w|^2)(x) > N_1^2 \},$$
  

$$S_2 := \{ x \in B_1 : \mathcal{M}(|\mathbf{G}|^2)(x) > \epsilon^2 \} \cup \{ x \in B_1 : \mathcal{M}(|\nabla w|^2) > 1 \}$$

From assumption we have  $\mu(S_1) < \eta \mu(B_1)$ . From Lemma 6.5 we have that whenever  $\mu(S_1 \cap B_r(x)) \ge \eta \mu(B_r(x))$  with 0 < r < 1, then

$$B_r(x) \cap B_1 \subset S_2.$$

From the Vitali covering lemma (see [16]) we have

$$|S_1| \le C_1 \eta |S_2|,$$

with the constant only depending on the doubling constant of  $\alpha$ . We now assume that the conclusion is valid for some k > 1. Let  $w_1 = w/N_1$  and  $G_1 = G/N_1$ . Then with  $w_1$  as a weak solution we obtain

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w_1)|^2)(x) > N_1^2\}) < \eta \mu(B_1).$$

Then by the induction assumption, we have

$$\mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^{2(k+1)}\}) = \mu(\{x \in \Omega : \mathcal{M}(|\nabla w|^2)(x) > N_1^{2k}\})$$

$$\leq \sum_{i=1}^k \eta_1^i \mu(\{x \in B_1 : \mathcal{M}(|\mathbf{G}|^2)(x) > \epsilon^2 N_1^{2(k-i)}\}) + \eta_1^k \mu(\{x \in B_1 : \mathcal{M}(|\nabla w_1|^2)(x) > 1\})$$

$$\leq \sum_{i=1}^{k+1} \eta_1^i \mu(\{x \in B_1 : \mathcal{M}(|\mathbf{G}|^2)(x) > \epsilon^2 N_1^{2(k+1-i)}\}) + \eta_1^{k+1} \mu(\{x \in B_1 : \mathcal{M}(|\nabla w|^2)(x) > 1\}),$$

and the conclusion is proven.

We now prove Theorem 6.1.

*Proof.* By multiplying our solution w by a small constant, we can assume  $\|G\|_{L^p(B_6)}$  is small and that

$$\mu(x \in B_6 : \mathcal{M}(|\nabla w|^2)(x) > N_1^2 \cap B_1) < \eta \mu(B_1).$$

Since  $\mathbf{G} \in L^p(B_6)$  we have from the strong *p*-*p* estimates in Theorem 6.3 that  $\mathcal{M}(|\mathbf{G}|^2) \in L^{p/2}(B_6)$ . We now use a standard tool from measure theory (see Lemma 7.3 in [7]) to see there is a constant C depending on  $\alpha, \epsilon, p, N_1$  such that

$$\sum_{k=0}^{\infty} N_1^{pk} \mu(\{x \in B_1 : \mathcal{M}(|\boldsymbol{G}|^2)(x) > \epsilon^2 N_1^{2k}\}) \le C \|\mathcal{M}(|\boldsymbol{G}|^2)\|_{L^{p/2}(B_6)}^{p/2}.$$

Using the strong p-p estimates on the right hand side above as well as the smallness condition on  $\|g\|_{L^p}$  we have

$$\sum_{k=0}^{\infty} N_1^{pk} \mu(\{x \in B_1 : \mathcal{M}(|\boldsymbol{G}|^2)(x) > \epsilon^2 N_1^{2k}\}) \le 1.$$

We now utilize p > 2 in the following computation.

$$\begin{split} &\sum_{k=0}^{\infty} N_{1}^{pk} \mu(x \in B_{1} : \mathcal{M}(|\nabla w|^{2})(x) > N_{1}^{2k}) \\ &\leq \sum_{k=1}^{\infty} N_{1}^{pk} \bigg( \sum_{i=1}^{k} \eta_{1}^{i} \mu(\{x \in B_{1} : \mathcal{M}(|\mathbf{G}|^{2})(x) > \epsilon^{2} N_{1}^{2(k-i)}\}) \\ &+ \eta_{1}^{k} \mu(\{x \in B_{1} : \mathcal{M}(|\nabla w|^{2})(x) > 1\}) \bigg) \\ &= \sum_{i=1}^{\infty} (N_{1}^{p} \eta_{1})^{i} \bigg( \sum_{k=i}^{\infty} N_{1}^{p(k-i)} \mu(\{x \in B_{1} : \mathcal{M}(|\mathbf{G}|^{2})(x) > \epsilon^{2} N_{1}^{2(k-i)}\}) \bigg) \\ &+ \sum_{k=1}^{\infty} (N_{1}^{p} \eta_{1})^{k} \mu(\{x \in B_{1} : \mathcal{M}(|\nabla w|^{2})(x) > 1\}) \\ &\leq C \sum_{k=1}^{\infty} (N_{1}^{p} \eta_{1}^{k}) \\ &\leq +\infty. \end{split}$$

Thus using Lemma 7.3 in [7] we have that  $\mathcal{M}(|\nabla w|^2) \in L^{p/2}(B_6)$ . Therefore,  $\nabla w \in L^p(B_6)$ .

#### Acknowledgements

This project was initiated while the authors stayed at Institute Mittag Leffler (Sweden), during the program Geometric aspects of nonlinear PDE. The authors wish to thank Georg Weiss for helpful conversations on this topic. H. Shahgholian was supported by Swedish research Council. D. Kriventsov was supported by NSF DMS grant 2247096.

#### DECLARATIONS

Data availability statement: All data needed are contained in the manuscript.

Funding and/or Conflicts of interests/Competing interests: The authors declare that there are no financial, competing or conflict of interests.

#### References

- Mark Allen, Dennis Kriventsov, and Henrik Shahgholian. The inhomogeneous boundary Harnack principle for fully nonlinear and p-Laplace equations. Ann. Inst. H. Poincaré C Anal. Non Linéaire, 40(1):133–156, 2023.
- Mark Allen and Henrik Shahgholian. A new boundary Harnack principle (equations with right hand side). Arch. Ration. Mech. Anal., 234(3):1413–1444, 2019.
- [3] Hans W. Alt and Daniel Phillips. A free boundary problem for semilinear elliptic equations. J. Reine Angew. Math., 368:63–107, 1986.
- [4] Ioannis Athanasopoulos and Luis A. Caffarelli. A theorem of real analysis and its application to free boundary problems. Comm. Pure Appl. Math., 38(5):499–502, 1985.
- [5] Sun-Sig Byun. Elliptic equations with BMO coefficients in Lipschitz domains. Trans. Amer. Math. Soc., 357(3):1025–1046, 2005.
- [6] Luis A. Caffarelli. The regularity of free boundaries in higher dimensions. Acta Math., 139(3-4):155–184, 1977.
- [7] Luis A. Caffarelli and Xavier Cabré. Fully nonlinear elliptic equations, volume 43 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1995.
- [8] Luis A. Caffarelli and I. Peral. On W<sup>1,p</sup> estimates for elliptic equations in divergence form. Comm. Pure Appl. Math., 51(1):1–21, 1998.
- [9] Olivaine S. de Queiroz and Henrik Shahgholian. A free boundary problem with log-term singularity. Interfaces Free Bound., 19(3):351–369, 2017.
- [10] D. De Silva and O. Savin. The Alt-Phillips functional for negative powers. Bull. Lond. Math. Soc., 55(6):2749– 2777, 2023.
- [11] Daniela De Silva and Ovidiu Savin.  $C^{\infty}$  regularity of certain thin free boundaries. Indiana Univ. Math. J., 64(5):1575–1608, 2015.
- [12] Daniela De Silva and Ovidiu Savin. A note on higher regularity boundary Harnack inequality. Discrete Contin. Dyn. Syst., 35(12):6155–6163, 2015.
- [13] Alessio Figalli. Free boundary regularity in obstacle problems, 2018.
- [14] Mariano Giaquinta and Enrico Giusti. Differentiability of minima of nondifferentiable functionals. Invent. Math., 72(2):285–298, 1983.
- [15] Enrico Giusti. Direct methods in the calculus of variations. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [16] Juha Heinonen. Lectures on analysis on metric spaces. Universitext. Springer-Verlag, New York, 2001.
- [17] Yash Jhaveri and Robin Neumayer. Higher regularity of the free boundary in the obstacle problem for the fractional Laplacian. Adv. Math., 311:748–795, 2017.
- [18] David Kinderlehrer and Louis Nirenberg. Regularity in free boundary problems. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 4(2):373–391, 1977.
- [19] Dennis Kriventsov and Henrik Shahgholian. Optimal regularity for a two-phase obstacle-like problem with logarithmic singularity. Comm. Partial Differential Equations, 46(10):1831–1850, 2021.
- [20] Arshak Petrosyan, Henrik Shahgholian, and Nina Uraltseva. Regularity of free boundaries in obstacle-type problems, volume 136 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2012.
- [21] Xavier Ros-Oton and Clara Torres-Latorre. New boundary Harnack inequalities with right hand side. J. Differential Equations, 288:204–249, 2021.
- [22] Henrik Shahgholian and Georg S. Weiss. Aleksandrov and Kelvin reflection and the regularity of free boundaries. In Free boundary problems, volume 154 of Internat. Ser. Numer. Math., pages 391–401. Birkhäuser, Basel, 2007.
- [23] Yannick Sire, Susanna Terracini, and Stefano Vita. Liouville type theorems and regularity of solutions to degenerate or singular problems part I: even solutions. *Comm. Partial Differential Equations*, 46(2):310–361, 2021.
- [24] Clara Torres-Latorre. Parabolic boundary harnack inequalities with right-hand side, 2023.
- [25] Chilin Zhang. On higher order boundary harnack and analyticity of free boundaries, 2023.

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY, PROVO, UT 84602, US *Email address*: allen@mathematics.byu.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ *Email address*: dnk34@math.rutgers.edu

DEPARTMENT OF MATHEMATICS, KTH ROYAL INSTITUTE OF TECHNOLOGY, STOCKHOLM, SWEDEN *Email address*: henriksh@kth.se