

the fine structure of the spectra of Langmuir waves in plasma and describes, under suitable boundary conditions, the propagation of the spectral packet of Langmuir oscillations against the background of thermal noise (see [10], [11]).

Thus, we observe an intriguing fact: some of the Toda-type lattices in 3D admit a hierarchy of reductions that have increased integrability such as exact solvability. Initiated with this observation we suggested a classification algorithm of the integrable 3D lattices based on the existence of a hierarchy of the reductions satisfying the more weak requirement than exact solvability. We assume that the reduced systems have complete sets of the characteristic integrals corresponding to each characteristic direction (see [12], [13]). Recall that systems of the hyperbolic type equations admitting complete sets of integrals for each direction are called integrable in the sense of Darboux (for the details see, for instance, monograph [14]). An effective criterion for Darboux integrability of a PDE system of hyperbolic type, based on the concept of characteristic Lie algebras, was proposed by A.B. Shabat in 1980 in his report at the conference dedicated to the memory of I.G. Petrovsky. An important statement is proved that a system of the hyperbolic type equations is integrable in the sense of Darboux if and only if its characteristic Lie algebras corresponding to both characteristic directions are of finite dimension (see [15], [14]).

In the article [12] a problem of the complete description of the integrable equations of the form

$$u_{n,xy} = f(u_{n+1}, u_n, u_{n-1}, u_{n,x}, u_{n,y}) \quad (1.3)$$

by using the method of the characteristic Lie algebras is considered. Later in [13] the problem is formulated and discussed to describe all equations of the form (1.3) for which there exists a pair of functions $f_1 = f_1(u_1, u_2, u_{1,x}, u_{1,y})$ and $f_2 = f_2(u_m, u_{m-1}, u_{m,x}, u_{m,y})$ such that for any integer $m \geq 2$ the reduced system of the hyperbolic type equations

$$\begin{aligned} u_{1,xy} &= f_1(u_1, u_2, u_{1,x}, u_{1,y}), \\ u_{j,xy} &= f(u_{j+1}, u_j, u_{j-1}, u_{j,x}, u_{j,y}), \quad 1 < j < m, \\ u_{m,xy} &= f_2(u_m, u_{m-1}, u_{m,x}, u_{m,y}) \end{aligned} \quad (1.4)$$

is integrable in the sense of Darboux.

For lattices having the following particular quasilinear form

$$u_{n,xy} = A_1 u_{n,x} u_{n,y} + A_2 u_{n,x} + A_3 u_{n,y} + A_4 \quad (1.5)$$

where the coefficients are smooth functions of the dynamical variables $A_i = A_i(u_{n+1}, u_n, u_{n-1})$ for $i = 1, 2, 3, 4$, the classification problem mentioned above was completely solved in the works [13], [16], [17]. This requirement was motivated by the fact that all examples of integrable chains in 3D known in the literature depend linearly on each of the first derivatives. Here we give the complete list of the lattices in the class (1.5) which pass the test formulated above:

- (E1) $u_{n,xy} = e^{u_{n+1}-2u_n+u_{n-1}},$
- (E2) $u_{n,xy} = e^{u_{n+1}} - 2e^{u_n} + e^{u_{n-1}},$
- (E3) $u_{n,xy} = e^{u_{n+1}-u_n} - e^{u_n-u_{n-1}},$
- (E4) $u_{n,xy} = (u_{n+1} - 2u_n + u_{n-1}) u_{n,x},$
- (E5) $u_{n,xy} = (e^{u_{n+1}-u_n} - e^{u_n-u_{n-1}}) u_{n,x},$
- (E6) $u_{n,xy} = \alpha_n u_{n,x} u_{n,y}, \quad \alpha_n = \frac{1}{u_n - u_{n-1}} - \frac{1}{u_{n+1} - u_n}$
- (E7) $u_{n,xy} = \alpha_n (u_{n,x} + u_n^2 - 1)(u_{n,y} + u_n^2 - 1) - 2u_n(u_{n,x} + u_{n,y} + u_n^2 - 1).$

The classification problem in the general case of (1.3) remains open. Alternative approaches to the problem of classifying integrable lattices in 3D are discussed, for example, in the works [18], [19].

Equations (E1)-(E6) are well-known integrable models (see, for instance, [4]). Equation (E7) was obtained during the classification process in [13]. The Lax pair for it was found in [20]. For the lattices (E6), (E7) the reductions of the form (1.4) are investigated in [21] and correspondingly in [13] admitting finite-dimensional characteristic Lie algebras for arbitrary m . However in these articles the integrals are not presented. Reductions of the form (1.4) for the lattice (E1) coincide with the exponential type systems related to the Cartan matrices of the simple Lie algebras of the series A. As it was shown in [15] they are certainly Darboux integrable for arbitrary m . In the case of the lattice (E3) the Darboux integrable reductions are studied in [22], where a generating function for the characteristic integrals for the reduced systems are obtained.

Characteristic Lie algebras provide an effective (see [14], [23]), but rather labor-intensive method for constructing integrals, so it is not convenient for large values of m . Various alternative approaches to the problem of constructing characteristic integrals, including Laplace's cascade method in scalar and matrix versions, were previously used in the works [22], [24], [25], [26], [27], [28]. The purpose of the present article is to develop a method for constructing characteristic integrals based on the use of the Lax pair and the Miura type transformations and then apply it to some key equations of the list above.

It is well known that in the case of the soliton systems, in order to find the densities of the integrals of motion, one can use the formal asymptotic expansion of the Lax operator eigenfunction in powers of the spectral parameter. This technique is not suitable for searching for characteristic integrals, since Lax pairs for Darboux integrable systems do not contain any spectral parameter.

Let us briefly discuss on the content of the article. In the second section, the definition of the complete set of independent characteristic integrals of a system of hyperbolic type differential equations is given, and a convenient criterion for the completeness is formulated. Using the Liouville equation as an example, it is shown that the presence of a complete set of integrals ensures separation of the variables, which, in turn, reduces the problem of integrating PDEs to the problem of finding explicit solutions to ODEs (see Remark below). In Section 3, we briefly discuss the work of S. V. Smirnov [22], where generating functions of characteristic integrals for several finite-field versions of the Toda lattice (E3) are presented. The approach of [22] is closely related to the sequence of the Laplace transformations of a linear equation of hyperbolic type with variable coefficients. We modified this approach and adapted it to the case of equations that are not directly related to the Laplace cascade method. In sections 4-6 we apply this algorithm for finding integrals to specific systems that are reductions of the lattices (E6), (E7) and, correspondingly, the coupled system (1.2). In all three cases generating functions are given, allowing one to construct complete sets of characteristic integrals for reductions of the form (1.4) of arbitrary order m .

2 Complete sets of characteristic integrals

Let us consider a system of the hyperbolic type equations of the form

$$u_{i,xy} = F_i(u_1, \dots, u_N, u_{1,x}, \dots, u_{N,x}, u_{1,y}, \dots, u_{N,y}) \quad i = 1, 2, \dots, N, \quad (2.1)$$

where F_i are analytic functions defined in some region of the space \mathbf{C}^{3N} .

We recall some basic definitions (see, for example, [14]). Smooth function of the form

$$I = I(u_1, \dots, u_N, u_{1,x}, \dots, u_{N,x}, u_{1,xx}, \dots, u_{N,xx}, \dots, u_{1,r}, \dots, u_{N,r}),$$

where the notation is accepted $u_{j,s} := \frac{\partial^s}{\partial x^s} u_j$, is called a y -integral of the order r of the system (2.1), if the condition is satisfied $D_y I = 0$, where the derivative with respect to the variable y is evaluated by means of the system (2.1). In other words to look for the y -integral we have a relation

$$D_y I = \sum_i (u_{i,y} \frac{\partial}{\partial u_i} + F_i \frac{\partial}{\partial u_{i,x}} + D_x(F_i) \frac{\partial}{\partial u_{i,xx}} + \dots) I = Y I = 0.$$

A set of y -integrals $\{I_i\}_{i=1}^{i=N}$ having the orders r_i for the system of equations (2.1) constitutes a complete set of independent y -integrals, if the condition holds

$$\det \left(\frac{\partial I_j}{\partial u_{i,m}} \right) = \begin{vmatrix} \frac{\partial I_1}{\partial u_{1,r_1}} & \frac{\partial I_1}{\partial u_{2,r_1}} & \dots & \frac{\partial I_1}{\partial u_{N,r_1}} \\ \dots & \dots & \dots & \dots \\ \frac{\partial I_N}{\partial u_{1,r_N}} & \frac{\partial I_N}{\partial u_{2,r_N}} & \dots & \frac{\partial I_N}{\partial u_{N,r_N}} \end{vmatrix} \neq 0. \quad (2.2)$$

Let's formulate an effective algebraic criterion (see [14]) for the existence of a complete set of integrals of the system (2.1).

Theorem 1. *System of equations (2.1) admits a complete set of independent y -integrals if and only if the characteristic Lie algebra over the ring of locally analytic functions of the variables $\bar{u}_y, \bar{u}, \bar{u}_x, \bar{u}_{xx}, \dots$, where $\bar{u} = (u_1, u_2, \dots, u_N)$, generated by the characteristic operators*

$$X_j = \frac{\partial}{\partial u_{j,y}}, \quad j = 1, \dots, N; \quad Y = \sum_i u_{i,y} \frac{\partial}{\partial u_i} + F_i \frac{\partial}{\partial u_{i,x}} + D_x(F_i) \frac{\partial}{\partial u_{i,xx}} + \dots$$

is of finite dimension.

Integrals and characteristic algebra in the direction x are defined similarly. As an illustrative example, we consider the well-known Liouville equation

$$u_{xy} = e^u. \quad (2.3)$$

Differentiating it with respect to x and using the equation (2.3) once again, we find $u_{xxy} = u_x u_{xy}$. Next, we integrate it over y and arrive at the relation

$$u_{xx} - \frac{1}{2} u_x^2 = h(x), \quad (2.4)$$

where $h(x)$ is an arbitrary function. Similarly we have

$$u_{yy} - \frac{1}{2} u_y^2 = g(y). \quad (2.5)$$

Now it's clear that the functions $I = u_{xx} - \frac{1}{2} u_x^2$ and $J = u_{yy} - \frac{1}{2} u_y^2$ are, respectively, y - and x -integrals. Due to the presence of the complete sets of integrals, the variables in the Liouville equation are separated in such a way that the problem of integrating the equation is reduced to finding a joint solution of two ordinary differential equations. From the relations (2.4), (2.5) it is easy to derive the formula for the general solution of the equation (2.3), found by Liouville:

$$u(x, y) = \ln \frac{2p'(x)q'(y)}{(p(x) + q(y))^2},$$

$$\begin{aligned}
u_{2,xy} &= u_{2,x}u_{2,y} \left(\frac{1}{u_2 - u_1} - \frac{1}{u_3 - u_2} \right), \\
&\dots\dots\dots \\
u_{m,xy} &= u_{m,x}u_{m,y} \left(\frac{1}{u_m - u_{m-1}} - \frac{1}{c_{m+1} - u_m} \right).
\end{aligned} \tag{4.2}$$

It was proved in [16] that for an arbitrary $m \geq 1$ the characteristic Lie algebras corresponding to the directions x and y have finite dimension that guarantees the existence of complete sets of characteristic integrals in both directions. However the integrals themselves were not found. Below we present the generating function for these integrals in terms of the Lax pair.

To begin with, we note that the first-order integral can be easily derived from elementary considerations. Indeed, let us rewrite the system in the form:

$$\begin{aligned}
\frac{u_{1,xy}}{u_{1,x}} &= u_{1,y} \left(\frac{1}{u_1 - c_0} - \frac{1}{u_2 - u_1} \right), \\
\frac{u_{2,xy}}{u_{2,x}} &= u_{2,y} \left(\frac{1}{u_2 - u_1} - \frac{1}{u_3 - u_2} \right), \\
&\dots\dots\dots \\
\frac{u_{m,xy}}{u_{m,x}} &= u_{m,y} \left(\frac{1}{u_m - u_{m-1}} - \frac{1}{c_{m+1} - u_m} \right)
\end{aligned} \tag{4.3}$$

and then adding all the equations we arrive at the relation

$$D_y \log(u_{1x}u_{2x} \dots u_{mx}) = D_y \log(g_1g_2 \dots g_{m+1}).$$

It obviously follows from this fact that the function

$$J = u_{1x}u_{2x} \dots u_{mx}(g_1g_2 \dots g_{m+1}) \tag{4.4}$$

is a y -integral. Here $g_1 = \frac{1}{u_1 - c_0}$, $g_2 = \frac{1}{u_2 - u_1}$, \dots $g_m = \frac{1}{u_m - u_{m-1}}$, $g_{m+1} = \frac{1}{c_{m+1} - u_m}$. To search for higher order integrals, we will use the Lax pair, which was found by E.V. Ferapontov¹:

$$\psi_{n,x} = a_n(\psi_{n+1} - \psi_n), \quad \psi_{n,y} = b_n(\psi_n - \psi_{n-1}), \tag{4.5}$$

where $a_n = \frac{u_{n,x}}{u_{n+1} - u_n}$, $b_n = \frac{u_{n,y}}{u_n - u_{n-1}}$. By imposing on the system (4.5), in addition to the equalities (4.1), also conditions of the form $\psi_0 = 0$, $\psi_{m+1} = 0$ we can obtain a Lax pair for system (4.2) which is given by the following two systems of linear equations, the first one is:

$$\begin{aligned}
\psi_{1,x} &= a_1(\psi_2 - \psi_1), \\
\psi_{2,x} &= a_2(\psi_3 - \psi_2), \\
&\dots\dots\dots \\
\psi_{m-1,x} &= a_{m-1}(\psi_m - \psi_{m-1}), \\
\psi_{m,x} &= -a_m\psi_m.
\end{aligned} \tag{4.6}$$

Likewise, the second system

$$\begin{aligned}
\psi_{1,y} &= b_1\psi_1, \\
\psi_{2,y} &= b_2(\psi_2 - \psi_1),
\end{aligned}$$

¹Private communication, 2020.

$$\begin{aligned} & \dots\dots\dots (4.7) \\ \psi_{m-1,y} &= b_{m-1}(\psi_{m-1} - \psi_{m-2}), \\ \psi_{m,y} &= b_m(\psi_m - \psi_{m-1}). \end{aligned}$$

From the system (4.6) one can easily obtain a differential relation connecting the functions ψ_1 and ψ_m :

$$\psi_m = \left(\frac{1}{a_{m-1}} D_x + 1 \right) \left(\frac{1}{a_{m-2}} D_x + 1 \right) \dots \left(\frac{1}{a_1} D_x + 1 \right) \psi_1. \quad (4.8)$$

Now it is easy to see that from the last equation of the system (4.6) and equation (4.8) the relation follows

$$R\psi_1 = 0, \quad (4.9)$$

where $R = (D_x + a_m) \left(\frac{1}{a_{m-1}} D_x + 1 \right) \left(\frac{1}{a_{m-2}} D_x + 1 \right) \dots \left(\frac{1}{a_1} D_x + 1 \right) \psi_1$. Therefore, the eigenfunction ψ_1 is a joint solution to the overdetermined system consisting of (4.9) and the equation

$$(D_y - b_1) \psi_1 = 0. \quad (4.10)$$

By virtue of the cutoff condition (4.1) the system (4.9)-(4.10) can be essentially simplified. Let us use an explicit expression for the potential b_1 and integrate the equation (4.10), rewritten as follows

$$\frac{d\psi_1}{\psi_1} = \frac{u_{1,y} dy}{u_1 - C_0}.$$

As a result, we obtain an expression for the eigenfunction $\psi_1 = (u_1 - C_0) \varphi_1(x)$. Afterwards in (4.9)-(4.10) we move on to the new sought function $\varphi_1(x)$ and arrive at a system of the form

$$\hat{R}\varphi_1 = 0, \quad \frac{\partial}{\partial y}\varphi_1 = 0, \quad (4.11)$$

where

$$\hat{R} = (D_x + a_m) \left(\frac{1}{a_{m-1}} D_x + 1 \right) \dots \left(\frac{1}{a_1} D_x + 1 \right) (u_1 - C_0).$$

Repeatedly applying the ratio

$$(D_x + f) \frac{1}{g} = \frac{1}{g} (D_x + f - (\ln g)_x), \quad (4.12)$$

we rewrite the operator \hat{R} as follows

$$\hat{R} = \frac{u_1 - C_0}{a_1 a_2 \dots a_m} B,$$

where the operator B has the form

$$B = (D_x + r_m) (D_x + r_{m-1}) \dots (D_x + r_1). \quad (4.13)$$

Here $r_1 = a_1 + \frac{u_{1,x}}{u_1 - C_0}$, $r_2 = a_2 - (\ln a_1)_x + \frac{u_{1,x}}{u_1 - C_0}$, but in the general case

$$r_k = a_k - \sum_{i=1}^{k-1} (\ln a_i)_x + \frac{u_{1,x}}{u_1 - C_0}, \quad k \leq m. \quad (4.14)$$

After all transformations, the system (4.11) converts into

$$B\varphi_1 = 0, \quad \frac{\partial}{\partial y}\varphi_1 = 0. \quad (4.15)$$

Let us represent the differential operator B in the following expanded form

$$B = D_x^m + \beta_1 D_x^{m-1} + \dots + \beta_m. \quad (4.16)$$

It is easy to verify, by analogy with the previous section, that the coefficients β_j of this expansion are characteristic integrals of the system of nonlinear equations (4.2).

The following example is really instructive.

Example 1. Let us assume that $m = 2$ in (4.2), then we arrive at the system:

$$\begin{aligned} \frac{u_{1,xy}}{u_{1,x}} &= u_{1,y} \left(\frac{1}{u_1 - C_0} - \frac{1}{u_2 - u_1} \right), \\ \frac{u_{2,xy}}{u_{2,x}} &= u_{2,y} \left(\frac{1}{u_2 - u_1} - \frac{1}{C_3 - u_2} \right). \end{aligned} \quad (4.17)$$

Calculations show that the operator B in this case is reduced to the following form:

$$B = D_x^2 + \left(-\frac{u_{1,xx}}{u_{1,x}} + 2g_1 u_{1,x} + (g_2 + g_3) u_{2,x} \right) D_x + (C_3 - C_0) u_{1,x} u_{2,x} g_1 g_2 g_3, \quad (4.18)$$

where

$$g_1 = \frac{1}{u_1 - C_0}, \quad g_2 = \frac{1}{u_2 - u_2}, \quad g_3 = \frac{1}{C_3 - u_2}. \quad (4.19)$$

Therefore, the required integrals are

$$\beta_1 = -\frac{u_{1,xx}}{u_{1,x}} + 2g_1 u_{1,x} + (g_2 + g_3) u_{2,x}, \quad \beta_2 = (C_3 - C_0) u_{1,x} u_{2,x} g_1 g_2 g_3.$$

It is worth noting that for $C_3 = C_0$ the integral β_2 becomes trivial, i.e. $\beta_2 = 0$. In other words formula (4.16) does not always define a complete set of independent integrals, since the first-order integral J given by the formula (4.4) might be lost. However, even in this case, the system of equations (4.17) admits a complete set of y -integrals

$$I_1 = -\frac{u_{1,xx}}{u_{1,x}} + 2g_1 u_{1,x} + (g_2 + g_3) u_{2,x}, \quad I_2 = u_{1,x} u_{2,x} g_1 g_2 g_3.$$

Let us examine the structure of the set of integrals obtained from the expansion (4.16) in order to construct a complete set of integrals of the system (4.2). As it can be seen from the formula (2.2), expressing the criterion for the independence of integrals, it is necessary to clarify the form of the dependence of the integrals on the highest derivatives. From explicit formulas (4.14) we can derive some simplified representations for the functions r_2, r_3, \dots, r_m , focusing on the dependence of these functions on the highest derivatives

$$r_k = -\sum_{i=1}^{k-1} \frac{u_{i,xx}}{u_{i,x}} + h[u_x], \quad 2 \leq k \leq m.$$

Here the symbol $h[u_x]$ indicates that the remaining terms can only depend on the dynamical variables u_1, u_2, \dots, u_m and their first-order derivatives. Our goal is to obtain similar representations for integrals β_j defined in terms of the operator B (see (4.16)). Let us introduce the notation

$$H_k := (D_x + r_k)(D_x + r_{k-1}) \dots (D_x + r_1). \quad (4.20)$$

For the operator H_2 the representation obviously holds

$$H_2 = D_x^2 - \left(\frac{u_{1,xx}}{u_{1,x}} + h[u_x] \right) D_x + h[u_{xx}].$$

5 On reductions of the lattice (E7)

Lattice (E7) was found in [13] as a result of the integrable classification of nonlinear lattices. In [20] a Lax pair was found for this equation

$$\begin{aligned}\psi_{n,x} &= \frac{v_{n,x} + v_n^2 - 1}{v_{n+1} - v_n} (\psi_{n+1} - \psi_n) - v_n \psi_n, \\ \psi_{n,y} &= \frac{v_{n,y} + v_n^2 - 1}{v_n - v_{n-1}} (\psi_n - \psi_{n-1}) - v_n \psi_n.\end{aligned}\tag{5.1}$$

In other words system (5.1) is consistent if and only if the function $v_n = v_n(x, y)$ solves equation (E7).

Let's consider a finite-field system of equations of hyperbolic type

$$\begin{aligned}v_{1,xy} &= \alpha_1 (v_{1,x} + v_1^2 - 1) (v_{1,y} + v_1^2 - 1) - 2v_1 (v_{1,x} + v_{1,y} + v_1^2 - 1), \\ v_{k,xy} &= \alpha_k (v_{k,x} + v_k^2 - 1) (v_{k,y} + v_k^2 - 1) - 2v_k (v_{k,x} + v_{k,y} + v_k^2 - 1), \\ v_{m,xy} &= \alpha_m (v_{m,x} + v_m^2 - 1) (v_{m,y} + v_m^2 - 1) - 2v_m (v_{m,x} + v_{m,y} + v_m^2 - 1)\end{aligned}\tag{5.2}$$

obtained from the lattice (E7) by imposing the following termination conditions:

$$v_0 = \epsilon, \quad v_{m+1} = \delta.\tag{5.3}$$

Here $2 \leq k \leq m - 1$ and

$$\begin{aligned}\alpha_1 &= \frac{1}{v_1 - \epsilon} - \frac{1}{v_2 - v_1}, \\ \alpha_k &= \frac{1}{v_k - v_{k-1}} - \frac{1}{v_{k+1} - v_k}, \\ \alpha_m &= \frac{1}{v_m - v_{m-1}} - \frac{1}{\delta - v_m}.\end{aligned}$$

Note that each of the parameters ϵ and δ can take only one of two values $\epsilon = \pm 1$, $\delta = \pm 1$. We emphasize that in this case the linear system of equations (5.1) allows us to obtain the Lax pair for the reduced system (5.2). To do this, in addition to the conditions (5.3) for cutting off the chain, it is necessary to impose conditions for terminating the Lax pair:

$$\psi_0 = 0, \quad \psi_{m+1} = 0.\tag{5.4}$$

As a result, we arrive at the linear problem in x :

$$\begin{aligned}\psi_{1,x} &= \frac{v_{1,x} + v_1^2 - 1}{v_2 - v_1} (\psi_2 - \psi_1) - v_1 \psi_1 \\ &\dots\dots\dots \\ \psi_{m-1,x} &= \frac{v_{m-1,x} + v_{m-1}^2 - 1}{v_m - v_{m-1}} (\psi_m - \psi_{m-1}) - v_{m-1} \psi_{m-1}, \\ \psi_{m,x} &= \left(\frac{v_{m,x}}{v_m - \delta} + \delta \right) \psi_m,\end{aligned}\tag{5.5}$$

and the linear problem in y

$$\psi_{1,y} = \left(\frac{v_{1,y}}{v_1 - \epsilon} + \epsilon \right) \psi_1,$$

As a result, the system of equations (5.11) above takes the form:

$$\frac{\partial}{\partial y}\varphi_1 = 0, \quad L\varphi_1 = 0, \quad (5.15)$$

where

$$L = (D_x + r_m)(D_x + r_{m-1}) \dots (D_x + r_1).$$

Let us present the operator L in an expanded form

$$L = D_x^m + \beta_1 D_x^{m-1} + \dots + \beta_m. \quad (5.16)$$

Using arguments similar to those given above in Section 3, we can use the system (5.15) to show that the coefficients $\beta_1, \beta_2, \dots, \beta_m$ are y -integrals of the system of equations (5.2). In order to construct a complete set of independent y -integrals of the system (5.2) we need the following auxiliary statement.

Lemma 1. *The function*

$$J = \frac{(v_{1,x} + v_1^2 - 1)(v_{2,x} + v_2^2 - 1) \dots (v_{m,x} + v_m^2 - 1)}{(v_1 - \epsilon)(v_2 - v_1)(v_3 - v_2) \dots (v_m - v_{m-1})(v_m - \delta)} e^{-(\epsilon + \delta)y} \quad (5.17)$$

is a first order y -integral of the system of equations (5.2).

The lemma is proved by a simple but cumbersome calculation, so we present it in the specific case $m = 3$. Consider the system (5.2) for $m = 3$ and reduce it to the following form:

$$\begin{aligned} D_y \log(v_{1,x} + v_1^2 - 1) &= \frac{v_{1,y}}{v_1 - \epsilon} - v_1 + \epsilon - \frac{v_{1,y} + v_1^2 - 1}{v_2 - v_1} \\ D_y \log(v_{2,x} + v_2^2 - 1) &= \frac{v_{2,y} + v_2^2 - 1}{v_2 - v_1} - 2v_2 - \frac{v_{2,y} + v_2^2 - 1}{v_3 - v_2} \\ D_y \log(v_{3,x} + v_3^2 - 1) &= \frac{v_{3,y} + v_3^2 - 1}{v_3 - v_2} - 2v_3 - \frac{v_{3,y} + v_3^2 - 1}{\delta - v_3}. \end{aligned} \quad (5.18)$$

Let's add up all the equations of the system (5.18) and get the relation

$$\begin{aligned} D_y \log(v_{1,x} + v_1^2 - 1)(v_{2,x} + v_2^2 - 1)(v_{3,x} + v_3^2 - 1) = \\ D_y \log(v_1 - \epsilon)(v_2 - v_1)(v_3 - v_2)(v_3 - \delta) + \epsilon + \delta, \end{aligned}$$

from which it follows that

$$D_y \log \frac{(v_{1,x} + v_1^2 - 1)(v_{2,x} + v_2^2 - 1)(v_{3,x} + v_3^2 - 1)}{(v_1 - \epsilon)(v_2 - v_1)(v_3 - v_2)(v_3 - \delta)} = \epsilon + \delta. \quad (5.19)$$

Now it is very easy to get the statement of the lemma.

Consider the following set of y -integrals of the system of equations (5.2):

$$I_1 := -\beta_{m-1}, I_2 := \beta_{m-2}, \dots, I_{m-1} = -\beta_1, I_m = J, \quad (5.20)$$

where the functions β_j coincide with the expansion coefficients (5.16) of the operator L , and the integral J is given by (5.17).

Theorem 3. *The functions I_1, I_2, \dots, I_m form a complete set of independent y -integrals of the system of equations (5.2).*

To prove the theorem, it is enough to verify that the determinant of the matrix constructed according to the rule

$$\begin{vmatrix} \frac{\partial I_1}{\partial u_{1,[m]}} & \frac{\partial I_1}{\partial u_{2,[m]}} & \cdots & \frac{\partial I_1}{\partial u_{m,[m]}} \\ \frac{\partial I_2}{\partial u_{1,[m-1]}} & \frac{\partial I_2}{\partial u_{2,[m-1]}} & \cdots & \frac{\partial I_2}{\partial u_{m,[m-1]}} \\ \cdots & \cdots & \cdots & \cdots \\ \frac{\partial I_m}{\partial u_{1,x}} & \frac{\partial I_m}{\partial u_{2,x}} & \cdots & \frac{\partial I_m}{\partial u_{m,x}} \end{vmatrix} \quad (5.21)$$

differs from identical zero. To do this, it is necessary to find out the dependence of the integrals on the higher order derivatives of the dynamical variables included in them. From explicit representations (5.14) for functions r_k we have

$$r_k = -\frac{v_{1,xx}}{v_{1,x} + v_1^2 - 1} - \frac{v_{2,xx}}{v_{2,x} + v_2^2 - 1} - \dots - \frac{v_{k-1,xx}}{v_{k-1,x} + v_{k-1}^2 - 1} + h[v_x], \quad (5.22)$$

where $2 \leq k \leq m$. Here the symbol $h[v_x]$ means that remaining terms contain dynamical variables and their first order derivatives. By analogy with the formula (4.21) the operator L can be represented as:

$$\begin{aligned} L = & D_x^m - \left(\frac{(m-1)v_{1,xx}}{v_{1,x} + v_1^2 - 1} + \frac{(m-2)v_{2,xx}}{v_{2,x} + v_2^2 - 1} - \dots - \frac{v_{k-1,xx}}{v_{k-1,x} + v_{k-1}^2 - 1} + h[v_x] \right) D_x^{m-1} - \\ & - \left(\frac{(m-2)v_{1,xxx}}{v_{1,x} + v_1^2 - 1} + \frac{(m-3)v_{2,xxx}}{v_{2,x} + v_2^2 - 1} + \dots + \frac{v_{m-2,xxx}}{v_{m-2,x}} + h[v_{xx}] \right) D_x^{k-2} - \dots - \\ & - \left(\frac{v_{1,[m]}}{v_{1,x} + v_1^2 - 1} + h[v_{[k-1]}] \right) D_x + h[v_{[k]}], \end{aligned}$$

from which it follows that the integrals can be represented in the following form, expressing the dependence of the integrals on the highest order derivatives

$$\begin{aligned} I_1 &= \frac{v_{1,[m]}}{v_{1,x} + v_1^2 - 1} + \dots \\ I_2 &= \frac{2v_{1,[m-1]}}{v_{1,x} + v_1^2 - 1} + \frac{v_{2,[m-1]}}{v_{2,x} + v_2^2 - 1} + \dots \\ &\dots\dots\dots \\ I_{m-1} &= \frac{(m-1)v_{1,xx}}{v_{1,x} + v_1^2 - 1} + \frac{(m-2)v_{2,xx}}{v_{2,x} + v_2^2 - 1} + \dots + \frac{v_{m-1,xx}}{v_{m-1,x}} + \dots \end{aligned}$$

Recall that $I_m = J$ is given explicitly by the equality (5.17). Now to complete the proof of the theorem we have to note that the determinant of the matrix (5.21) is equal to the expression:

$$\frac{e^{-(\epsilon+\delta)y}}{(v_1 - \epsilon)(v_2 - v_1)(v_3 - v_2) \dots (v_m - v_{m-1})(v_m - \delta)}.$$

Below we present an illustrative example assuming $m = 2$.

Example 2. For $m = 2$ the system (5.2) takes the form

$$\begin{aligned} v_{1,xy} &= \left(\frac{1}{v_1 - \epsilon} - \frac{1}{v_2 - v_1} \right) (v_{1,x} + v_1^2 - 1)(v_{1,y} + v_1^2 - 1) - 2v_1(v_{1,x} + v_{1,y} + v_1^2 - 1) \\ v_{2,xy} &= \left(\frac{1}{v_2 - v_1} - \frac{1}{\delta - v_2} \right) (v_{2,x} + v_2^2 - 1)(v_{2,y} + v_2^2 - 1) - 2v_2(v_{2,x} + v_{2,y} + v_2^2 - 1), \end{aligned} \quad (5.23)$$

The corresponding operator L is represented as follows

$$L = (D_x + r_2)(D_x + r_1) = D_x^2 + (r_1 + r_2)D_x + r_1r_2 + r_{1,x}, \quad (5.24)$$

where

$$r_1 = \frac{v_{1,x}}{v_1 - \epsilon} - B_1 = \frac{v_{1,x}}{v_1 - \epsilon} + \frac{v_{1,x}}{v_2 - v_1} - \frac{1 - v_1v_2}{v_2 - v_1} \quad (5.25)$$

and

$$r_2 = \left(\log \frac{v_1 - \epsilon}{A_1} \right)_x - B_2 = \frac{v_{1,x}}{v_1 - \epsilon} - \frac{v_{1,xx} + 2v_1v_{1,x}}{v_{1,x} + v_1^2 - 1} + \frac{v_{2,x} + v_2\delta - 1}{\delta - v_2} + \frac{v_{2,x} - v_{1,x}}{v_2 - v_1}. \quad (5.26)$$

The integral $\beta_1 = I$ is easily evaluated:

$$I = r_1 + r_2 = \frac{2v_{1,x}}{v_1 - \epsilon} - \frac{v_{1,xx} + 2v_1v_{1,x}}{v_{1,x} + v_1^2 - 1} + \frac{v_{2,x} + v_1v_2 - 1}{v_2 - v_1} + \frac{v_{2,x} + v_2\delta - 1}{\delta - v_2}.$$

Due to the Lemma 1 we have the integral $\beta_2 = J$ (evidently it is non-autonomous for $\epsilon \neq -\delta$):

$$J = \frac{(v_{1,x} + v_1^2 - 1)(v_{2,x} + v_2^2 - 1)}{(v_1 - \epsilon)(v_2 - v_1)(v_2 - \delta)} e^{-(\epsilon+\delta)y}. \quad (5.27)$$

These two functions I and J constitute the complete set of the y -integrals for the system (5.23).

6 On the reductions of the two-dimensional Volterra equation

It is well known that the two-dimensional Volterra equation

$$\begin{aligned} a_{n,y} &= a_n (b_n - b_{n+1}), \\ b_{n,x} &= b_n (a_{n-1} - a_n) \end{aligned} \quad (6.1)$$

admits a Lax pair of the following form (see, for instance, [4], [32]):

$$\psi_{n,x} = a_n (\psi_{n+1} - \psi_n), \quad \psi_{n,y} = b_n (\psi_n - \psi_{n-1}). \quad (6.2)$$

By imposing the boundary conditions $a_0 = 0$, $b_{m+1} = 0$ on the lattice (6.1), we arrive at a finite system of hyperbolic equations

$$\begin{aligned} a_{1,y} &= a_1 (b_1 - b_2), \\ a_{2,y} &= a_2 (b_2 - b_3), \\ &\dots\dots\dots \\ a_{m,y} &= a_m b_m, \\ b_{1,x} &= -b_1 a_1, \\ b_{2,x} &= b_2 (a_1 - a_2), \\ &\dots\dots\dots \\ b_{m,x} &= b_m (a_{m-1} - a_m), \end{aligned} \quad (6.3)$$

which is also integrable (see [8]). Its Lax pair can be easily derived from the linear system (6.2) by setting $\psi_0 = 0$, $\psi_{m+1} = 0$. Then the linear system in x takes the form:

$$\begin{aligned} \psi_{1,x} &= a_1 (\psi_2 - \psi_1), \\ \psi_{2,x} &= a_2 (\psi_3 - \psi_2), \\ &\dots\dots\dots \\ \psi_{m,x} &= -a_m \psi_m. \end{aligned} \quad (6.4)$$

In a similar way we get the linear system in y :

$$\begin{aligned} \psi_{1,y} &= b_1 \psi_1, \\ \psi_{2,y} &= b_2 (\psi_2 - \psi_1), \\ &\dots\dots\dots \\ \psi_{m,y} &= b_m (\psi_m - \psi_{m-1}). \end{aligned} \quad (6.5)$$

It is worth noting that the system (6.4), (6.5) exactly goes into the system (4.6)-(4.7), studied in section 4, if the relations are satisfied

$$a_n = \frac{u_{n,x}}{u_{n+1} - u_n}, \quad b_n = \frac{u_{n,y}}{u_n - u_{n-1}}, \quad (6.6)$$

as well as boundary conditions (4.1). The matter is that the relations (6.6) implement the Miura-type transformation between equations (E6) and (6.1). Let's rewrite some of the equations (6.3) in the following form:

$$\begin{aligned} \frac{a_{1,y}}{a_1} &= b_1 - b_2, \\ \frac{a_{2,y}}{a_2} &= b_2 - b_3, \\ &\dots\dots\dots \\ \frac{a_{m,y}}{a_m} &= b_m. \end{aligned} \quad (6.7)$$

Then, summing these equalities term by term, we obtain $D_y \log a_1 a_2 \dots a_m = b_1$. Further, keeping in mind that due to (4.1) and (6.6) the equality $b_1 = D_y \log(u_1 - C_0)$ holds we find $D_y \log(u_1 - C_0) = D_y \log a_1 a_2 \dots a_m$. Afterwards by integrating, we find

$$u_1 = C_0 + a_1 a_2 \dots a_m p(x), \quad (6.8)$$

where $p(x)$ is an arbitrary function.

Now, to construct the integrals of the system (6.3), one can use the scheme outlined in Section 4, where the key role is played by the operator B (4.13). To do this, in explicit representations (4.14) for the coefficients r_k it is necessary to exclude the variable u_1 due to (6.8). Simple calculations lead to the following expressions

$$r_k = a_k + \sum_{i=k}^m \frac{a_{i,x}}{a_i} + \frac{p'(x)}{p(x)}, \quad k = 1, 2, \dots, m. \quad (6.9)$$

For the coefficients of the operator (4.20) we find convenient representations in order to clarify their dependence on the higher order derivatives of the dynamical variables a_1, a_2, \dots, a_m . For example, for H_2 we have

$$\begin{aligned} H_2 &= \left(D_x + \sum_{i=2}^m \frac{a_{i,x}}{a_i} + h[a] \right) \left(D_x + \sum_{i=1}^m \frac{a_{i,x}}{a_i} + h[a] \right) = \\ &D_x^2 + \left(\sum_{i=2}^m \frac{a_{i,x}}{a_i} + \sum_{i=1}^m \frac{a_{i,x}}{a_i} + h[a] \right) D_x + \sum_{i=1}^m \frac{a_{i,xx}}{a_i} + h[a_x] \\ H_3 &= D_x^3 + \left(\sum_{i=3}^m \frac{a_{i,x}}{a_i} + \sum_{i=2}^m \frac{a_{i,x}}{a_i} + \sum_{i=1}^m \frac{a_{i,x}}{a_i} + h[a] \right) D_x^2 + \\ &\left(\sum_{i=2}^m \frac{a_{i,xx}}{a_i} + \sum_{i=1}^m \frac{a_{i,xx}}{a_i} + h[a_x] \right) D_x + \sum_{i=1}^m \frac{a_{i,xxx}}{a_i} + h[a_{xx}]. \end{aligned}$$

Here the symbols $h[a]$, $h[a_x]$ and $h[a_{xx}]$ indicate that the terms located at this place may depend on x , the dynamical variables a_1, a_2, \dots, a_m themselves and their derivatives with

respect to x of the order no higher than 2. By induction it can be shown that the operator H_m , coinciding with the operator B (4.13), admits the following representation:

$$\begin{aligned}
H_m = D_x^m + & \left(\frac{a_{m,x}}{a_m} + \sum_{i=m-1}^m \frac{a_{i,x}}{a_i} + \sum_{i=m-2}^m \frac{a_{i,x}}{a_i} + \dots + \sum_{i=1}^m \frac{a_{i,x}}{a_i} + h[a] \right) D_x^{m-1} + \\
& \left(\sum_{i=m-1}^m \frac{a_{i,xx}}{a_i} + \sum_{i=m-2}^m \frac{a_{i,xx}}{a_i} + \dots + \sum_{i=1}^m \frac{a_{i,xx}}{a_i} + h[a_x] \right) + \dots + \sum_{i=1}^m \frac{a_{i,[m]}}{a_i} + h[a_{[m-1]}],
\end{aligned} \tag{6.10}$$

where $a_{i,[s]} = \frac{\partial^s}{\partial x^s} a_i$, and the symbol $h[a_{[s]}]$ indicates that the terms located in this place depend on x , on the dynamical variables a_1, a_2, \dots, a_m and on their derivatives with respect to x of order no higher than s .

From the latter we find some special representation modulo the lower order derivatives for the coefficients of the operator $H_m = B$:

$$H_m = D_x^m + \beta_1 D_x^{m-1} + \beta_2 D_x^{m-2} + \dots + \beta_m,$$

which are y -integrals of the system of equations (6.3),

$$\begin{aligned}
\beta_1 &= \frac{a_{m,x}}{a_m} + \sum_{i=m-1}^m \frac{a_{i,x}}{a_i} + \dots + \sum_{i=1}^m \frac{a_{i,x}}{a_i} + \dots \\
\beta_2 &= \sum_{i=m-1}^m \frac{a_{i,xx}}{a_i} + \sum_{i=m-2}^m \frac{a_{i,xx}}{a_i} + \dots + \sum_{i=1}^m \frac{a_{i,xx}}{a_i} + \dots \\
&\dots\dots\dots \\
\beta_{m-1} &= \sum_{i=2}^m \frac{a_{i,[m-1]}}{a_i} + \sum_{i=1}^m \frac{a_{i,[m-1]}}{a_i} + \dots \\
\beta_m &= \sum_{i=1}^m \frac{a_{i,[m]}}{a_i} + \dots
\end{aligned} \tag{6.11}$$

Now we calculate the determinant of a matrix of the form (2.2) for the integrals $I_j = \beta_j$. Let's first consider a simple example, assuming $m = 2$. For $m = 2$ we have

$$\begin{aligned}
\frac{\partial \beta_1}{\partial a_{1,x}} &= \frac{1}{a_1}, \\
\frac{\partial \beta_1}{\partial a_{2,x}} &= \frac{2}{a_2}, \\
\frac{\partial \beta_2}{\partial a_{1,xx}} &= \frac{1}{a_1}, \\
\frac{\partial \beta_2}{\partial a_{2,xx}} &= \frac{1}{a_2}.
\end{aligned} \tag{6.12}$$

The required determinant has the form:

$$\begin{vmatrix} \frac{1}{a_1} & \frac{2}{a_2} \\ \frac{1}{a_1} & \frac{1}{a_2} \end{vmatrix} = -\frac{1}{a_1 a_2}.$$
 \tag{6.13}

Similarly, for $m = 3$ one can check that the determinant equals $-\frac{1}{a_1 a_2 a_3}$. In the case of an arbitrary m , the determinant of the form (2.2) coincides with the determinant of the

matrix

$$\left(\begin{array}{ccccccc} \frac{1}{a_1} & \frac{2}{a_2} & \frac{3}{a_3} & \cdots & \frac{m-2}{a_{m-2}} & \frac{m-1}{a_{m-1}} & \frac{m}{a_m} \\ \frac{1}{a_1} & \frac{2}{a_2} & \frac{3}{a_3} & \cdots & \frac{m-2}{a_{m-2}} & \frac{m-1}{a_{m-1}} & \frac{m-1}{a_m} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \frac{1}{a_1} & \frac{2}{a_2} & \frac{2}{a_3} & \cdots & \frac{2}{a_{m-2}} & \frac{2}{a_{m-1}} & \frac{2}{a_m} \\ \frac{1}{a_1} & \frac{a_2}{1} & \frac{a_3}{1} & \cdots & \frac{a_{m-2}}{1} & \frac{a_{m-1}}{1} & \frac{a_m}{1} \\ \frac{1}{a_1} & \frac{1}{a_2} & \frac{1}{a_3} & \cdots & \frac{1}{a_{m-2}} & \frac{1}{a_{m-1}} & \frac{1}{a_m} \end{array} \right) \quad (6.14)$$

By simple linear transformations, namely, subtracting the second row from the first one, the third row from the second one, etc., the matrix (6.14) can be reduced to a triangular form and make sure that its determinant is equal to $-\frac{1}{a_1 a_2 \dots a_m}$. Consequently, the set of y -integrals (6.11) forms a complete set of independent integrals.

Example 3. For $m = 2$ the system (6.3) takes the form

$$\begin{aligned} a_{1,y} &= a_1 (b_1 - b_2), \\ a_{2,y} &= a_2 b_2, \\ b_{1,x} &= -b_1 a_1, \\ b_{2,x} &= b_2 (a_1 - a_2). \end{aligned} \quad (6.15)$$

Let's find the complete set of the y -integrals for this system. In this case the functions r_1 and r_2 are given by

$$r_1 = a_1 + \frac{a_{1,x}}{a_1} + \frac{a_{2,x}}{a_2} + \frac{p'(x)}{p(x)}, \quad (6.16)$$

$$r_2 = a_2 + \frac{a_{2,x}}{a_2} + \frac{p'(x)}{p(x)}. \quad (6.17)$$

Without loss of generality we can put $p(x) = 1$. For the operator H_2 we have the following representation:

$$H_2 = D_x^2 + (r_1 + r_2)D_x + r_1 r_2 + r_{1,x}, \quad (6.18)$$

where functions $\beta_1 = r_1 + r_2$ and $\beta_2 = r_1 r_2 + r_{1,x}$ are the desired integrals. In virtue of the formulas (6.16), (6.17) we have the complete set of y -integrals for the system (6.15)

$$\begin{aligned} \beta_1 &= a_2 + a_1 + 2\frac{a_{2,x}}{a_2} + \frac{a_{1,x}}{a_1}, \\ \beta_2 &= \frac{a_{1,xx}}{a_1} + \frac{a_{2,xx}}{a_2} + a_1 a_2 + a_{2,x} + a_{1,x} + \frac{a_2 a_{1,x}}{a_1} + \frac{a_1 a_{2,x}}{a_2} + \frac{a_{2,x} a_{1,x}}{a_2 a_1} - \left(\frac{a_{1,x}}{a_1} \right)^2. \end{aligned}$$

References

- [1] A.V.Mikhailov, R.I.Yamilov, "On integrable two-dimensional generalizations of non-linear Schrödinger type equations", Physics Letters, Section A: General, Atomic and Solid State Physics, 230:5-6 (1997), 295-300.
- [2] D.J.Benney and G.J.Roskes, Wave instabilities, Stud. Appl. Math. 48 (1969) 377-385.
- [3] A.Davey and K.Stewartson, Proc. Roy. Soc. London A, 338, (1974) pp. 101-110.
- [4] A.B.Shabat, R.I.Yamilov, "To a transformation theory of two-dimensional integrable systems", Phys. Lett., A, 227:1-2 (1997), 15-23.
- [5] K.Ueno and K.Takasaki, Adv. Stud. Pure Math. 4 (1984) 1.
- [6] Laplace P S 1776 *Recherches sur le calcul intégral aux différences partielles* (Mémoires de l'académie royale de Sciences de Paris).

- [7] E.I.Ganzha and S.P.Tsarev 2001 *Integration of classical series A_n, B_n, C_n of exponential systems* (Krasnoyarsk: Krasnoyarsk State Pedagogical University Press).
- [8] A.N.Leznov, M.V.Saveliev, and V.G.Smirnov. "General solutions of the two-dimensional system of Volterra equations which realize the Bäcklund transformation for the Toda lattice." *Theoretical and Mathematical Physics* 47.2 (1981): 417-422.
- [9] V.Volterra, "Lecons sur la theorie Mathematique de la Lutte pour la Vie". Paris: Gauthier-Villars, 1931.
- [10] B.N.Breizman, V.E.Zakharov, S.L.Musher, "On kinetics of stimulated scattering of Langmuir waves by plasma ions"- *Zhurnal Eksperimental'noj i Teoreticheskoy Fiziki* 64 (4), 1297-1313, 1973.
- [11] V.E.Zakharov, S.L.Musher, A.M.Rubenchik, "Nonlinear stage of parametric wave excitation in a plasma".- *Jetp Lett* 19 (5), 151-152, 1974.
- [12] I.T.Habibullin, "Characteristic Lie rings, finitely-generated modules and integrability conditions for $(2+1)$ -dimensional lattices", *Physica Scripta*, 87:6 (2013), 065005, IOP Publishing.
- [13] M.N.Poptsova, I.T.Habibullin, "Algebraic properties of quasilinear two-dimensional lattices connected with integrability", *Ufa Math. J.*, 10:3 (2018), 86–105.
- [14] A.V.Zhiber, R.D.Murtazina, I.T.Habibullin, A.B.Shabat, "Characteristic Lie rings and nonlinear integrable equations", M.Izhevsk, 2012).
- [15] A.B.Shabat, R.I.Yamilov. "Exponential systems of type I and Cartan matrices" // Preprint. Bashkir Branch of Academy of Sciences of USSR. (1981).
- [16] I.T.Habibullin, M.N.Kuznetsova, "A classification algorithm for integrable two-dimensional lattices via Lie–Rinehart algebras", *Theoret. and Math. Phys.*, 203:1 (2020), 569–581.
- [17] M.N.Kuznetsova, "Classification of a subclass of quasilinear two-dimensional lattices by means of characteristic algebras", *Ufa Math. J.*, 11:3 (2019), 109–131.
- [18] E. V. Ferapontov, V. S. Novikov, and Ilia Roustemoglou. "On the classification of discrete Hirota-type equations in 3D." *International Mathematics Research Notices* 2015.13 (2015): 4933-4974.
- [19] E. V. Ferapontov, I. T. Habibullin, M. N. Kuznetsova and V. S. Novikov, "On a class of 2D integrable lattice equations". *J. Math. Phys.* 61, 073505 (2020); doi: 10.1063/5.0013697.
- [20] M.N.Kuznetsova, "Lax Pair for a Novel Two-Dimensional Lattice", *SIGMA*, 17 (2021), 088, 13 pp.
- [21] I.T.Habibullin, M.N.Poptsova, "Classification of a Subclass of Two-Dimensional Lattices via Characteristic Lie Rings", *SIGMA*, 13 (2017), 73 , 26 pp.
- [22] S.V.Smirnov, "Darboux integrability of discrete two-dimensional Toda lattices", *Theoret. and Math. Phys.*, 182:2 (2015), 189–210.
- [23] M.Gurses, A.V.Zhiber, I.T.Habibullin, "Characteristic Lie rings of differential equations", *Ufmsk. Mat. Zh.*, 4:1 (2012), 53–62.
- [24] D.K.Demskoi, "Integrals of open two-dimensional lattices", *Theoret. and Math. Phys.*, 163:1 (2010), 466–471.
- [25] I.M.Anderson and N.Kamran 1997 "The variational bicomplex for second order scalar partial differential equations in the plane" *Duke Math. J.* **87** 265-319.
- [26] A.V.Zhiber and V.V.Sokolov 2001 "Exactly integrable hyperbolic equations of Liouville type" *Russ. Math. Surv.* **56** 61-101.

- [27] A.M.Gurieva, A.V.Zhiber, “Laplace Invariants of Two-Dimensional Open Toda Lattices”, *Theoret. and Math. Phys.*, 138:3 (2004), 338–355.
- [28] I.T.Habibullin, A.R.Khakimova, “Characteristic Lie algebras of integrable differential-difference equations in 3D”, *Journal of Physics A: Mathematical and Theoretical*, 54:29 (2021), 295202 , 34 pp.
- [29] M. N. Kuznetsova, “Construction of localized particular solutions of chains with three independent variables”, *Theoret. and Math. Phys.*, 216:2 (2023), 1158–1167.
- [30] M. N. Kuznetsova, I. T. Habibullin, A. R. Khakimova, “On the problem of classifying integrable chains with three independent variables”, *Theoret. and Math. Phys.*, 215:2 (2023), 667–690.
- [31] A.V. Mikhailov, “Integrability of a Two-Dimensional Generalization of the Toda Chain”. *Jetp Lett.*, 30:7 (1979), 414–418.
- [32] E.V.Ferapontov, “Laplace transformations of hydrodynamic-type systems in Riemann invariants”, *Theoret. and Math. Phys.*, 110:1 (1997), 68–77.